

## Correcting models with long-range electron interaction using generalized cusp conditions

Andreas Savin<sup>\*1</sup> and Jacek Karwowski<sup>2</sup>

<sup>1</sup>*Laboratoire de Chimie Théorique, CNRS and Sorbonne University 4 place Jussieu, 75252 Paris cedex 05, France<sup>a)</sup>*

<sup>2</sup>*Institute of Physics, Faculty of Physics, Astronomy and Informatics, Nicolaus Copernicus University, Grudziadzka 5, 87-100 Toruń, Poland<sup>b)</sup>*

Sources of energy errors resulting from the replacement of the physical Coulomb interaction by its long-range  $\text{erfc}(\mu r)/r$  approximation are explored. It is demonstrated that the results can be dramatically improved and the range of  $\mu$  giving energies within chemical accuracy limits significantly extended, if the generalized cusp conditions are used to represent the wave function at small  $r$ . The numerical results for two-electron harmonium are presented and discussed.

Keywords: range separation; long-range interaction; short-range interaction, cusp conditions; Schrödinger equation; harmonium; chemical accuracy

---

<sup>a)</sup>Electronic mail: [andreas.savin@lct.jussieu.fr](mailto:andreas.savin@lct.jussieu.fr)

<sup>b)</sup>Electronic mail: [jka@umk.pl](mailto:jka@umk.pl)

## I. THE PROBLEM TO BE SOLVED

We have a model system,  $H(\mathbf{R}; \mu)$ , and a corresponding Schrödinger equation,

$$H(\mathbf{R}; \mu)\Psi(\mathbf{R}, \boldsymbol{\sigma}; \mu) = E(\mu)\Psi(\mathbf{R}, \boldsymbol{\sigma}; \mu). \quad (1)$$

The system is composed of  $N$  electrons confined by an external potential,  $\mathbf{R}$  and  $\boldsymbol{\sigma}$  stand, respectively, for their orbital and spin coordinates. All quantities characterizing the system (e.g. energy or wave function) depend on the external potential, but we show this dependence explicitly only when the form of this potential is specified (e.g. the dependence on  $\omega$  in the section "The model system").

The interaction between electrons is described by a  $\mu$ -dependent model potential  $v_{\text{int}}(r; \mu)$ :

- $\mu = 0$ : there is no interaction between electrons, so  $v_{\text{int}}(r; 0) = 0$ ,
- $\mu = \infty$ : we have the physical, Coulomb interaction, so  $v_{\text{int}}(r; \infty) = 1/r$ ,
- $\mu \in (0, \infty)$ : we choose

$$v_{\text{int}}(r; \mu) = w(r; \mu) = \frac{\text{erf}(\mu r)}{r}, \quad (2)$$

where  $r = r_{12} = |\mathbf{r}_1 - \mathbf{r}_2|$ . Exploring other forms of interaction may be both interesting and useful as, for example, in ref [1](#).

To simplify the notation, we drop  $\mu$  when  $\mu = \infty$ .

We assume that the solutions of eq (1) are accessible for selected finite values of  $\mu$ . However, we are not interested in the model system energy,  $E(\mu)$ . We aim at determining  $E$  corresponding to the physical interaction. Stated differently, we are interested in

$$\bar{E}(\mu) = E - E(\mu) \equiv -\Delta_0 E(\mu), \quad (3)$$

where  $\Delta_0 E(\mu)$  is referred to as the *error of the energy of the model system*.

To get an idea about the change of  $\bar{E}(\mu)$  with respect to  $\mu$ , we show in Figure [1](#) some situations where  $E$  is known for arbitrary precision. By construction,  $\bar{E}(\infty) = 0$ . As  $\mu$  decreases, the interaction weakens and disappears for  $\mu = 0$ . This effect, not compensated by any change in the external potential, leads to the absolute values of  $\bar{E}(\mu)$  increasing with decreasing  $\mu$  and becoming very large for sufficiently small  $\mu$ .

In this paper we explore how much one can lower the values of  $\mu$  and, by correcting the model, still retain approximations of  $\bar{E}$  within the *chemical accuracy* ( $\pm 1$  kcal/mol) error bars.




FIG. 1. Error of the energy of the model system,  $\Delta_0 E(\mu)$ , for the lowest energy state of harmonium with  $\omega = 1/2$  (red curves), and  $\omega = 1$  (thin blue curve), for  $\ell = 0$ , full curves;  $\ell = 1$ , dot-dashed curve;  $\ell = 2$ , dashed curve.

## II. CORRECTING MODELS

### A. Energy extrapolation – a historic solution

A way proposed in ref 2 is to expand  $\bar{E}(\mu)$  in the following basis,

$$\bar{E}(\mu) \approx \sum_{k=1}^M \bar{e}_k \chi_k(\mu). \quad (4)$$

Here  $\chi_k(\mu)$ ,  $k = 1, \dots, M$  are some basis functions, and  $\bar{e}_k$  are coefficients to be determined. There are different ways to determine these coefficients, once the basis functions are given. One option is to obtain them from the derivatives of  $E(\mu)$  with respect to  $\mu$ . This corresponds to (generalized) Taylor expansions, or to perturbation theory. Another possibility is calculating  $E(\mu)$  for several values of  $\mu$ , and then matching the results to the expansion. This method is called *energy extrapolation*<sup>2</sup>: from model information, we aim to reach the physical result.

One way to achieve our aim is to introduce more parameters into the Hamiltonian<sup>3</sup>, e.g.,

$$H(\mathbf{R}; \lambda, \mu) = H(\mathbf{R}; \mu) + \lambda [H(\mathbf{R}) - H(\mathbf{R}; \mu)]. \quad (5)$$

The eigenvalue and the corresponding eigenfunction of  $H(\mathbf{R}; \lambda, \mu)$  are, respectively,  $E(\lambda, \mu)$  and  $\Psi(\mathbf{R}, \boldsymbol{\sigma}; \lambda, \mu)$  – notice that  $E(\mu) = E(\lambda, \mu)|_{\lambda=0}$ . In the Hamiltonian (5) the interaction

potential is

$$v_{\text{int}}(r; \lambda, \mu) = w(r; \mu) + \lambda \bar{w}(r; \mu),$$

where

$$\bar{w}(r; \mu) = \frac{1}{r} - w(r; \mu) = \frac{1 - \text{erf}(\mu r)}{r} = \frac{\text{erfc}(\mu r)}{r}. \quad (6)$$

Therefore,

$$v_{\text{int}}(r; \lambda, \mu) = (1 - \lambda) \frac{\text{erf}(\mu r)}{r} + \frac{\lambda}{r} \quad (7)$$

We see that we can reach the physical result either with  $\lambda = 1$ , or with  $\mu = \infty$ . "Shooting" from different points to the same target may simplify our task. However, this is not further discussed in this paper.

Energy extrapolation has an important problem: we do not know how to choose  $\chi_k(\mu)$ . What makes the problem worse is that we are not willing to use many basis functions. Ideally, we should use a single function, that is to perform a single model calculation,  $M = 1$  in eq (4).

## B. The adiabatic connection

For  $\|\Psi(\mathbf{R}, \boldsymbol{\sigma}; \lambda, \mu)\| = 1$ , the Hellmann-Feynman theorem yields

$$\partial_\lambda E(\lambda, \mu) = \langle \Psi(\mathbf{R}, \boldsymbol{\sigma}; \lambda, \mu) | \partial_\lambda H(\mathbf{R}; \lambda, \mu) | \Psi(\mathbf{R}, \boldsymbol{\sigma}; \lambda, \mu) \rangle = \langle \bar{w}(\lambda, \mu) \rangle, \quad (8)$$

where

$$\langle \bar{w}(\lambda, \mu) \rangle = \langle \Psi(\mathbf{R}, \boldsymbol{\sigma}; \lambda, \mu) | \bar{W}(\mathbf{R}; \mu) | \Psi(\mathbf{R}, \boldsymbol{\sigma}; \lambda, \mu) \rangle, \quad (9)$$

and

$$\bar{W}(\mathbf{R}; \mu) \equiv H(\mathbf{R}) - H(\mathbf{R}; \mu) = \sum_{1 \leq i < j \leq N} \bar{w}(r_{ij}; \mu), \quad (10)$$

By integrating eq (8) over  $\lambda$  one obtains that <sup>1</sup>

$$\bar{E}(\mu) = \int_0^1 \langle \bar{w}(\lambda, \mu) \rangle d\lambda. \quad (11)$$

The integrand in eq (11), an integral over  $3N$ -dimensional configuration space and  $2^N$ -dimensional spin space, can be reduced to a one-dimensional radial integral. Exploiting

---

<sup>1</sup> Notice that  $E$  could have also been obtained by integration over  $\mu$ , as  $E = E(1, \mu) = E(\lambda, \infty)$ .

the antisymmetry of the wave function and integrating over spin, and over coordinates of electrons 3, 4,  $\dots$ ,  $N$ , yields<sup>4</sup>

$$\langle \bar{w}(\lambda, \mu) \rangle = \int_{\mathbb{R}^6} \bar{w}(r_{12}; \mu) \Gamma_{\lambda, \mu}(\mathbf{r}_1, \mathbf{r}_2) d\mathbf{r}_1 d\mathbf{r}_2, \quad (12)$$

where

$$\Gamma_{\lambda, \mu}(\mathbf{r}_1, \mathbf{r}_2) = \binom{N}{2} \sum_{\sigma_1, \dots, \sigma_N} \int_{\mathbb{R}^{3N-6}} |\Psi(\mathbf{R}, \boldsymbol{\sigma}; \lambda, \mu)|^2 d\mathbf{r}_3 d\mathbf{r}_4 \cdots d\mathbf{r}_N$$

is the diagonal part of the second-order reduced density matrix, 2RDM, corresponding to  $\Psi(\mathbf{R}, \boldsymbol{\sigma}; \lambda, \mu)$ ; the sum is extended over spin coordinates of all electrons.

After introducing the relative-motion variables

$$\mathbf{r} = \mathbf{r}_1 - \mathbf{r}_2, \quad \mathbf{r}^+ = \frac{\mathbf{r}_1 + \mathbf{r}_2}{2}, \quad (13)$$

performing integration over  $\mathbf{r}^+$ , and expressing  $\mathbf{r}$  in spherical coordinates,  $\mathbf{r}(r, \theta, \phi)$ , eq (12) can be rewritten as

$$\langle \bar{w}(\lambda, \mu) \rangle = \int_{\mathbb{R}^6} \bar{w}(r; \mu) \Gamma_{\lambda, \mu}(\mathbf{r}, \mathbf{r}^+) d\mathbf{r} d\mathbf{r}^+ = \int_{\mathbb{R}^3} \bar{w}(r; \mu) \gamma(\mathbf{r}; \lambda, \mu) d\mathbf{r}, \quad (14)$$

where  $d\mathbf{r} = r^2 dr \sin \theta d\theta d\phi$ ,  $r = r_{12} = |\mathbf{r}_1 - \mathbf{r}_2|$ , and  $\gamma(\mathbf{r}; \lambda, \mu)$  is the diagonal part of the first-order reduced density matrix, 1RDM. Since in the coordinate space  $\bar{w}(r; \mu)$  depends on the radial coordinate  $r$  only, we can do the spherical averaging. In effect the integrand of eq (11) is simplified to

$$\langle \bar{w}(\lambda, \mu) \rangle = \int_0^\infty \bar{w}(r; \mu) \tilde{\gamma}(r; \lambda, \mu) r^2 dr, \quad (15)$$

where

$$\tilde{\gamma}(r; \lambda, \mu) = \int_0^{2\pi} \int_0^\pi \gamma(\mathbf{r}; \lambda, \mu) \sin \theta d\theta d\phi.$$

The adiabatic connection defined in eq (11) carries no practical information, as it requires the knowledge of  $\langle \bar{w}(\lambda, \mu) \rangle$  for all values of  $\lambda$  while in the present approach it is known only for  $\lambda = 0$ .

At the limit of  $r \rightarrow 0$ ,  $\bar{w}(r; \mu) \sim 1/r$ . Therefore, also for large  $\mu$ ,  $\bar{w}(r; \mu)$  is non-negligible if  $r$  is small enough. As is shown hereafter, the information necessary for correcting the model at small  $r$  can be derived from the generalized cusp conditions (GCC).

### C. Generalized cusp conditions

The information about the behavior of the wave function in the vicinity of the coalescence point, i.e., for  $r = r_{12} \ll 1$ , can be derived from general properties of the Schrödinger equation at  $r \rightarrow 0$ . In general, the approach is based on the expansion of the wave function and of the potential as the power series of  $r$  and deriving conditions that have to be fulfilled by the expansion coefficients in order to retain the consistency of the Hamiltonian eigenvalue problem. The simplest and most commonly known is the Kato's cusp condition<sup>5</sup>, which can be derived from the requirement that in the case of the electrostatic interaction the local energy at  $r = 0$  is nonsingular. Higher order (generalized) cusp conditions can be obtained from the demand that the local energies generated by powers of the Hamiltonian are nonsingular (energy-independent conditions) and ratios of the local energies generated by the consecutive powers are constant (energy-dependent conditions)<sup>6</sup>. Alternatively one can use the expansion of the wave function in powers of  $r$  and require that the Schrödinger equation is satisfied<sup>7-9</sup>. Both approaches are equivalent, but the conditions derived from the former one, though more complicated, have more transparent physical meaning.

In this paper the GCCs are applied to describe the  $r$ -dependence of the wave function in the area of small  $r$ , where the model interaction potential departs from the physical one.

### III. THE MODEL SYSTEM

The simplest nontrivial model system containing one pair of electrons is composed of three particles: two electrons interacting by a repulsive model potential, and a third particle, "nucleus", which interacts with electrons by an attractive force. Commonly known examples of such systems are helium atom - the nucleus attracts electrons by the Coulomb force, and harmonium (Hooke atom) - the nucleus attracts electrons by the Hooke force. After the separation of the center of mass, the system is reduced to two interacting particles in an external potential  $v_{\text{ext}}$ . In the case of harmonium,

$$v_{\text{ext}}(r_1, r_2; \omega) = \frac{\omega^2}{2} (r_1^2 + r_2^2). \quad (16)$$

The potential depends on a parameter,  $\omega$ , which defines the strength of the confinement. In the case of quantities that depend on this potential, the dependence on  $\omega$  is explicitly

shown. For example,  $E(\omega, \mu)$  stands for the special case of  $E(\mu)$ , corresponding to the external potential (16).

To our knowledge, harmonium is the only bound system containing a pair of interacting electrons for which the Schrödinger equation is known to be separable. Apart from the three-dimensional free-particle equation describing the motion of the center of mass, the two-particle Schrödinger equation for harmonium is separable into six one-dimensional equations - five are exactly solvable, and the sixth one can be solved numerically to an arbitrary precision (for some specific values of  $\omega$  it is also solvable analytically). It is important to note that the separability holds for all forms of the interaction potential including the form given by  $v_{\text{int}}(r; \lambda, \mu)$ .<sup>2</sup> Therefore, harmonium is particularly suitable for pilot studies of the consequences of using various non-Coulombic forms of the interaction potentials. Motivated by these observations, at this stage, we explore our problem using harmonium as the model system.

The Schrödinger equation for harmonium

$$[T(\mathbf{r}_1, \mathbf{r}_2) + v_{\text{ext}}(r_1, r_2; \omega) + v_{\text{int}}(r; \lambda, \mu) - \mathcal{E}(\mathbf{p})] \Psi(\mathbf{r}_1, \mathbf{r}_2; \mathbf{p}) = 0, \quad (17)$$

where  $T(\mathbf{r}_1, \mathbf{r}_2)$  is two-particle kinetic energy operator, depends on three parameters, collectively denoted  $\mathbf{p} = \{\omega, \lambda, \mu\}$ , and  $\Psi(\mathbf{r}_1, \mathbf{r}_2; \mathbf{p})$  is the orbital part of the two-electron wave function. After transformation (13), eq (17) can be split into two spherically-symmetric equations. The first one depends on the interaction potential and describes the relative motion of electrons:

$$[-\Delta_{\mathbf{r}} + v(r; \mathbf{p}) - E(\mathbf{p})] \psi_{\text{rel}}(\mathbf{r}; \mathbf{p}) = 0, \quad (18)$$

where

$$v(r; \mathbf{p}) = \frac{\omega^2 r^2}{4} + v_{\text{int}}(r; \lambda, \mu) = \frac{\omega^2 r^2}{4} + (1 - \lambda) \frac{\text{erf}(\mu r)}{r} + \frac{\lambda}{r}. \quad (19)$$

The second equation describes the motion of the center of mass of the electron pair in the external potential:

$$\left[ -\frac{\Delta_{\mathbf{r}^+}}{4} + \omega^2 (r^+)^2 - \mathfrak{E}(\omega) \right] \psi_{\text{cm}}(\mathbf{r}^+; \omega) = 0, \quad (20)$$

where  $r^+ = |\mathbf{r}^+|$ . By construction, we have

$$\mathcal{E}(\mathbf{p}) = E(\mathbf{p}) + \mathfrak{E}(\omega), \quad \Psi(\mathbf{r}_1, \mathbf{r}_2; \mathbf{p}) = \psi_{\text{rel}}(\mathbf{r}; \mathbf{p}) \psi_{\text{cm}}(\mathbf{r}^+; \omega). \quad (21)$$

---

<sup>2</sup> In principle the term 'harmonium' refers to two confined Coulomb-interacting electrons. In this paper we extend this term to the case of model potentials  $v_{\text{int}}$ .

The interaction potential appears only in eq (18). So, for our study, we deal with this equation only. The potential is spherically symmetric. Therefore,

$$\psi_{\text{rel}}(\mathbf{r}; \mathbf{p}) = \psi_\ell(r; \mathbf{p}) Y_{\ell m}(\theta, \phi), \quad (22)$$

where  $\psi_\ell(r; \mathbf{p})$  is determined by the radial equation

$$\left[ -\frac{d^2}{dr^2} + \frac{\ell(\ell+1)}{r^2} + v(r; \mathbf{p}) - E(\mathbf{p}) \right] [r \psi_\ell(r; \mathbf{p})] = 0. \quad (23)$$

The two-electron wave function,  $\Psi(\mathbf{r}_1, \mathbf{r}_2; \mathbf{p})$ , symmetric/antisymmetric with respect to the transposition of  $(\mathbf{r}_1, \mathbf{r}_2)$  correspond to singlet/triplet. As one can see, singlet states correspond to the even parity (even  $\ell$ ) spherical harmonics in eq (22), and triplet states - to the odd ones.

### A. Generalized cusp conditions for the model system

Using<sup>10</sup>

$$\frac{\text{erf}(\mu r)}{r} = \frac{2\mu}{\sqrt{\pi}} \left( 1 - \frac{(\mu r)^2}{3 \cdot 1!} + \frac{(\mu r)^4}{5 \cdot 2!} - \frac{(\mu r)^6}{7 \cdot 3!} + \dots \right)$$

one can expand the potential (19) as

$$v(r; \mathbf{p}) = \sum_{i=-1}^{\infty} v_i(\mathbf{p}) r^i, \quad (24)$$

with

$$v_{-1} = \lambda, \quad v_0 = (1 - \lambda) \frac{2\mu}{\sqrt{\pi}}, \quad v_1 = 0, \quad v_2 = \frac{\omega^2}{4} - (1 - \lambda) \frac{2\mu^3}{3\sqrt{\pi}}, \quad v_3 = 0, \dots \quad (25)$$

The wave function, for small  $r$ , can be represented by the following power series

$$\psi_\ell(r; \mathbf{p}) \approx r^\ell \sum_{k=0}^K c_k(\mathbf{p}) r^k = c_0(\mathbf{p}) r^\ell \sum_{k=0}^K \tilde{c}_k(\mathbf{p}) r^k, \quad (26)$$

where  $\tilde{c}_k(\mathbf{p}) = c_k(\mathbf{p})/c_0(\mathbf{p})$ . General formulas for GCC are given in refs 6–9. Here we give equations defining  $\tilde{c}_k$  for  $k \leq 4$ :

$$\begin{aligned} A_1 \tilde{c}_1 + v_{-1} &= 0, \\ A_2 \tilde{c}_2 + v_{-1} \tilde{c}_1 - \epsilon &= 0, \\ A_1 A_3 \tilde{c}_3 + (A_1 + A_2) v_{-1} \tilde{c}_2 + v_{-1}^2 \tilde{c}_1 + A_1 v_1 &= 0, \\ A_2 A_4 \tilde{c}_4 + (A_2 + A_3) v_{-1} \tilde{c}_3 + v_{-1}^2 \tilde{c}_2 + A_2 v_1 \tilde{c}_1 + (v_{-1} v_1 + A_2 v_2 - \epsilon^2) &= 0, \\ &\dots \dots \dots \end{aligned} \quad (27)$$



where  $A_i = -i(2\ell+i+1)$ , and  $\epsilon = E - v_0$ . Coefficients  $\tilde{c}_k$ , energy parameter  $\epsilon$ , and coefficients  $v_i$ , depend on the parameters  $\omega$ ,  $\lambda$  and  $\mu$ . For simplicity, in eqs (27) this dependence has not been shown explicitly. All coefficients  $\tilde{c}_k$  depend on  $\ell$  and on  $v_{-1}$ . The coefficients  $\tilde{c}_k$  with  $k \geq 3$  depend on  $v_1$ . In general,  $v_i$  shows up in  $\tilde{c}_k$  with  $k \geq (i+2)$ <sup>6</sup>. The external potential is proportional to  $r^2$  and vanishes at  $r = 0$ . Therefore, in expansion (26) the  $\omega$  dependence begins at  $\tilde{c}_4$ .

For the construction of  $\tilde{c}_k$  with even values of  $k$ , the state energy is needed. In these cases we use the expectation value of the Hamiltonian defined in eq (5):

$$E(\omega, \lambda, \mu) \approx E(\omega, \mu) + \lambda \langle \bar{w}(r; \omega, 0, \mu) \rangle,$$

where  $E(\omega, \mu) = E(\omega, \lambda, \mu)|_{\lambda=0}$ . For definition of  $\langle \bar{w}(r; \omega, \lambda, \mu) \rangle$  see eq (28). Numerical tests using the exact  $E(\omega, \lambda, \mu)$  have shown that this approximation is negligible in comparison to the other approximations made in the present paper.

## B. Dependence on $\lambda$ and normalization

The GCC provide only ratios  $c_k(\mathbf{p})/c_0(\mathbf{p}) \equiv \tilde{c}_k(\mathbf{p})$ . If  $\psi_\ell(r; \mathbf{p})$  is known in the whole range of  $r$  then  $c_0(\mathbf{p})$  can be determined by the normalization condition. In our case this approach is nonapplicable since only the small- $r$  part of  $\psi_\ell(r; \mathbf{p})$  is defined by the cusp conditions. But  $c_0(\mathbf{p})$  appears as a prefactor in the approximation (26) for  $\psi_\ell(r; \mathbf{p})$ , and its value is necessary for any practical use of this approximation. Therefore,  $c_0(\mathbf{p})$  has to be estimated in a different way using only the information about the short-range behavior of the wave function.

We have to introduce additional information to deal with this issue. Let us first consider the dependence of  $c_0$  on  $\lambda$ . It is needed for the adiabatic connection expression [eqs (11) and (15)]. We select  $\|\psi_{\text{cm}}(\mathbf{r}^+; \omega)\| = 1$ . Then the substitution of the wave function defined in eqs (21) and (22), and of its expansion (26), to eq (14) yields

$$\begin{aligned} \langle \bar{w}(\mathbf{p}) \rangle &= \int_0^{2\pi} \int_0^\pi \int_0^\infty |\psi_{\text{rel}}(\mathbf{r}; \mathbf{p})|^2 \bar{w}(r; \mu) r^2 dr \sin \theta d\theta d\phi \\ &= \int_0^\infty |\psi_\ell(r; \mathbf{p})|^2 \bar{w}(r; \mu) r^2 dr, \\ &\approx |c_0(\mathbf{p})|^2 \int_0^\infty \left| r^{\ell+1} \sum_{k=0}^K \tilde{c}_k(\mathbf{p}) r^k \right|^2 \bar{w}(r; \mu) dr, \end{aligned} \tag{28}$$

and, according to eq (11),

$$\bar{E}(\omega, \mu) = \int_0^1 \langle \bar{w}(\omega, \lambda, \mu) \rangle d\lambda. \quad (29)$$

For models close to the exact interaction (for large  $\mu$ ) one can derive the following relationship<sup>3,11</sup> (a derivation is given in the Appendix)

$$c_0(\omega, \lambda, \mu) \underset{\mu \rightarrow \infty}{\approx} \mathcal{N} \left( 1 + \frac{1 - \lambda}{\sqrt{\pi}\mu} + O(\mu^{-2}) \right), \quad (30)$$

where  $\mathcal{N}$  is a still unknown normalization constant. As one can see,

$$\mathcal{N} = c_0(\omega, 1, \mu) = c_0(\omega, \lambda, \infty) \equiv c_0(\omega).$$

Notice that  $c_0(\omega, 1, \mu)$  and  $c_0(\omega, \lambda, \infty)$  do not depend, respectively, on  $\mu$  and on  $\lambda$  and are equal to  $c_0(\omega)$  corresponding to the physical (Coulomb) interaction potential.

We introduce the notation

$$\mathcal{I}_K(\omega, \lambda, \mu) = \left( 1 + \frac{1 - \lambda}{\sqrt{\pi}\mu} \right)^2 \int_0^\infty \left[ r^{\ell+1} \sum_{k=0}^K \tilde{c}_k(\omega, \lambda, \mu) r^k \right]^2 \bar{w}(r, \mu) dr. \quad (31)$$

As stated at the beginning of this article, we know the model wave function at  $\lambda = 0$ . Therefore, we can calculate

$$\langle \bar{w}(\omega, 0, \mu) \rangle = \int_0^\infty |\psi(r, \omega, 0, \mu)|^2 \bar{w}(r, \mu) r^2 dr. \quad (32)$$

Since  $\mathcal{N}$  does not depend on  $\lambda$ , we can assume that

$$\mathcal{N}^2 \approx \frac{\langle \bar{w}(\omega, 0, \mu) \rangle}{\mathcal{I}_K(\omega, 0, \mu)}. \quad (33)$$

Combining eqs (29), (28), and (33) we get

$$\bar{E}(\omega, \mu) \approx \frac{\langle \bar{w}(\omega, 0, \mu) \rangle}{\mathcal{I}_K(\omega, 0, \mu)} \int_0^1 \mathcal{I}(\omega, \lambda, \mu) d\lambda \quad (34)$$

### C. Corrections to the model

Assume that  $\omega$  is fixed and  $\lambda = 0$ . For simplicity, in the next equations, we skip these parameters. Also the  $r$ -dependence, in most cases obvious, is not shown explicitly. We designate  $\mu$ -dependence only, retaining the convention that we drop  $\mu$ , if  $\mu = \infty$ . Our model is defined by the analog of eq (1):

$$H(\mu)\psi(\mu) = E(\mu)\psi(\mu), \quad \text{i.e.} \quad E(\mu) = \langle \psi(\mu) | H(\mu) | \psi(\mu) \rangle. \quad (35)$$

For the physical (Coulombic) interaction,  $\mu = \infty$  and we have  $E = \langle \psi | H | \psi \rangle$ .

As we already said, the solutions of the crude model,  $E(\mu)$  and  $\psi(\mu)$ , are known. The error of this model,  $\Delta_0 E(\mu)$ , has been defined in eq (3). Our aim is to add corrections to the model. The first-order perturbation correction to  $E(\mu)$  is

$$\langle \psi(\mu) | H - H(\mu) | \psi(\mu) \rangle = \langle \psi(\mu) | \bar{w}(\mu) | \psi(\mu) \rangle. \quad (36)$$

From here we have the correction:

$$\Delta_H E(\mu) = \langle \psi(\mu) | H | \psi(\mu) \rangle - E = E(\mu) - E + \langle \bar{w}(\mu) \rangle. \quad (37)$$




FIG. 2. Errors of the model energy,  $\Delta_0 E(\mu)$ , eq (3), gray curve, and of the expectation value of the physical Hamiltonian,  $H$ , with the model wave function,  $\psi(\mu)$ ,  $\Delta_H E(\mu)$ , eq (37), black curve, for the ground state of harmonium with  $\omega = 0.5$ . The inset zooms on the same curves (by two orders of magnitude), the horizontal dashed lines indicating the region of "chemical accuracy" ( $\pm 1$  kcal/mol).

Plots of  $\Delta_0 E(\mu)$  and  $\Delta_H E(\mu)$  are given in Figure 2. At  $\mu = 0$ , that is for the non-interacting system,  $\Delta_0 E(\mu)$  is huge. For the ground state of harmonium with  $\omega = 1/2$ ,  $E(\mu = 0) = 0.75$  instead of  $E = 1.25$  hartree, giving  $\Delta_0 E(\mu) = 0.5$  hartree. But,  $\Delta_0 E(\mu)$

does not fall to the chemical accuracy error bars in the entire range of  $\mu$  – up to 3 bohr<sup>-1</sup>. The improvement due to the first-order perturbation (without any reference to a specific structure of the wave function) is impressive. For  $\Delta_{\text{H}}E(\mu)$  the chemical accuracy is reached if  $\mu > 1.5$  bohr<sup>-1</sup>. The error at  $\mu = 0$  is reduced from 0.5 hartree to  $\sim 0.06$  hartree, that is of the same order of magnitude as obtained with mean-field approximations ( $\sim 0.04$  hartree for Hartree-Fock or Kohn-Sham). The sign of the errors in Figure 2 is (i) negative for  $E(\mu) - E$  (as  $w(r, \mu) \leq 1/r$ ), and (ii) positive for  $\langle \psi(\mu) | H | \psi(\mu) \rangle - E$  (by the variational principle).

Eq (34) may be rewritten as

$$\Delta_{\text{K}}E(\mu) = E(\mu) - E + C_{\mathcal{I}}^K \langle \bar{w}(\mu) \rangle, \quad (38)$$

where

$$C_{\mathcal{I}}^K = \frac{1}{\mathcal{I}_K(\omega, 0, \mu)} \int_0^1 \mathcal{I}(\omega, \lambda, \mu) d\lambda. \quad (39)$$

Equation (38) can be interpreted as a generalization of eqs (3) and (37). The prefactor of  $\langle \bar{w}(\mu) \rangle$  in these two equations is, respectively, 0 and 1. Prefactor  $C_{\mathcal{I}}^K$  in eq (38) has been derived using some information about the structure of the wave function, specifically, GCC and the adiabatic connection. Therefore, one can expect that

$$|\Delta_0 E(\mu)| \gg |\Delta_{\text{H}}E(\mu)| \gg |\Delta_{\text{K}}E(\mu)|. \quad (40)$$

The correctness of this expectation is demonstrated in the next section.

The integrals over  $r$  in eq (31) can be computed analytically<sup>10</sup>

$$\int_0^\infty \text{erfc}(\mu r) r^k dr = \frac{k \Gamma(k/2)}{2\sqrt{\pi}(k+1)\mu^{k+1}}. \quad (41)$$

Therefore  $\mathcal{I}_K(\omega, \lambda, \mu)$  can be expressed as an algebraic function of  $\omega$ ,  $\lambda$ , and  $\mu$ . Explicit formulas for  $\tilde{c}_k(\mathbf{p})$  can be easily deduced from eq (27). Consequently, the effort for computing the prefactor  $C_{\mathcal{I}}^K$  in eq (38) is negligible. Notice that in eq (41))  $\Gamma(k/2)$  introduces a fast increase of the absolute value of  $\mathcal{I}_K(\omega, \lambda, \mu)$  with  $K$  (the maximum power of  $r$  in the expansion).

The approximation is valid only asymptotically, for sufficiently large  $\mu$ . Notice that expressions for  $c_0(\mathbf{p})$  and  $\mathcal{I}(\mathbf{p})$  given in eqs (30), and (31), are divergent at  $\mu = 0$ .

## IV. RESULTS

### A. General considerations

In the following we will consider only harmonium systems. Of course, this can be seen as futile, because solving the one-dimensional differential Schrödinger equation, eq (23), is trivial (to obtain accurate results we did it on a grid of the order of  $10^5$  points). The choice of this simple systems is motivated by the desire to study uniquely the effect of the approximations introduced without adding other effects such as the dependence on the one-particle basis set, or the expansion in terms of Slater determinants. To avoid introducing new effects, the one-particle (external) potential is independent of  $\mu$ . In view of future practical applications, we are interested to see how weak the model interaction can be taken (how small  $\mu$  can be chosen), and still obtain an estimate of the energy within "chemical accuracy".

We now define the "smallest acceptable  $\mu$ ",  $SA\mu$ : for  $\mu \geq SA\mu$  the absolute errors are smaller than the chemical accuracy (1 kcal/mol). The range of errors between  $\pm 1$  kcal/mol is indicated in the plots by horizontal dashed lines. For the system considered in Figure 2,  $E(\mu)$  has a  $SA\mu$  slightly above  $3 \text{ bohr}^{-1}$ . So, it is not interesting to discuss corrections if the interaction already reaches this strength. Weaker interactions can be considered if we simply correct the energy to first order – with  $\langle \psi(\mu) | H | \psi(\mu) \rangle$ ,  $SA\mu \approx 1.5$ .

In the following, the presented plots show the errors of the approximations of  $E$  as functions of  $\mu$  using the same scale as the inset in Figure 2.

### B. How weak can the interaction be?

Let us consider the ground state of harmonium with  $\omega = 1/2$ . Now we use the GCC to correct the model energy,  $E(\mu)$  as in eqs (34) and (38), and cut off the expansion of the wave function  $\psi(r, \mathbf{p})$ , limiting the expansion in eq (26) to a maximal power of  $r$ ,  $K = 1, 2, 3, 4$  – see Figure 3a. When  $K = 1$  we satisfy only Kato's cusp condition. This improves over  $E(\mu)$ , and also over first-order perturbation theory, bringing the  $SA\mu$  to  $1.3 \text{ bohr}^{-1}$ . Increasing  $K$  further reduces the  $SA\mu$ , until a "wall" at around  $0.5 \text{ bohr}^{-1}$  is reached. Notice that not only the error is reduced by increasing  $K$ , but also the stability with respect to the change of the results by changing  $\mu$  is increased when going beyond satisfying only the Kato cusp




FIG. 3. Errors of the different approximations for the energy of harmonium systems. The errors of the model energy,  $\Delta_0 E(\mu)$ , eq (3) are represented as gray curves, that of the expectation value of the physical Hamiltonian with the model wave function,  $\Delta_H E(\mu)$ , eq (37), by black curves. Panels a – g show the errors for different orders  $K$  in the GCC expansion, eq (26), the value of  $K$  being indicated next to the curve. Panel h shows the error of the approximation of order  $\mu^{-4}$ , as given in eq (42). Chemical accuracy ( $\pm 1$  kcal/mol) is indicated by horizontal dashed lines. a:  $\omega = 1/2, n = 1, \ell = 0$ ; b:  $\omega = 1/2, n = 1, \ell = 1$ ; c:  $\omega = 1/2, n = 1, \ell = 2$ ; d:  $\omega = 1/2, n = 2, \ell = 0$ ; e:  $\omega = 1/2, n = 3, \ell = 0$ ; f:  $\omega = 1, n = 1, \ell = 0$ ; g:  $\omega = 1/4, n = 1, \ell = 0$ ; h:  $\omega = 1/4, 1/2$ , or  $1$ , and  $n = 1, \ell = 0$ .




FIG. 4. Errors in the ground state energy estimate for harmonium ( $\omega = 1/2$ ) using density functional approximations ( $\mu$ -LDA and  $\mu$ -PBE). Also shown are the errors of the model energy,  $\Delta_0 E(\mu)$ , gray curve, and of the expectation value of the physical Hamiltonian,  $\Delta_H E(\mu)$ , black curve.

condition. This is important because (i) in practice, there is an arbitrariness in the choice of  $\mu$ , and (ii) the SA $\mu$  is system-dependent, as will be illustrated further down. The latter is of importance for size-consistency.

The derivations presented above never supposed that the state considered corresponds to the ground state. So, let us now consider some excited states. We consider first the lowest energy states with  $\ell = 1$  and  $\ell = 2$ . The first corresponds to a triplet state, (Figure 3b), the second to a non-natural singlet state<sup>12</sup> (Figure 3c); for  $\ell = 2$  the singlet cusp condition does not give a prefactor  $(1 + r/2)$  but  $(1 + r/6)$ . King<sup>13</sup> remarks that, in an orbital picture, the  $\ell = 2$  state corresponds to a strong mixture of sd and  $p^2$  configurations. Notice that both the values provided by the model, and the expectation value of the physical Hamiltonian, are now in much better agreement with the exact value than for  $\ell = 0$ : the prefactor  $r^\ell$  appearing in eq (26) already keeps the electrons apart. Even for  $K = 4$ , for this system and these states, there is no significant gain over using just  $\langle \psi(\mu) | H | \psi(\mu) \rangle$ . For  $\ell = 2$ , the SA $\mu$ s

obtained for  $K \leq 4$  are not improved over that of the expectation value of the Hamiltonian.

We also present the first two excited states with  $\ell = 0$ , Figure 3d,e. The model system has much larger errors which fall outside the domain of the plots. We notice an overall worsening of the quality of the approximations. The model shows errors that fall outside the range of the plots. The SA $\mu$  for the expectation value of the Hamiltonian is around 2 bohr<sup>-1</sup>. Using the Kato cusp condition reduces it to  $\approx 1.5$  bohr<sup>-1</sup>. Increasing the order of the expansion,  $K$ , moves the SA $\mu$  down to somewhere between 1 and 0.5 bohr<sup>-1</sup>.

We have seen above that the SA $\mu$ s can be quite sensitive to the state described. They can also be sensitive to the system. Modifying  $\omega$  affects the SA $\mu$ s. Figure 3f,g show the effect of making the system more compact ( $\omega = 1$  au), or diffuse ( $\omega = 1/4$  au), for the respective ground states.

### C. Comparisons

Using the GCC is not the only way to describe the short-range behavior. One can replace the expansion in powers of  $r$ , eq (26) by a simple form that does not diverge as  $r \rightarrow \infty$ . For example, one can ignore all terms arising from the external potential and the energy in the Schrödinger equation, in the limit  $\mu \rightarrow \infty$ , and we get, to order  $\mu^{-4}$  (see ref 11, and eq (21) in ref 3),

$$\frac{\int_0^1 d\lambda \mathcal{I}(\lambda, \mu)}{\mathcal{I}(\lambda = 0, \mu)} = \frac{\mu^2 + 1.06385\mu + 0.31982}{\mu^2 + 1.37544\mu + 0.487806} \quad (42)$$

A comparison of Figures 3a, 3f, or 3g to 3h shows that this approximation performs very well compared to those obtained from the GCC: the "walls" are at comparable values of  $\mu$ . This is encouraging because the GCC require, in general, taking into account the external potential and the energy.

Density functional approximations have been used for many years to correct models for missing short-range interaction.<sup>14,15</sup> Figure 4 shows the corrections provided for two approximations, the local density approximation, LDA, and that of Perdew, Burke, and Ernzerhof, PBE, modified to depend on  $\mu$ .<sup>15-17</sup> The SA $\mu$ s are around 1 bohr<sup>-1</sup>, thus improving over using only Kato's cusp condition (cf. Figure 3a). It may be at first surprising that in Figure 4 LDA is slightly better than PBE. However, PBE becomes better than LDA for small  $\mu$ . This cannot be seen in Figure 4 because it shows only the region of "chemical accuracy"; the errors for small  $\mu$  are much larger.



## V. CONCLUSIONS

### A. Summary

We have considered model systems, where electrons interact only via a long-range potential, eq (2). In order to obtain the physical energy we explored corrections based upon the short-range behavior of the wave function, both for the ground and excited states. The numerical results were all obtained for harmonium, eqs (16) – (22) where accurate results are easy to reach due to separability.

We are interested in having corrections to models where the interaction is very weak. However, the approximations proposed can be systematically improved in an asymptotic sense: as the order of the approximation increases, there is a domain of models close enough to the exact solution that gets systematically better. Unfortunately, at present, the approximations fail when the range of the interaction between particles becomes too large (our parameter  $\mu$  becomes too small). We attribute it to imposing only the short-range behavior of the wave function. For short range, the results turned out to be more reliable than obtained by correcting the models with density functional approximations. Unfortunately, the range of validity is system- and state-dependent.

### B. Perspectives

In quantum chemistry there are different approaches to tackle the problem raised by the singularity of the Coulomb potential. A “brute force” pathway, is to use a large expansion in Slater determinants. The burden can be reduced significantly by using “selected configuration interaction” techniques (see, e.g., ref 20 and references therein). Another way is to improve the description of the wave function by the use of correlation factors, like Jastrow factors in Quantum Monte Carlo (see, e.g., refs 21–24), or F12 methods (see, e.g., refs 25,26). Still another way to approach this problem is to use density functionals that transfer the short-range behavior from other systems (as a rule, from the uniform electron gas).

In this paper, we combine the spirit of the last two approaches. As in density functional calculations, we compute a model system for which the interaction has no singularity (and thus is expected to converge faster, in general). However, in contrast to density functional calculations, no density functional approximations are present, and the Hohenberg-Kohn

theorem (formulated for ground states) is not used. The approach is applicable to ground and excited states.

As in methods using correlation factors we use the exact short-range behavior of the wave function. The trick allowing to use it comes from having a correction that depends only on the missing part of the interaction that is short-ranged.

The present paper did only show exploratory calculations. However, it is possible to extend considerations presented in this paper to other systems. Already Kurokawa *et al.*<sup>8</sup> have shown that the GCC can be applied when the Schrödinger equation for the relative coordinate cannot be separated from that of the center of mass (the He atom).

Before extending our approach there are several issues to be explored. Probably the first one is its formulation in terms of reduced density matrices. In this paper we use GCC for the wave function. To generalize our approach, it might be useful to express the GCC in terms of the 2RDM<sup>18,19</sup>.

One may ask whether it is not more convenient to re-express our formulation in terms of the 1RDM. In 1975 Kimball found a relationship between the 2RDM at coalescence and the distribution of momentum, in fact, the behavior of the 1RDM<sup>27</sup>. At the same time Yasuhara demonstrated that the energy of the uniform electron gas can be expressed in terms of the kinetic energy (one-particle operators) instead of the two-body interactions<sup>28</sup>. This gave rise to studies on using the adiabatic connection on the kinetic energy (i.e. the 1RDM) rather than on two-body interactions<sup>29-31</sup>. Recently, the effects of the electron-electron coalescence on the properties of the natural orbitals and structure of the 1RDM have been revealed<sup>32,33</sup>.

Another issue is the dependence of the GCC on the coalescence point<sup>6-9</sup>. For this, we can get inspiration from density functional approximation: in each point of space one has a different approximation. Even the Kato's cusp condition (that seemingly is universal) contains a state dependence (through  $\ell$ ). In density functional calculation it is treated by using the spin polarization<sup>34</sup>, but it can be treated in the context of a pair density (see, e.g., ref 35).

We illustrate the problematic with two two-electron harmonium systems (A and B) that instead of being treated separately are treated together (for example, having two quantum dots). The system may be described by a double-well potential with two wells sufficiently separated and deep, so that each of them can be approximated by a harmonic potential. We assume that this separation is large enough to neglect the exchange effects between electrons

in A and B. Consequently, the spin part may be separated and, in effect, we can consider the total wave function which depends on the orbital variables only. It can be written as a product of two harmonium wave functions:

$$\Psi_{\text{total}}(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3, \mathbf{r}_4) = \Psi_A(\mathbf{r}_1, \mathbf{r}_2)\Psi_B(\mathbf{r}_3, \mathbf{r}_4) \quad (43)$$

Let us first consider that subsystem A is in a state with  $\ell = 0$ , while subsystem B is in a state with  $\ell = 1$ . In region A the coefficient  $\tilde{c}_1 = 1/2$ , while in region B,  $\tilde{c}_1 = 1/4$ . Let us now consider another example. In both subsystems we have  $\ell = 0$ . However, the values of  $\omega$  differ:  $\omega_A \neq \omega_B$ . We can see from Figures 3a, 3f-h that the range of where results are reliable depend on the model chosen. For example, we see in Figure 3h that if we choose  $\mu = 0.5$  we will have chemical accuracy for a subsystem with  $\omega = 1/4$ , but that the error is significantly larger for a subsystem with  $\omega = 1$  where chemical accuracy is reached only for  $\mu = 1$ . Do we choose to make the more expensive global calculation, say with  $\mu = 1$ , or do we decide to lose quality and make the cheaper calculation at  $\mu = 1/2$ ? Alternatively, we might consider making calculations with models that change from one region of space to another.

In how far do we need to go beyond the Kato cusp condition? Using eq (42), Figure 3h, shows that this might not be needed. Furthermore, eq (50) of the Appendix (see also discussion in ref 11) shows that there are terms that are important when  $r > 1/\mu$  even when  $r$  is small and they may need a more careful treatment.

Another question is raised by analyzing the dependence on the external potential. It was shown by Kurokawa *et al.*<sup>8</sup> that for small interelectronic distance  $r$ , the leading term in the expansion of the Coulomb potential of the nuclei is quadratic. So, some of the conclusions drawn from the present treatment of harmonium might be applied in systems where the external potential is Coulombic.

## VI. APPENDIX: A DERIVATION OF EQ (30)

We repeat here the argumentation given in ref 11, section III, where the result was obtained for  $\lambda = 0$ .

We consider the behavior of the Schrödinger equation

$$\left( -\frac{d^2}{dr^2} - \frac{2}{r} \frac{d}{dr} + \frac{(1-\lambda) \operatorname{erf}(\mu r) + \lambda}{r} + \frac{\omega^2 r^2}{2} - E \right) \psi(r) = 0 \quad (44)$$

at the limit of large  $\mu$ . After a change of the variable  $r$  to  $x = \mu r$ , and defining  $u(x) = \psi(x/\mu)$  one obtains

$$\left(-\frac{d^2}{dx^2} - \frac{2}{x} \frac{d}{dx} + \frac{(1-\lambda)\text{erf}(x) + \lambda}{\mu x} + \frac{\omega^2 x^2}{2\mu^4} - \frac{E}{\mu^2}\right) u(x) = 0. \quad (45)$$

For large  $\mu$ , we neglect terms proportional to  $(1/\mu)^n$  with  $n > 1$  and to the resulting equation apply perturbation theory up to the first order in  $1/\mu$ . We set:

$$u(x) = u^{(0)}(x) + \frac{1}{\mu} u^{(1)}(x). \quad (46)$$

The 0th order in  $1/\mu$  gives

$$\left(-\frac{d^2}{dx^2} - \frac{2}{x} \frac{d}{dx}\right) u^{(0)}(x) = 0. \quad (47)$$

From here we get

$$u^0(x) = \mathcal{A}_0 + \frac{\mathcal{A}_1}{x} \quad (48)$$

where  $\mathcal{A}_0$  and  $\mathcal{A}_1$  are integration constants. The wave function has to be finite at  $x = 0$ . Therefore,  $\mathcal{A}_1 = 0$ . Furthermore, as  $u^{(0)}$  corresponds to the solution at  $\mu = \infty$ , namely to the exact solution independent of either  $\lambda$  or  $\mu$ , we set  $\mathcal{A}_0 = \mathcal{N}$ . Consequently, the equation for  $u^{(1)}(x)$  is

$$\left(-\frac{d^2}{dx^2} - \frac{2}{x} \frac{d}{dx}\right) u^{(1)}(x) + \frac{(1-\lambda)\text{erf}(x) + \lambda}{x} \mathcal{N} = 0. \quad (49)$$

After solving this equation and changing variable  $x$  to  $r$ , we get

$$\psi(r) = \mathcal{N} \left(1 + \frac{\lambda}{2} r + \frac{1-\lambda}{2\sqrt{\pi}\mu} e^{-\mu^2 r^2} + (1-\lambda) \left(\frac{r}{2} + \frac{1}{4\mu^2 r}\right) \text{erf}(\mu r)\right) + \mathcal{B}_0 + \frac{\mathcal{B}_1}{\mu r}, \quad (50)$$

where  $\mathcal{B}_0$  and  $\mathcal{B}_1$  are integration constants. To avoid singularity of the wave function at  $r = 0$   $\mathcal{B}_1 = 0$ . In order to recover results for  $\lambda = 0$ <sup>11</sup> we have to set  $\mathcal{B}_0 = 0$ . Finally, in the limit  $r \rightarrow 0$ , up to the first-order in  $1/\mu$ , we have

$$\psi(r)|_{r=0} = \mathcal{N} \left(1 + \frac{1-\lambda}{\sqrt{\pi}\mu}\right), \quad (51)$$

where  $\mathcal{N}$  is a normalization constant independent of either  $\lambda$  or  $\mu$ .

## VII. ACKNOWLEDGEMENT

This work was done without specific financial support. It was presented in part at the MQM 2022 honoring Profs. Gustavo E. Scuseria (Rice University, Houston, Texas, U.S.A.) and Martin Head-Gordon (University of California, Berkeley, California, U.S.A.) held at Virginia Tech (Blacksburg, Virginia, U.S.A.).

## REFERENCES

- <sup>1</sup>González-Espinoza, C. E.; Ayers, P. W.; Karwowski, J.; Savin A. Smooth Models for the Coulomb Potential. *Theor. Chem. Acc.* **2016**, *135* (12), 256; DOI: 10.1007/s00214-016-2007-5.
- <sup>2</sup>Savin, A. Correcting Model Energies by Numerically Integrating Along an Adiabatic Connection and a Link to Density Functional Approximations. *J. Chem. Phys.* **2011**, *134* (21), 214108; DOI: 10.1063/1.3592782; DOI: 10.1063/1.3592782.
- <sup>3</sup>Savin A. Models and Corrections: Range Separation for Electronic Interaction — Lessons from Density Functional Theory. *J. Chem. Phys.* **2020**, *153* (16), 160901; DOI: 10.1063/5.0028060.
- <sup>4</sup>Löwdin, P. O. Quantum Theory of Many-Particle Systems. I. Physical Interpretations by Means of Density Matrices, Natural Spin-Orbitals, and Convergence Problems in the Method of Configurational Interaction. *Phys. Rev.* **1955**, *97* (6), 1474-1489; DOI: 10.1103/PhysRev.97.1474.
- <sup>5</sup>Kato, T. On the Eigenfunctions of Many-Particle Systems in Quantum Mechanics. *Commun. Pure Appl. Math.* **1957** *10* (2), 151-177; DOI: 10.1002/cpa.3160100201.
- <sup>6</sup>Karwowski, J.; Savin A. Two-Particle Coalescence Conditions Revisited. *Mol. Phys.* **2022**, *120* (19-20), e2069055; DOI: 10.1080/00268976.2022.2069055.
- <sup>7</sup>Kurokawa, Y. I.; Nakashima, H.; Nakatsuji, H. General Coalescence Conditions for the Exact Wave Functions: Higher-Order Relations for Two-Particle Systems. *J. Chem. Phys.* **2013**, *139* (4), 044114; DOI: 10.1063/1.4816281.
- <sup>8</sup>Kurokawa, Y. I.; Nakashima, H.; Nakatsuji, H. General Coalescence Conditions for the Exact Wave Functions. II. Higher-Order Relations for Many-Particle Systems. *J. Chem. Phys.* **2014**, *140* (21), 214103; DOI: 10.1063/1.4879266.
- <sup>9</sup>Kurokawa, Y. I.; Nakashima, H.; Nakatsuji, H. General Coalescence conditions for the Exact Wave Functions: Higher-Order Relations for Coulombic and Non-Coulombic Systems. *Adv. Quant. Chem.* **2016**, *73* (2), 59-79; DOI: 10.1016/bs.aiq.2015.09.001.
- <sup>10</sup>Gradshteyn, I. S.; Ryzhik, I. M. *Table of Integrals, Series and Products*; Jeffrey A.; Zwillinger D. Eds.; Elsevier/Academic Press, 2007, eqs 8.253, and 6.281.1.
- <sup>11</sup>Gori-Giorgi, P.; Savin, A. Properties of Short-Range and Long-Range Correlation Energy Density Functionals from Electron-Electron Coalescence. *Phys. Rev. A* **2006** *73* (3),

- 032506; DOI: 10.1103/PhysRevA.73.032506.
- <sup>12</sup>Kutzelnigg, W.; Morgan III, J. D. Rates of Convergence of the Partial-Wave Expansions of Atomic Correlation Energies. *J. Chem. Phys.* **1992**, *96* (6), 4484-4508; *J. Chem. Phys.* **1992**, *97* (11), 8821; DOI: 10.1063/1.463358.
- <sup>13</sup>King, H. F. The Electron Correlation Cusp. I. Overview and Partial Wave Analysis of the Kais Function. *Theor. Chim. Acta* **1996**, *94*, 345-381; DOI: 10.1007/BF00186448.
- <sup>14</sup>Savin, A.; Flad, H. J. Density Functionals for the Yukawa Electron-Electron Interaction. *Int. J. Quantum Chem.* **1995**, *56* (4), 327-332; DOI: 10.1002/qua.560560417.
- <sup>15</sup>Savin, A. In *Recent Developments of Modern Density Functional Theory*; Seminario, J. M., Ed.; Elsevier: Amsterdam, 1996; pp. 327–357.
- <sup>16</sup>Paziani, S.; Moroni, S.; Gori-Giorgi, P.; Bachelet, G. B. Local-Spin-Density Functional for Multideterminant Density Functional Theory. *Phys. Rev. B* **2006**, *73* (15), 155111; DOI: 10.1103/PhysRevB.73.155111.
- <sup>17</sup>Goll, E.; Werner, H.-J.; Stoll, H. A Short-Range Gradient-Corrected Density Functional in Long-Range Coupled-Cluster Calculations for Rare Gas Dimers. *Phys. Chem. Chem. Phys.* **2005**, *7* (23), 3917-3923; DOI: 10.1039/b509242f.
- <sup>18</sup>Nagy, A.; Sen, K. D. Ground- and Excited-State Cusp Conditions for the Electron Density. *J. Chem. Phys.* **2001**, *115* (14), 6300-6308; DOI: 10.1063/1.1402165.
- <sup>19</sup>Nagy, A.; Amovilli, C. Ground- and Excited-State Cusp Conditions for the Pair Density. *Phys. Rev. A* **2010**, *82* (4), 042510; DOI: 10.1103/PhysRevA.82.042510.
- <sup>20</sup>Chilkuri, V. G.; Neese, F. Comparison of Many-Particle Representations for Selected-CI I: A Tree Based Approach. *J. Comput. Chem.* **2021**, *42* (14), 982-1005; DOI: 10.1002/jcc.26518.
- <sup>21</sup>Jastrow, R. Many-Body Problem with Strong Forces. *Phys. Rev.* **1955**, *98* (5), 1479-1484; DOI: 10.1103/PhysRev.98.1479.
- <sup>22</sup>Luchow, A.; Sturm, A.; Schulte, C.; Mood, K. H. Generic Expansion of the Jastrow Correlation Factor in Polynomials Satisfying Symmetry and Cusp Conditions. *J. Chem. Phys.* **2015**, *142* (8), 084111; DOI: 10.1063/1.4909554.
- <sup>23</sup>Petruzielo, F. R.; Toulouse, J.; Umrigar, C. J. Approaching Chemical Accuracy with Quantum Monte Carlo. *J. Chem. Phys.* **2012**, *136* (12), 124116; DOI: 10.1063/1.3697846.
- <sup>24</sup>Booth, G. H.; Thom, A. J. W.; Alavi, A. Fermion Monte Carlo without Fixed Nodes: A Game of Life, Death, and Annihilation in Slater Determinant Space. *J. Chem. Phys.*

- 2009**, *131* (5), 054106; DOI: 10.1063/1.3193710.
- <sup>25</sup>Ten-no, S. Explicitly Correlated Wave Functions: Summary and Perspective. *Theor. Chem. Acc.* **2012**, *131* (1), 1070; DOI: 10.1007/s00214-011-1070-1.
- <sup>26</sup>Tew, D. P.; Klopper, W. Open-Shell Explicitly Correlated F12 Methods. *Mol. Phys.* **2010**, *108* (3-4), 315-325; DOI: 10.1080/00268970903449388.
- <sup>27</sup>Kimball, J. C. Short-Range Correlations and the Structure Factor and Momentum Distribution of Electrons. *J. Phys. A: Math. Gen.* **1975**, *8*, 1513-1517; DOI: 10.1088/0305-4470/8/9/021.
- <sup>28</sup>Yasuhara, H. Comment on Ground-State Energy Formula for Interacting Many-Particle Systems. *Lett. Nuovo Cimento* **1975**, *12*, 418-420; DOI: 10.1007/BF02782230.
- <sup>29</sup>Savin, A. Expression of the Exact Electron-Correlation-Energy Density-Functional in Terms of First-Order Density-Matrices. *Phys. Rev. A* **1995**, *52* (3), R1805-R1807; DOI: 10.1103/PhysRevA.52.R1805.
- <sup>30</sup>Levy, M.; Görling, A. Correlation-Energy Density-Functional Formulas from Correlating First-Order Density-Matrices. *Phys. Rev. A* **1995**, *52* (3), R1808-R1810; DOI: 10.1103/PhysRevA.52.R1808.
- <sup>31</sup>Teale, A. M.; Helgaker, T. U.; Savin, A. Alternative Representations of the Correlation Energy in Density-Functional Theory: A Kinetic-Energy Based Adiabatic Connection. *J. Chin. Chem. Soc.* **2016**, *63* (1), 121-128; DOI: 10.1002/jccs.201500132.
- <sup>32</sup>Cioslowski, J.; Strasburger, K. From Fredholm to Schrödinger via Eikonal: A New Formalism for Revealing Unknown Properties of Natural Orbitals. *J. Chem. Theory Comput.* **2021**, *17* (11), 6918-6933; DOI: 10.1021/acs.jctc.1c00709.
- <sup>33</sup>Cioslowski, J. Reverse Engineering in Quantum Chemistry: How to Reveal the Fifth-Order Off-Diagonal Cusp in the One-Electron Reduced Density Matrix Without Actually Calculating it. *Int. J. Quantum Chem.* **2021**, *122* (8), e26651; DOI: 10.1002/qua.26651.
- <sup>34</sup>Becke, A. D.; Savin, A.; Stoll, H. Extension of the Local-Spin-Density Exchange-Correlation Approximation to Multiplet States. *Theor. Chim. Acta* **1995**, *91* (3-4), 147-156; DOI: 10.1007/BF01114982.
- <sup>35</sup>Perdew, J. P.; Savin, A.; Burke, K. Escaping the Symmetry Dilemma through a Pair-Density Interpretation of Spin-Density Functional Theory. *Phys. Rev. A* **1995**, *51* (6), 4531-4541; DOI: 10.1103/PhysRevA.51.4531.