

Introduction to density-functional theory

Julien Toulouse
Laboratoire de Chimie Théorique
Sorbonne Université and CNRS, 75005 Paris, France
Institut Universitaire de France, F-75005 Paris, France
toulouse@lct.jussieu.fr

June 27, 2022

http://www.lct.jussieu.fr/pagesperso/toulouse/enseignement/introduction_dft.pdf

These lecture notes grew out of courses given at:

- 3rd edition of the International summer School in electronic structure Theory: electron correlation in Physics and Chemistry (ISTPC), 19 June - 2 July 2022, Aussois, France.
- 17th European Summerschool in Quantum Chemistry (ESQC), 8-21 September 2019, Sicily, Italy.
- 2nd edition of the International summer School in electronic structure Theory: electron correlation in Physics and Chemistry (ISTPC), 18 June - 1 July 2017, Aussois, France.
- ICS Summer School 2015 “Scientific Trends at the Interfaces: Mathematics - Chemistry - High Performance Computing”, 15 July - 14 August 2015, Roscoff, France.
- International summer School in electronic structure Theory: electron correlation in Physics and Chemistry (ISTPC), 14-27 June 2015, Aussois, France.

Contents

1	Basic density-functional theory	3
1.1	The many-body problem	3
1.2	The universal density functional	4
1.2.1	The Hohenberg-Kohn theorem	4
1.2.2	Levy-Lieb constrained-search formulation	6
1.3	The Kohn-Sham method	7
1.3.1	Decomposition of the universal functional	7
1.3.2	The Kohn-Sham equations	8
1.3.3	Practical calculations in an atomic basis	10
1.3.4	Extension to spin density-functional theory	12
1.4	The generalized Kohn-Sham method	13
2	Exact expressions and constraints for the exchange and correlation functionals	15
2.1	The exchange and correlation functionals in terms of the exchange and correlation holes	15
2.1.1	The exchange and correlation holes	15
2.1.2	The adiabatic connection	17

2.2	Uniform coordinate scaling	18
2.3	One-orbital spatial regions and self-interaction	20
2.4	Lieb-Oxford lower bound	20
2.5	Fractional electron numbers and frontier orbital energies	21
2.5.1	Quantum mechanics with fractional electron numbers	21
2.5.2	Density-functional theory with fractional electron numbers	22
2.5.3	The HOMO energy and the ionization energy	24
2.5.4	The LUMO energy, the electron affinity, and the derivative discontinuity	25
2.5.5	Fundamental gap	26
3	Usual approximations for the exchange-correlation energy	27
3.1	The local-density approximation	27
3.2	Generalized-gradient approximations	29
3.3	Meta-generalized-gradient approximations	30
3.4	Single-determinant hybrid approximations	31
3.4.1	Hybrid approximations	31
3.4.2	Range-separated hybrid approximations	33
3.5	Multideterminant hybrid approximations	34
3.5.1	Double-hybrid approximations	34
3.5.2	Range-separated double-hybrid approximations	35
3.6	Dispersion corrections	36
3.6.1	Semiempirical dispersion corrections	36
3.6.2	Nonlocal van der Waals density functionals	36
4	Some less usual orbital-dependent exchange-correlation functionals	37
4.1	Exact exchange	37
4.2	Second-order Görling-Levy perturbation theory	38
4.3	Adiabatic-connection fluctuation-dissipation approach	39
4.3.1	Exact adiabatic-connection fluctuation-dissipation expression	39
4.3.2	Direct random-phase approximation	41
4.3.3	Practical calculation in a spin orbital basis	42
4.3.4	Random-phase approximation with exchange and beyond	45
	Appendices	46
	A The Hamiltonian in second quantization	46
	B A brief introduction to functional derivatives	48
	Solutions to the exercises	51
	References	63

These notes provide an introduction to ground-state density-functional theory (DFT) of electronic systems. For more intensive coverages of the subject, see e.g. the books of Refs. [1–6] and the review articles of Refs. [7–17].

1 Basic density-functional theory

1.1 The many-body problem

We consider a N -electron system (atom, molecule, or solid) in the Born-Oppenheimer and non-relativistic approximations. The electronic Hamiltonian in the position representation is, in atomic units,

$$H = -\frac{1}{2} \sum_{i=1}^N \nabla_{\mathbf{r}_i}^2 + \frac{1}{2} \sum_{i=1}^N \sum_{\substack{j=1 \\ i \neq j}}^N \frac{1}{|\mathbf{r}_i - \mathbf{r}_j|} + \sum_{i=1}^N v_{\text{ne}}(\mathbf{r}_i), \quad (1.1)$$

where $v_{\text{ne}}(\mathbf{r}_i) = -\sum_{\alpha} Z_{\alpha}/|\mathbf{r}_i - \mathbf{R}_{\alpha}|$ is the nuclei-electron interaction (\mathbf{R}_{α} and Z_{α} are the positions and charges of the nuclei). The stationary electronic states are determined by the time-independent Schrödinger equation

$$H\Psi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N) = E\Psi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N), \quad (1.2)$$

where $\Psi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N)$ is a wave function written with space-spin coordinates $\mathbf{x}_i = (\mathbf{r}_i, \sigma_i)$ (with $\mathbf{r}_i \in \mathbb{R}^3$ and $\sigma_i \in \{\uparrow, \downarrow\}$) which is antisymmetric with respect to the exchange of two coordinates, and E is the associated energy. The one-electron Hilbert space is thus $L^2(\mathbb{R}^3 \times \{\uparrow, \downarrow\}, \mathbb{C})$ and the N -electron Hilbert space is given by the N -fold antisymmetric tensor product $\mathcal{H} = \bigwedge^N L^2(\mathbb{R}^3 \times \{\uparrow, \downarrow\}, \mathbb{C})$.

Using Dirac notations, the Schrödinger equation (1.2) can be rewritten in a representation-independent formalism

$$\hat{H}|\Psi\rangle = E|\Psi\rangle, \quad (1.3)$$

where the Hamiltonian is formally written as

$$\hat{H} = \hat{T} + \hat{W}_{\text{ee}} + \hat{V}_{\text{ne}}, \quad (1.4)$$

with the kinetic-energy operator \hat{T} , the electron-electron interaction operator \hat{W}_{ee} , and the nuclei-electron interaction operator \hat{V}_{ne} . These operators can be conveniently expressed in second quantization (see Appendix A).

The quantity of primary interest is the ground-state energy E_0 . The variational theorem establishes that E_0 can be obtained by the following minimization

$$E_0 = \min_{\Psi \in \mathcal{W}} \langle \Psi | \hat{H} | \Psi \rangle, \quad (1.5)$$

where the search is over the set of admissible N -electron antisymmetric wave functions Ψ normalized to unity, $\mathcal{W} = \{\Psi \in \bigwedge^N H^1(\mathbb{R}^3 \times \{\uparrow, \downarrow\}, \mathbb{C}) \mid \langle \Psi | \Psi \rangle = 1\}$, where $H^1(\mathbb{R}^3 \times \{\uparrow, \downarrow\}, \mathbb{C}) = \{\psi \in L^2(\mathbb{R}^3 \times \{\uparrow, \downarrow\}, \mathbb{C}) \mid \nabla \psi \in (L^2(\mathbb{R}^3 \times \{\uparrow, \downarrow\}, \mathbb{C}))^3\}$ is the first Sobolev space (ensuring a finite kinetic energy). DFT is based on a reformulation of the variational theorem in terms of the one-electron density defined as¹

$$n(\mathbf{r}) = N \int \dots \int |\Psi(\mathbf{x}, \mathbf{x}_2, \dots, \mathbf{x}_N)|^2 d\sigma d\mathbf{x}_2 \dots d\mathbf{x}_N, \quad (1.6)$$

which is normalized to the electron number, $\int n(\mathbf{r}) d\mathbf{r} = N$.

¹In Eq. (1.6), an integration over a spin coordinate σ just means a sum over the two values $\sigma \in \{\uparrow, \downarrow\}$.

1.2 The universal density functional

1.2.1 The Hohenberg-Kohn theorem

Consider an electronic system obtained by replacing the potential $v_{\text{ne}}(\mathbf{r})$ by an arbitrary external local potential $v(\mathbf{r})$ such that the corresponding Hamiltonian $\hat{T} + \hat{W}_{\text{ee}} + \hat{V}$ has a (bound) N -electron ground state. A ground-state wave function Ψ (there can be several of them if the ground state is degenerate) can be obtained by solving the Schrödinger equation, from which the associated ground-state density $n(\mathbf{r})$ can be deduced. Therefore, one has a mapping from the potential $v(\mathbf{r})$ to the considered ground-state density $n(\mathbf{r})$

$$v(\mathbf{r}) \longrightarrow n(\mathbf{r}). \quad (1.7)$$

In 1964, Hohenberg and Kohn [18] showed that this mapping can be inverted, i.e. the ground-state density $n(\mathbf{r})$ determines the potential $v(\mathbf{r})$ up to an arbitrary additive real-valued constant

$$\boxed{n(\mathbf{r}) \xrightarrow{\text{Hohenberg-Kohn}} v(\mathbf{r}) + \text{const.}} \quad (1.8)$$

Proof: The two-step proof by contradiction proceeds as follows (see, e.g., Ref. [2]).

We consider two local potentials $v_1(\mathbf{r})$ and $v_2(\mathbf{r})$ differing by more than an additive constant, $v_1(\mathbf{r}) \neq v_2(\mathbf{r}) + \text{const}$, and we note E_1 and E_2 the ground-state energies of the Hamiltonians $\hat{H}_1 = \hat{T} + \hat{W}_{\text{ee}} + \hat{V}_1$ and $\hat{H}_2 = \hat{T} + \hat{W}_{\text{ee}} + \hat{V}_2$, respectively.

(1) Assume that \hat{H}_1 and \hat{H}_2 have the same ground-state wave function Ψ , i.e. $\hat{H}_1|\Psi\rangle = E_1|\Psi\rangle$ and $\hat{H}_2|\Psi\rangle = E_2|\Psi\rangle$. Then, subtracting these two equations gives

$$(\hat{V}_1 - \hat{V}_2)|\Psi\rangle = (E_1 - E_2)|\Psi\rangle, \quad (1.9)$$

or, in position representation,

$$\sum_{i=1}^N [v_1(\mathbf{r}_i) - v_2(\mathbf{r}_i)] \Psi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N) = (E_1 - E_2) \Psi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N), \quad (1.10)$$

which implies $v_1(\mathbf{r}) - v_2(\mathbf{r}) = \text{const}$, in contradiction with the initial hypothesis. Note that, to eliminate Ψ in Eq. (1.10), it is assumed that $\Psi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N) \neq 0$ for at least one fixed set of spin coordinates $(\sigma_1, \sigma_2, \dots, \sigma_N)$ and for “almost” all spatial coordinates $(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N)$ (i.e., except possibly on a set of zero measure). This is in fact true for “reasonably well behaved” potentials. In this case, we thus conclude that two local potentials differing by more than an additive constant cannot share the same ground-state wave function.

(2) Let then Ψ_1 and Ψ_2 be (necessarily different) ground-state wave functions of \hat{H}_1 and \hat{H}_2 , respectively, and assume that Ψ_1 and Ψ_2 have the same ground-state density $n(\mathbf{r})$. The variational theorem leads to the following inequality

$$E_1 = \langle \Psi_1 | \hat{H}_1 | \Psi_1 \rangle < \langle \Psi_2 | \hat{H}_1 | \Psi_2 \rangle = \langle \Psi_2 | \hat{H}_2 + \hat{V}_1 - \hat{V}_2 | \Psi_2 \rangle = E_2 + \int [v_1(\mathbf{r}) - v_2(\mathbf{r})] n(\mathbf{r}) \text{d}\mathbf{r}, \quad (1.11)$$

where the strict inequality comes from the fact that Ψ_2 cannot be a ground-state wave function of \hat{H}_1 , as shown in the first step of the proof. Symmetrically, by exchanging the role of systems 1 and 2, we have the strict inequality

$$E_2 < E_1 + \int [v_2(\mathbf{r}) - v_1(\mathbf{r})] n(\mathbf{r}) \text{d}\mathbf{r}. \quad (1.12)$$

Adding Eqs. (1.11) and (1.12) gives the inconsistent result

$$E_1 + E_2 < E_1 + E_2, \quad (1.13)$$

which finally leads to the conclusion that there cannot exist two local potentials differing by more than an additive constant which have the same ground-state density. Note that this proof does not assume non-degenerate ground states (contrary to the original Hohenberg-Kohn proof). ■

So, the ground-state density $n(\mathbf{r})$ determines the potential $v(\mathbf{r})$, which in turn determines the Hamiltonian, and thus everything about the many-body problem. In other words, the potential v is a unique (up to an additive constant) functional of the ground-state density n , and all other properties as well. The ground-state wave function Ψ for the potential $v(\mathbf{r})$ is itself a functional of n , denoted by $\Psi[n]$, which was exploited by Hohenberg and Kohn to define the universal (i.e., independent from the external potential) density functional

$$F[n] = \langle \Psi[n] | \hat{T} + \hat{W}_{\text{ee}} | \Psi[n] \rangle, \quad (1.14)$$

which can be used to define the total electronic energy functional

$$E[n] = F[n] + \int v_{\text{ne}}(\mathbf{r})n(\mathbf{r})d\mathbf{r}, \quad (1.15)$$

for the specific external potential $v_{\text{ne}}(\mathbf{r})$ of the system considered. Note that, for degenerate ground states, $\Psi[n]$ is not unique but stands for any degenerate ground-state wave function. However, all $\Psi[n]$ give the same $F[n]$, which is thus a unique functional of n . Note also that the Hohenberg-Kohn functional in Eq. (1.14) is only defined for N -electron densities n that are ground-state densities associated with some local potential, the so-called set of v -representable densities which we will denote by \mathcal{A} .

Hohenberg and Kohn further showed that the density functional $E[n]$ satisfies a variational property: the ground-state energy E_0 of the system considered is obtained by minimizing this functional with respect to v -representable densities

$$E_0 = \min_{n \in \mathcal{A}} \left\{ F[n] + \int v_{\text{ne}}(\mathbf{r})n(\mathbf{r})d\mathbf{r} \right\}, \quad (1.16)$$

the minimum being reached for a ground-state density $n_0(\mathbf{r})$ corresponding to the potential $v_{\text{ne}}(\mathbf{r})$.

The existence of a mapping from a ground-state density to a local potential, the existence of the universal density functional, and the variational property with respect to the density constitutes the *Hohenberg-Kohn theorem*.

Exercise 1 : For the special case of Coulombic potentials $v_{\text{ne}}(\mathbf{r})$ there is a simple argument due to E. Bright Wilson showing that the ground-state density $n_0(\mathbf{r})$ fully determines $v_{\text{ne}}(\mathbf{r})$. Can you find it yourself?

Exercise 2 : Prove the variational property [Eq. (1.16)] of the density functional $E[n]$.

1.2.2 Levy-Lieb constrained-search formulation

In 1979 Levy [19,20], and later Lieb [21], proposed to redefine the universal density functional $F[n]$ using a *constrained-search formulation*

$$F[n] = \min_{\substack{\Psi \in \mathcal{W} \\ \Psi \rightarrow n}} \langle \Psi | \hat{T} + \hat{W}_{ee} | \Psi \rangle = \langle \Psi[n] | \hat{T} + \hat{W}_{ee} | \Psi[n] \rangle, \quad (1.17)$$

where $\Psi \rightarrow n$ means that the wave function Ψ is constrained to yield the fixed density n [via Eq. (1.6)]. For a given density n , the minimizing wave function is denoted by $\Psi[n]$ (possibly non unique in case of degeneracy). This so-called Levy-Lieb functional $F[n]$ does not require the existence of a local potential associated with the density. It is an extension of the Hohenberg-Kohn functional: it is defined on the larger set of N -electron densities coming from a wave function $\Psi \in \mathcal{W}$, the so-called set of N -representable densities which we will denote by \mathcal{D} . It turns out that the set \mathcal{D} is known explicitly: $\mathcal{D} = \{n \in L^1(\mathbb{R}^3) \mid n \geq 0, \int n(\mathbf{r})d\mathbf{r} = N, \sqrt{n} \in H^1(\mathbb{R}^3)\}$.

The variational property of the total electronic energy functional can easily be demonstrated using the constrained-search formulation. One starts from the usual variational theorem and decomposes the minimization over Ψ in two steps: a constrained minimization over Ψ giving a fixed density n , followed by a minimization over n ,

$$\begin{aligned} E_0 &= \min_{\Psi \in \mathcal{W}} \langle \Psi | \hat{T} + \hat{W}_{ee} + \hat{V}_{ne} | \Psi \rangle \\ &= \min_{n \in \mathcal{D}} \min_{\substack{\Psi \in \mathcal{W} \\ \Psi \rightarrow n}} \langle \Psi | \hat{T} + \hat{W}_{ee} + \hat{V}_{ne} | \Psi \rangle \\ &= \min_{n \in \mathcal{D}} \left\{ \min_{\substack{\Psi \in \mathcal{W} \\ \Psi \rightarrow n}} \langle \Psi | \hat{T} + \hat{W}_{ee} | \Psi \rangle + \int v_{ne}(\mathbf{r})n(\mathbf{r})d\mathbf{r} \right\} \\ &= \min_{n \in \mathcal{D}} \left\{ F[n] + \int v_{ne}(\mathbf{r})n(\mathbf{r})d\mathbf{r} \right\}, \end{aligned} \quad (1.18)$$

and again the minimum is reached for a ground-state density $n_0(\mathbf{r})$ corresponding to the potential $v_{ne}(\mathbf{r})$.

The ground-state energy and density can then be in principle obtained by minimizing over the density $n(\mathbf{r})$, i.e. a simple function of 3 variables, which is a tremendous simplification compared to the minimization over a complicated many-body wave function Ψ . However, the explicit expression of $F[n]$ in terms of the density is not known, and the direct approximations for $F[n]$ that have been tried so far turn out not to be accurate enough, especially for the kinetic energy part $T[n]$ included in $F[n]$

$$F[n] = T[n] + W_{ee}[n], \quad (1.19)$$

where $T[n] = \langle \Psi[n] | \hat{T} | \Psi[n] \rangle$ and $W_{ee}[n] = \langle \Psi[n] | \hat{W}_{ee} | \Psi[n] \rangle$. Note that the decomposition of Eq. (1.19) is well defined if the minimizing wave function $\Psi[n]$ is unique (up to an irrelevant phase factor).

Remark:

Valone [22] and Lieb [21] generalized the constrained-search approach from pure states to ensemble density matrices $\hat{\Gamma}$, resulting in a new universal density functional referred to as the density-matrix (DM) density functional or Lieb density functional

$$F_{\text{DM}}[n] = \min_{\substack{\hat{\Gamma} \in \mathcal{D}_{\text{DM}} \\ \hat{\Gamma} \rightarrow n}} \text{Tr}[(\hat{T} + \hat{W}_{ee})\hat{\Gamma}], \quad (1.20)$$

where Tr denotes the trace² and the minimization is done over admissible N -electron ensemble density matrices $\hat{\Gamma} \in \mathcal{D}_{\text{DM}} = \{\hat{\Gamma} = \sum_i w_i |\Psi_i\rangle\langle\Psi_i|, 0 \leq w_i \leq 1, \sum_i w_i = 1, \Psi_i \in \mathcal{W}, \langle\Psi_i|\Psi_j\rangle = \delta_{i,j}\}$ yielding the fixed density n . This density functional is defined over the same set of N -representable densities and it is always lower than the Levy-Lieb density functional, i.e. $F_{\text{DM}}[n] \leq F[n]$, with equality if the minimizing wave function $\Psi[n]$ in Eq. (1.17) is non-degenerate. Introducing the ground-state energy $E_0[v]$ as a functional of the external potential v , Lieb showed that this functional is the Legendre-Fenchel transform of $E_0[v]$

$$F_{\text{DM}}[n] = \sup_{v \in \mathcal{V}} \left(E_0[v] - \int v(\mathbf{r})n(\mathbf{r})d\mathbf{r} \right), \quad (1.21)$$

where the space of potentials is $\mathcal{V} = L^{3/2}(\mathbb{R}^3) + L^\infty(\mathbb{R}^3)$. The density functional $F_{\text{DM}}[n]$ has the mathematical advantage of being convex. However, in the rest of these notes, we will only use the Levy-Lieb density functional.

1.3 The Kohn-Sham method

1.3.1 Decomposition of the universal functional

Faced with the difficulty of approximating directly $F[n]$, Kohn and Sham (KS) [23] proposed to decompose $F[n]$ as

$$F[n] = T_s[n] + E_{\text{Hxc}}[n], \quad (1.22)$$

where $T_s[n]$ is the non-interacting kinetic-energy functional which can be defined with a constrained-search formulation

$$T_s[n] = \min_{\substack{\Phi \in \mathcal{S} \\ \Phi \rightarrow n}} \langle \Phi | \hat{T} | \Phi \rangle = \langle \Phi[n] | \hat{T} | \Phi[n] \rangle, \quad (1.23)$$

where the minimization is done over *single-determinant* wave functions $\Phi \in \mathcal{S} = \{\Phi = \psi_1 \wedge \psi_2 \wedge \dots \wedge \psi_N \mid \psi_i \in H^1(\mathbb{R}^3 \times \{\uparrow, \downarrow\}, \mathbb{C}), \langle \psi_i | \psi_j \rangle = \delta_{i,j}\}$ yielding the fixed density n . For a given density n , the (non necessarily unique) minimizing single-determinant wave function is called the *KS wave function* and is denoted by $\Phi[n]$. The remaining functional $E_{\text{Hxc}}[n]$ in Eq. (1.22) is called the Hartree-exchange-correlation functional.

Even though the wave function is restricted to be a single-determinant wave function in Eq. (1.23), the functional $T_s[n]$ is still defined over the entire set of N -representable densities \mathcal{D} . This is because any N -representable density can be obtained from a single-determinant wave function. Therefore, the decomposition in Eq. (1.22) does not introduce any approximation. The idea of the KS method is then to use the exact expression of $T_s[n]$ by reformulating the variational property of $F[n]$ in terms of single-determinant wave functions Φ

$$\begin{aligned} E_0 &= \min_{n \in \mathcal{D}} \left\{ F[n] + \int v_{\text{ne}}(\mathbf{r})n(\mathbf{r})d\mathbf{r} \right\}, \\ &= \min_{n \in \mathcal{D}} \left\{ \min_{\substack{\Phi \in \mathcal{S} \\ \Phi \rightarrow n}} \langle \Phi | \hat{T} | \Phi \rangle + E_{\text{Hxc}}[n] + \int v_{\text{ne}}(\mathbf{r})n(\mathbf{r})d\mathbf{r} \right\} \\ &= \min_{n \in \mathcal{D}} \min_{\substack{\Phi \in \mathcal{S} \\ \Phi \rightarrow n}} \left\{ \langle \Phi | \hat{T} + \hat{V}_{\text{ne}} | \Phi \rangle + E_{\text{Hxc}}[n_\Phi] \right\} \\ &= \min_{\Phi \in \mathcal{S}} \left\{ \langle \Phi | \hat{T} + \hat{V}_{\text{ne}} | \Phi \rangle + E_{\text{Hxc}}[n_\Phi] \right\}, \end{aligned} \quad (1.24)$$

²The trace of an operator \hat{A} is defined as $\text{Tr}[\hat{A}] = \sum_n \langle \Psi_n | \hat{A} | \Psi_n \rangle$ where $\{\Psi_n\}$ is an orthonormal basis of states.

the minimizing single-determinant KS wave function giving the exact ground-state density $n_0(\mathbf{r})$. Thus, the exact ground-state energy and density can in principle be obtained by minimizing over single-determinant wave functions only. Even though a wave function has been reintroduced compared to Eq. (1.18), it is only a single-determinant wave function Φ and therefore it still represents a tremendous simplification over the usual variational theorem involving a multi-determinant wave function Ψ . The advantage of Eq. (1.24) over Eq. (1.18) is that a major part of the kinetic energy can be treated explicitly with the single-determinant wave function Φ , and only $E_{\text{Hxc}}[n]$ needs to be approximated as a functional of the density.

In practice, $E_{\text{Hxc}}[n]$ is written as

$$\boxed{E_{\text{Hxc}}[n] = E_{\text{H}}[n] + E_{\text{xc}}[n]}, \quad (1.25)$$

where $E_{\text{H}}[n]$ is the Hartree energy functional

$$\boxed{E_{\text{H}}[n] = \frac{1}{2} \iint \frac{n(\mathbf{r}_1)n(\mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2}, \quad (1.26)$$

representing the classical electrostatic repulsion energy for the charge distribution $n(\mathbf{r})$, and $E_{\text{xc}}[n]$ is the exchange-correlation energy functional that remains to approximate. Assuming that the KS wave function $\Phi[n]$ is unique (up to a phase factor), the latter functional can be further decomposed as

$$\boxed{E_{\text{xc}}[n] = E_{\text{x}}[n] + E_{\text{c}}[n]}, \quad (1.27)$$

where $E_{\text{x}}[n]$ is the exchange energy functional

$$\boxed{E_{\text{x}}[n] = \langle \Phi[n] | \hat{W}_{\text{ee}} | \Phi[n] \rangle - E_{\text{H}}[n]}, \quad (1.28)$$

and $E_{\text{c}}[n]$ is the correlation energy functional

$$E_{\text{c}}[n] = \langle \Psi[n] | \hat{T} + \hat{W}_{\text{ee}} | \Psi[n] \rangle - \langle \Phi[n] | \hat{T} + \hat{W}_{\text{ee}} | \Phi[n] \rangle = T_{\text{c}}[n] + U_{\text{c}}[n], \quad (1.29)$$

which contains a kinetic contribution $T_{\text{c}}[n] = \langle \Psi[n] | \hat{T} | \Psi[n] \rangle - \langle \Phi[n] | \hat{T} | \Phi[n] \rangle$ and a potential contribution $U_{\text{c}}[n] = \langle \Psi[n] | \hat{W}_{\text{ee}} | \Psi[n] \rangle - \langle \Phi[n] | \hat{W}_{\text{ee}} | \Phi[n] \rangle$. The exchange and correlation functionals are always negative, i.e. $E_{\text{x}}[n] \leq 0$ and $E_{\text{c}}[n] \leq 0$.

1.3.2 The Kohn-Sham equations

The single-determinant wave function Φ is constructed from a set of N orthonormal occupied spin-orbitals $\{\psi_i(\mathbf{x})\}_{i=1,\dots,N}$. To enforce \hat{S}_z symmetry, each spin-orbital is factorized as $\psi_i(\mathbf{x}) = \varphi_i(\mathbf{r})\chi_{\sigma_i}(\sigma)$ where $\varphi_i(\mathbf{r})$ is a spatial orbital and $\chi_{\sigma_i}(\sigma) = \delta_{\sigma_i,\sigma}$ is a spin function (σ_i is the spin of the spin-orbital i). Alternatively, when this is convenient, we will sometimes reindex the spatial orbitals, $\{\varphi_i(\mathbf{r})\} \rightarrow \{\varphi_{i\sigma}(\mathbf{r})\}$, including explicitly the spin σ in the index. Writing the total electronic energy in Eq. (1.24) in terms of spin-orbitals and integrating over the spin variables, we obtain

$$\boxed{E[\{\varphi_i\}] = \sum_{i=1}^N \int \varphi_i^*(\mathbf{r}) \left(-\frac{1}{2} \nabla^2 + v_{\text{ne}}(\mathbf{r}) \right) \varphi_i(\mathbf{r}) d\mathbf{r} + E_{\text{Hxc}}[n]}, \quad (1.30)$$

where the density is expressed in terms of the orbitals as

$$\boxed{n(\mathbf{r}) = \sum_{i=1}^N |\varphi_i(\mathbf{r})|^2}. \quad (1.31)$$

The minimization over Φ can then be recast into a minimization of $E[\{\varphi_i\}]$ with respect to the spatial orbitals $\varphi_i(\mathbf{r})$ with the constraint of keeping the orbitals orthonormalized. Using the method of Lagrange multipliers, we introduce the following Lagrangian

$$\mathcal{L}[\{\varphi_i\}] = E[\{\varphi_i\}] - \sum_{i=1}^N \varepsilon_i \left(\int \varphi_i^*(\mathbf{r}) \varphi_i(\mathbf{r}) d\mathbf{r} - 1 \right), \quad (1.32)$$

where ε_i is the Lagrange multiplier associated to the normalization condition of $\varphi_i(\mathbf{r})$ ³. The Lagrangian should then be stationary with respect to variations of the orbitals $\varphi_i(\mathbf{r})$ ⁴

$$\frac{\delta \mathcal{L}}{\delta \varphi_i^*(\mathbf{r})} = 0, \quad (1.33)$$

where $\delta \mathcal{L} / \delta \varphi_i^*(\mathbf{r})$ is the functional derivative of \mathcal{L} with respect to $\varphi_i^*(\mathbf{r})$. Calculating this functional derivative gives (see Appendix B for an introduction to functional derivatives)

$$\left(-\frac{1}{2} \nabla^2 + v_{\text{ne}}(\mathbf{r}) \right) \varphi_i(\mathbf{r}) + \frac{\delta E_{\text{Hxc}}[n]}{\delta \varphi_i^*(\mathbf{r})} = \varepsilon_i \varphi_i(\mathbf{r}), \quad (1.34)$$

where the term $\delta E_{\text{Hxc}}[n] / \delta \varphi_i^*(\mathbf{r})$ can be expressed as, using the chain rule,

$$\frac{\delta E_{\text{Hxc}}[n]}{\delta \varphi_i^*(\mathbf{r})} = \int \frac{\delta E_{\text{Hxc}}[n]}{\delta n(\mathbf{r}')} \frac{\delta n(\mathbf{r}')}{\delta \varphi_i^*(\mathbf{r})} d\mathbf{r}'. \quad (1.35)$$

Noting that $\delta n(\mathbf{r}') / \delta \varphi_i^*(\mathbf{r}) = \varphi_i(\mathbf{r}) \delta(\mathbf{r} - \mathbf{r}')$ [from Eq. (1.31)], and defining the Hartree-exchange-correlation potential $v_{\text{Hxc}}(\mathbf{r})$ as the functional derivative of $E_{\text{Hxc}}[n]$ with respect to $n(\mathbf{r})$

$$\boxed{v_{\text{Hxc}}(\mathbf{r}) = \frac{\delta E_{\text{Hxc}}[n]}{\delta n(\mathbf{r})}}, \quad (1.36)$$

which is itself a functional of the density, we then arrive at the KS equations

$$\boxed{\left(-\frac{1}{2} \nabla^2 + v_{\text{ne}}(\mathbf{r}) + v_{\text{Hxc}}(\mathbf{r}) \right) \varphi_i(\mathbf{r}) = \varepsilon_i \varphi_i(\mathbf{r})}. \quad (1.37)$$

The orbitals satisfying Eq. (1.37) are called the KS orbitals. They are the eigenfunctions of the KS one-electron Hamiltonian

$$h_s = -\frac{1}{2} \nabla^2 + v_s(\mathbf{r}), \quad (1.38)$$

where

$$v_s(\mathbf{r}) = v_{\text{ne}}(\mathbf{r}) + v_{\text{Hxc}}(\mathbf{r}), \quad (1.39)$$

is the KS potential, and ε_i are then the KS orbital energies. Note that Eq. (1.37) constitutes a set of coupled self-consistent equations since the potential $v_{\text{Hxc}}(\mathbf{r})$ depends on all the occupied orbitals $\{\varphi_i\}_{i=1, \dots, N}$ through the density [Eq. (1.31)]. At convergence, the orbitals obtained by

³It turns out that it is not necessary to include the orthogonalization conditions $\int \varphi_i^*(\mathbf{r}) \varphi_j(\mathbf{r}) d\mathbf{r} = 0$ for $i \neq j$ in Eq. (1.32). Indeed, we will find that the minimizing orbitals $\varphi_i(\mathbf{r})$ are eigenfunctions of a self-adjoint operator, which implies that we can always find an orthogonal set of orbitals

⁴Here, the orbitals are assumed to take complex values, so that it is as if $\varphi_i(\mathbf{r})$ and $\varphi_i^*(\mathbf{r})$ were taken as independent functions. We then write the stationary equation for variations with respect to $\varphi_i^*(\mathbf{r})$ only, the second stationary equation for variations with respect to $\varphi_i(\mathbf{r})$ is just the complex conjugate of the first one.

solving Eq. (1.37) must be the same as the orbitals used to construct $v_{\text{Hxc}}(\mathbf{r})$. The operator h_s defines the KS system which is a system of N non-interacting electrons in an effective external potential $v_s(\mathbf{r})$ ensuring that its ground-state density $n(\mathbf{r})$ is the same as the exact ground-state density $n_0(\mathbf{r})$ of the physical system of N interacting electrons. The exact ground-state energy E_0 is then obtained by injecting the KS orbitals in Eq. (1.30). Note that Eq. (1.37) also permits to define virtual KS orbitals $\{\varphi_a\}_{a \geq N+1}$.

Note that to define the potential $v_{\text{Hxc}}(\mathbf{r})$ in Eq. (1.36) a form of differentiability of $E_{\text{Hxc}}[n]$ has been assumed. This can in fact only hold on a restricted set of densities. This is known as the v -representability problem. Also, note that the KS potential in Eq. (1.39) is defined only up to an additive constant. For atomic and molecular ground-state densities, we choose the constant so that the potential vanishes at infinity, $v_s(\infty) = 0$.

Following the decomposition of $E_{\text{Hxc}}[n]$ in Eq. (1.25), the potential $v_{\text{Hxc}}(\mathbf{r})$ is decomposed as

$$v_{\text{Hxc}}(\mathbf{r}) = v_{\text{H}}(\mathbf{r}) + v_{\text{xc}}(\mathbf{r}), \quad (1.40)$$

where $v_{\text{H}}(\mathbf{r}) = \delta E_{\text{H}}[n]/\delta n(\mathbf{r})$ is the Hartree potential and $v_{\text{xc}}(\mathbf{r}) = \delta E_{\text{xc}}[n]/\delta n(\mathbf{r})$ is the exchange-correlation potential. Likewise, following the decomposition of $E_{\text{xc}}[n]$ in Eq. (1.27), the potential $v_{\text{xc}}(\mathbf{r})$ can be decomposed as

$$v_{\text{xc}}(\mathbf{r}) = v_{\text{x}}(\mathbf{r}) + v_{\text{c}}(\mathbf{r}), \quad (1.41)$$

where $v_{\text{x}}(\mathbf{r}) = \delta E_{\text{x}}[n]/\delta n(\mathbf{r})$ is the exchange potential and $v_{\text{c}}(\mathbf{r}) = \delta E_{\text{c}}[n]/\delta n(\mathbf{r})$ is the correlation potential. Thus, the Kohn-Sham equations are similar to the Hartree-Fock equations, with the difference that they involve a local exchange potential $v_{\text{x}}(\mathbf{r})$ instead of a nonlocal one, and an additional correlation potential.

Exercise 3 : Show that the expression of the Hartree potential is

$$v_{\text{H}}(\mathbf{r}) = \int \frac{n(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r}'. \quad (1.42)$$

1.3.3 Practical calculations in an atomic basis

In practical calculations for molecular systems, we usually work in a basis of M atomic functions $\{\chi_\nu\}$, e.g. Gaussian-type basis functions centered on the nuclei. We then expand the orbitals as

$$\varphi_i(\mathbf{r}) = \sum_{\nu=1}^M c_{\nu i} \chi_\nu(\mathbf{r}), \quad (1.43)$$

and thus calculating the orbitals amounts to calculating the orbital coefficients $c_{\nu i}$. Inserting Eq. (1.43) into the KS equations

$$h_s \varphi_i(\mathbf{r}) = \varepsilon_i \varphi_i(\mathbf{r}), \quad (1.44)$$

multiplying on the left by $\chi_\mu^*(\mathbf{r})$ and integrating over \mathbf{r} , we arrive at

$$\sum_{\nu=1}^M F_{\mu\nu} c_{\nu i} = \varepsilon_i \sum_{\nu=1}^M S_{\mu\nu} c_{\nu i}, \quad (1.45)$$

where $F_{\mu\nu} = \int \chi_\mu^*(\mathbf{r}) h_s \chi_\nu(\mathbf{r}) d\mathbf{r}$ are the elements of the KS Fock matrix and $S_{\mu\nu} = \int \chi_\mu^*(\mathbf{r}) \chi_\nu(\mathbf{r}) d\mathbf{r}$ are the elements of the overlap matrix of the basis functions.

The Fock matrix is calculated as

$$F_{\mu\nu} = h_{\mu\nu} + J_{\mu\nu} + V_{xc,\mu\nu}, \quad (1.46)$$

where $h_{\mu\nu}$ are the one-electron integrals

$$h_{\mu\nu} = \int \chi_\mu^*(\mathbf{r}) \left(-\frac{1}{2} \nabla^2 + v_{ne}(\mathbf{r}) \right) \chi_\nu(\mathbf{r}) d\mathbf{r}, \quad (1.47)$$

$J_{\mu\nu}$ is the Hartree potential contribution

$$J_{\mu\nu} = \int \chi_\mu^*(\mathbf{r}) v_H(\mathbf{r}) \chi_\nu(\mathbf{r}) d\mathbf{r} = \sum_{\lambda=1}^M \sum_{\gamma=1}^M P_{\gamma\lambda} (\chi_\mu \chi_\nu | \chi_\lambda \chi_\gamma), \quad (1.48)$$

with the density matrix

$$P_{\gamma\lambda} = \sum_{i=1}^N c_{\gamma i} c_{\lambda i}^*, \quad (1.49)$$

and the two-electron integrals (in chemists' notation)

$$(\chi_\mu \chi_\nu | \chi_\lambda \chi_\gamma) = \iint \frac{\chi_\mu^*(\mathbf{r}_1) \chi_\nu(\mathbf{r}_1) \chi_\lambda^*(\mathbf{r}_2) \chi_\gamma(\mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2, \quad (1.50)$$

and $V_{xc,\mu\nu}$ is the exchange-correlation potential contribution

$$V_{xc,\mu\nu} = \int \chi_\mu^*(\mathbf{r}) v_{xc}(\mathbf{r}) \chi_\nu(\mathbf{r}) d\mathbf{r}. \quad (1.51)$$

In Eq. (1.51), the exchange-correlation potential $v_{xc}(\mathbf{r})$ is evaluated at the density calculated as

$$n(\mathbf{r}) = \sum_{\gamma=1}^M \sum_{\lambda=1}^M P_{\gamma\lambda} \chi_\gamma(\mathbf{r}) \chi_\lambda^*(\mathbf{r}). \quad (1.52)$$

Exercise 4 : Check Eq. (1.52) and prove the second equality in Eq. (1.48).

Eq. (1.45) is a self-consistent generalized eigenvalue equation that must be solved iteratively for finding the KS orbital coefficients and KS orbital energies. The converged density matrix can then be used to obtain the total electronic energy

$$E = \sum_{\mu=1}^M \sum_{\nu=1}^M P_{\nu\mu} h_{\mu\nu} + \frac{1}{2} \sum_{\mu=1}^M \sum_{\nu=1}^M P_{\nu\mu} J_{\mu\nu} + E_{xc}, \quad (1.53)$$

where E_{xc} is calculated with the density in Eq. (1.52).

In the simplest approximations (see Section 3), the exchange-correlation energy functional has a local form

$$E_{xc}^{\text{local}} = \int f(n(\mathbf{r})) d\mathbf{r}, \quad (1.54)$$

where $f(n(\mathbf{r}))$ has a complicated nonlinear dependence on the density $n(\mathbf{r})$. For example, in the local-density approximation (LDA) (see Section 3.1), the exchange energy is

$$E_x^{\text{LDA}} = C_x \int n(\mathbf{r})^{4/3} d\mathbf{r}, \quad (1.55)$$

where C_x is a constant, and the exchange potential is

$$v_x^{\text{LDA}}(\mathbf{r}) = \frac{4}{3} C_x n(\mathbf{r})^{1/3}. \quad (1.56)$$

Therefore, the integrals in Eq. (1.51) and Eq. (1.54) cannot be calculated analytically, but are instead evaluated by numerical integration

$$V_{xc,\mu\nu} \approx \sum_k w_k \chi_\mu^*(\mathbf{r}_k) v_{xc}(\mathbf{r}_k) \chi_\nu(\mathbf{r}_k), \quad (1.57)$$

and

$$E_{xc}^{\text{local}} \approx \sum_k w_k f(n(\mathbf{r}_k)), \quad (1.58)$$

where \mathbf{r}_k and w_k are quadrature points and weights. For example, for polyatomic molecules, the multicenter numerical integration scheme of Becke [24] is generally used.

1.3.4 Extension to spin density-functional theory

For dealing with an external magnetic field, DFT has been extended from the total density to spin-resolved densities [25, 26]. Without external magnetic fields, this spin density-functional theory is in principle not necessary, even for open-shell systems. In practice, however, the dependence on the spin densities allows one to construct approximate exchange-correlation functionals that are more accurate, and is therefore almost always used for open-shell systems.

The spin density $n_\sigma(\mathbf{r})$ for $\sigma \in \{\uparrow, \downarrow\}$ is defined as

$$n_\sigma(\mathbf{r}) = N \int \cdots \int |\Psi(\mathbf{r}\sigma, \mathbf{x}_2, \dots, \mathbf{x}_N)|^2 d\mathbf{x}_2 \dots d\mathbf{x}_N, \quad (1.59)$$

and integrates to the number of σ -spin electrons, i.e. $\int n_\sigma(\mathbf{r}) d\mathbf{r} = N_\sigma$. The universal density functional is now defined as [27]

$$F[n_\uparrow, n_\downarrow] = \min_{\substack{\Psi \in \mathcal{W} \\ \Psi \rightarrow n_\uparrow, n_\downarrow}} \langle \Psi | \hat{T} + \hat{W}_{ee} | \Psi \rangle, \quad (1.60)$$

where the search is over wave functions $\Psi \in \mathcal{W}$ with $N = N_\uparrow + N_\downarrow$ electrons and which yield fixed spin densities. A KS method is obtained by decomposing $F[n_\uparrow, n_\downarrow]$ as

$$F[n_\uparrow, n_\downarrow] = T_s[n_\uparrow, n_\downarrow] + E_H[n] + E_{xc}[n_\uparrow, n_\downarrow], \quad (1.61)$$

where $T_s[n_\uparrow, n_\downarrow]$ is defined with a constrained search over (spin-unrestricted) Slater determinants Φ

$$T_s[n_\uparrow, n_\downarrow] = \min_{\substack{\Phi \in \mathcal{S} \\ \Phi \rightarrow n_\uparrow, n_\downarrow}} \langle \Phi | \hat{T} | \Phi \rangle, \quad (1.62)$$

and $E_H[n]$ is the Hartree energy which is a functional of the total density $n = n_\uparrow + n_\downarrow$ only [Eq. (1.26)], and $E_{xc}[n_\uparrow, n_\downarrow]$ is the spin-resolved exchange-correlation energy functional. Writing the spatial orbitals of the spin-unrestricted determinant as $\varphi_{i\uparrow}(\mathbf{r})$ and $\varphi_{i\downarrow}(\mathbf{r})$ (with indices explicitly including spin now for clarity), we have now the spin-dependent KS equations

$$\left(-\frac{1}{2} \nabla^2 + v_{ne}(\mathbf{r}) + v_H(\mathbf{r}) + v_{xc,\sigma}(\mathbf{r}) \right) \varphi_{i\sigma}(\mathbf{r}) = \varepsilon_{i\sigma} \varphi_{i\sigma}(\mathbf{r}), \quad (1.63)$$

with the spin-dependent exchange-correlation potential

$$v_{\text{xc},\sigma}(\mathbf{r}) = \frac{\delta E_{\text{xc}}[n_{\uparrow}, n_{\downarrow}]}{\delta n_{\sigma}(\mathbf{r})}, \quad (1.64)$$

and the spin density

$$n_{\sigma}(\mathbf{r}) = \sum_{i=1}^{N_{\sigma}} |\varphi_{i\sigma}(\mathbf{r})|^2. \quad (1.65)$$

It turns out that the spin-dependent exchange functional $E_{\text{x}}[n_{\uparrow}, n_{\downarrow}]$ can be exactly expressed in terms of the spin-independent exchange functional $E_{\text{x}}[n]$ [28]

$$E_{\text{x}}[n_{\uparrow}, n_{\downarrow}] = \frac{1}{2} (E_{\text{x}}[2n_{\uparrow}] + E_{\text{x}}[2n_{\downarrow}]), \quad (1.66)$$

which is known as the *spin-scaling relation* and stems directly from the fact the \uparrow - and \downarrow -spin electrons are uncoupled in the exchange energy. Therefore, any approximation for the spin-independent exchange functional $E_{\text{x}}[n]$ can be easily extended to an approximation for the spin-dependent exchange functional $E_{\text{x}}[n_{\uparrow}, n_{\downarrow}]$. Unfortunately, there is no such relation for the correlation functional.

Exercise 5 : Prove the spin-scaling relation of Eq. (1.66).

Obviously, in the spin-unpolarized case, i.e. $n_{\uparrow} = n_{\downarrow} = n/2$, this spin-dependent formalism reduces to the spin-independent one.

1.4 The generalized Kohn-Sham method

An important extension of the KS method is the so-called *generalized Kohn-Sham* (GKS) method [29], which recognizes that the universal density functional $F[n]$ of Eq. (1.17) can be decomposed in other ways than in the KS decomposition of Eq. (1.22). In particular, we can decompose $F[n]$ as

$$F[n] = \min_{\substack{\Phi \in \mathcal{S} \\ \Phi \rightarrow n}} \left\{ \langle \Phi | \hat{T} | \Phi \rangle + E_{\text{H}}[n_{\Phi}] + S[\Phi] \right\} + \bar{S}[n], \quad (1.67)$$

where $S[\Phi]$ is any functional of a single-determinant wave function $\Phi \in \mathcal{S}$ leading to a minimum in Eq. (1.67), and $\bar{S}[n]$ is the corresponding complementary density functional that makes Eq. (1.67) exact. Defining the S -dependent GKS exchange-correlation functional as

$$E_{\text{xc}}^S[\Phi] = S[\Phi] + \bar{S}[n_{\Phi}], \quad (1.68)$$

we can express the exact ground-state energy as

$$E_0 = \min_{\Phi \in \mathcal{S}} \left\{ \langle \Phi | \hat{T} + \hat{V}_{\text{ne}} | \Phi \rangle + E_{\text{H}}[n_{\Phi}] + E_{\text{xc}}^S[\Phi] \right\}, \quad (1.69)$$

and any minimizing single-determinant wave function in Eq. (1.69) gives a ground-state density $n_0(\mathbf{r})$. Similarly to the KS equations [Eq. (1.37)], Eq. (1.69) leads to the one-electron GKS equations

$$\left(-\frac{1}{2} \nabla^2 + v_{\text{ne}}(\mathbf{r}) + v_{\text{H}}(\mathbf{r}) + v_{\bar{S}}(\mathbf{r}) \right) \varphi_{i\sigma}(\mathbf{r}) + \frac{\delta S[\Phi]}{\delta \varphi_{i\sigma}^*(\mathbf{r})} = \varepsilon_{i\sigma} \varphi_{i\sigma}(\mathbf{r}), \quad (1.70)$$

where $v_{\bar{S}}(\mathbf{r}) = \delta\bar{S}[n]/\delta n(\mathbf{r})$ is a local potential and $\delta S[\Phi]/\delta\varphi_{i\sigma}^*(\mathbf{r})$ generates a one-electron (possibly nonlocal) operator.

In the special case $S[\Phi] = 0$, we recover the KS exchange-correlation density functional

$$E_{\text{xc}}^{S=0}[\Phi] = E_{\text{xc}}[n_{\Phi}]. \quad (1.71)$$

Due to the freedom in the choice of $S[\Phi]$, there is an infinity of GKS exchange-correlation functionals $E_{\text{xc}}^S[\Phi]$ giving the exact ground-state energy via Eq. (1.69). This freedom and the fact that Φ carries more information than n_{Φ} gives the possibility to design more accurate approximations for the exchange-correlation energy.

Of course, by starting from the density functional $F[n_{\uparrow}, n_{\downarrow}]$ in Eq. (1.60), this GKS method can be extended to the spin-dependent case, leading to GKS exchange-correlation functionals of the form $E_{\text{xc}}^S[\Phi] = S[\Phi] + \bar{S}[n_{\uparrow, \Phi}, n_{\downarrow, \Phi}]$.

2 Exact expressions and constraints for the exchange and correlation functionals

2.1 The exchange and correlation functionals in terms of the exchange and correlation holes

2.1.1 The exchange and correlation holes

Let us consider the pair density associated with the wave function $\Psi[n]$ defined in Eq. (1.17)

$$n_2(\mathbf{r}_1, \mathbf{r}_2) = N(N-1) \int \cdots \int |\Psi[n](\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N)|^2 d\sigma_1 d\sigma_2 d\mathbf{x}_3 \dots d\mathbf{x}_N, \quad (2.1)$$

which is a functional of the density, and is normalized to the number of electron pairs, $\iint n_2(\mathbf{r}_1, \mathbf{r}_2) d\mathbf{r}_1 d\mathbf{r}_2 = N(N-1)$. It is proportional to the probability density of finding two electrons at positions $(\mathbf{r}_1, \mathbf{r}_2)$ with all the other electrons anywhere. The pair density is useful to express the expectation of the electron-electron interaction operator

$$\langle \Psi[n] | \hat{W}_{ee} | \Psi[n] \rangle = \frac{1}{2} \iint \frac{n_2(\mathbf{r}_1, \mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2. \quad (2.2)$$

Mirroring the decomposition of the Hartree-exchange-correlation energy performed in the KS method [Eq. (1.25)], the pair density can be decomposed as

$$n_2(\mathbf{r}_1, \mathbf{r}_2) = n(\mathbf{r}_1)n(\mathbf{r}_2) + n_{2,xc}(\mathbf{r}_1, \mathbf{r}_2). \quad (2.3)$$

The product of the densities $n(\mathbf{r}_1)n(\mathbf{r}_2)$ corresponds to the case of independent electrons and the exchange-correlation pair density $n_{2,xc}(\mathbf{r}_1, \mathbf{r}_2)$ represents the modification of the pair density due to exchange and correlation effects between the electrons. It can be further written as

$$n_{2,xc}(\mathbf{r}_1, \mathbf{r}_2) = n(\mathbf{r}_1)n_{xc}(\mathbf{r}_1, \mathbf{r}_2), \quad (2.4)$$

where $n_{xc}(\mathbf{r}_1, \mathbf{r}_2)$ is the *exchange-correlation hole*. It can be interpreted as the modification due to exchange and correlation effects of the conditional probability of finding an electron at \mathbf{r}_2 knowing that one has been found at \mathbf{r}_1 . The positivity of $n_2(\mathbf{r}_1, \mathbf{r}_2)$ implies that

$$n_{xc}(\mathbf{r}_1, \mathbf{r}_2) \geq -n(\mathbf{r}_2). \quad (2.5)$$

Moreover, we have the following sum rule

$$\int n_{xc}(\mathbf{r}_1, \mathbf{r}_2) d\mathbf{r}_2 = -1. \quad (2.6)$$

Exercise 6 : Prove the sum rule of Eq. (2.6).

We can separate the exchange and correlation contributions in the exchange-correlation hole. For this, consider the pair density $n_{2,KS}(\mathbf{r}_1, \mathbf{r}_2)$ associated with the KS single-determinant wave function $\Phi[n]$ defined in Eq. (1.23). It can be decomposed as

$$n_{2,KS}(\mathbf{r}_1, \mathbf{r}_2) = n(\mathbf{r}_1)n(\mathbf{r}_2) + n_{2,x}(\mathbf{r}_1, \mathbf{r}_2), \quad (2.7)$$

where $n_{2,x}(\mathbf{r}_1, \mathbf{r}_2)$ is the exchange pair density, which is further written as

$$n_{2,x}(\mathbf{r}_1, \mathbf{r}_2) = n(\mathbf{r}_1)n_x(\mathbf{r}_1, \mathbf{r}_2), \quad (2.8)$$

where $n_x(\mathbf{r}_1, \mathbf{r}_2)$ is the *exchange hole*. Just like the exchange-correlation hole, the exchange hole satisfies the conditions

$$n_x(\mathbf{r}_1, \mathbf{r}_2) \geq -n(\mathbf{r}_2), \quad (2.9)$$

and

$$\int n_x(\mathbf{r}_1, \mathbf{r}_2) d\mathbf{r}_2 = -1. \quad (2.10)$$

Moreover, by writing the exchange hole in terms of the one-electron KS density matrix $n_{1,\sigma}(\mathbf{r}_1, \mathbf{r}_2) = \sum_{j=1}^{N_\sigma} \varphi_{j\sigma}^*(\mathbf{r}_2)\varphi_{j\sigma}(\mathbf{r}_1)$,

$$n_x(\mathbf{r}_1, \mathbf{r}_2) = - \sum_{\sigma \in \{\uparrow, \downarrow\}} |n_{1,\sigma}(\mathbf{r}_1, \mathbf{r}_2)|^2 / n(\mathbf{r}_1), \quad (2.11)$$

we see that it is always negative

$$n_x(\mathbf{r}_1, \mathbf{r}_2) \leq 0. \quad (2.12)$$

From Eqs. (1.28), (2.2), (2.7), and (2.8), it can be seen that the exchange energy functional can be written in terms of the exchange hole

$$E_x[n] = \frac{1}{2} \iint \frac{n(\mathbf{r}_1)n_x(\mathbf{r}_1, \mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2, \quad (2.13)$$

leading to the interpretation of E_x as the electrostatic interaction energy of an electron and its exchange hole. It is also useful to write the exchange energy functional as

$$E_x[n] = \int n(\mathbf{r}_1)\varepsilon_x[n](\mathbf{r}_1) d\mathbf{r}_1, \quad (2.14)$$

where $\varepsilon_x[n](\mathbf{r}_1)$ is the exchange energy per particle

$$\varepsilon_x[n](\mathbf{r}_1) = \frac{1}{2} \int \frac{n_x(\mathbf{r}_1, \mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_2, \quad (2.15)$$

which is itself a functional of the density. In approximate exchange density functionals, the quantity $\varepsilon_x[n](\mathbf{r}_1)$ is usually what is approximated.

Exercise 7 : Show that, for finite systems, $\varepsilon_x[n](\mathbf{r}) \underset{r \rightarrow +\infty}{\sim} -1/(2r)$.

The *correlation hole* is defined as the difference

$$n_c(\mathbf{r}_1, \mathbf{r}_2) = n_{xc}(\mathbf{r}_1, \mathbf{r}_2) - n_x(\mathbf{r}_1, \mathbf{r}_2), \quad (2.16)$$

and, from Eqs. (2.6) and (2.10), satisfies the sum rule

$$\int n_c(\mathbf{r}_1, \mathbf{r}_2) d\mathbf{r}_2 = 0, \quad (2.17)$$

which implies that the correlation hole has negative and positive contributions⁵. The potential contribution to the correlation energy can be written in terms of the correlation hole

$$U_c[n] = \frac{1}{2} \iint \frac{n(\mathbf{r}_1)n_c(\mathbf{r}_1, \mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2. \quad (2.18)$$

In order to express the total correlation energy $E_c[n] = T_c[n] + U_c[n]$ in a form similar to Eq. (2.18), we need to introduce the adiabatic-connection formalism.

⁵Therefore, the correlation hole is really a “hole” only in some region of space, and a “bump” in other regions.

2.1.2 The adiabatic connection

The idea of the *adiabatic connection* is to have a continuous path between the non-interacting KS system and the physical system while keeping the ground-state density constant. An infinity of such paths are possible, but the one most often considered consists in switching on the electron-electron interaction linearly with a coupling constant $\lambda \in \mathbb{R}$. The Hamiltonian along this adiabatic connection is

$$\hat{H}^\lambda = \hat{T} + \lambda \hat{W}_{ee} + \hat{V}^\lambda, \quad (2.19)$$

where \hat{V}^λ is the external local potential operator imposing that the ground-state density is the same as the ground-state density of the physical system for all λ . The Hamiltonian (2.19) reduces to the KS non-interacting Hamiltonian for $\lambda = 0$ and to the physical Hamiltonian for $\lambda = 1$.

Just as for the physical system, it is possible to define a universal functional associated with the system of Eq. (2.19) for each value of the parameter λ

$$F^\lambda[n] = \min_{\substack{\Psi \in \mathcal{W} \\ \Psi \rightarrow n}} \langle \Psi | \hat{T} + \lambda \hat{W}_{ee} | \Psi \rangle = \langle \Psi^\lambda[n] | \hat{T} + \lambda \hat{W}_{ee} | \Psi^\lambda[n] \rangle, \quad (2.20)$$

the minimizing wave function being denoted by $\Psi^\lambda[n]$. Clearly, for $\lambda = 1$ we recover the physical universal functional, $F^{\lambda=1}[n] = F[n]$, while for $\lambda = 0$, assuming that the KS wave function is non-degenerate, we recover the non-interacting kinetic energy functional $F^{\lambda=0}[n] = T_s[n]$. The functional $F^\lambda[n]$ can be decomposed as

$$F^\lambda[n] = T_s[n] + E_H^\lambda[n] + E_{xc}^\lambda[n], \quad (2.21)$$

where $E_H^\lambda[n]$ is the Hartree energy functional associated with the interaction $\lambda \hat{W}_{ee}$ and is simply linear in λ

$$E_H^\lambda[n] = \frac{1}{2} \iint n(\mathbf{r}_1)n(\mathbf{r}_2) \frac{\lambda}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2 = \lambda E_H[n], \quad (2.22)$$

and $E_{xc}^\lambda[n]$ is a remaining exchange-correlation functional. It can be decomposed as a sum of an exchange contribution, which is also linear in λ ,

$$E_x^\lambda[n] = \langle \Phi[n] | \lambda \hat{W}_{ee} | \Phi[n] \rangle - E_H^\lambda[n] = \lambda E_x[n], \quad (2.23)$$

and a correlation contribution, which is nonlinear in λ ,

$$E_c^\lambda[n] = \langle \Psi^\lambda[n] | \hat{T} + \lambda \hat{W}_{ee} | \Psi^\lambda[n] \rangle - \langle \Phi[n] | \hat{T} + \lambda \hat{W}_{ee} | \Phi[n] \rangle. \quad (2.24)$$

Taking the derivative of Eq. (2.24) with respect to λ and using the Hellmann-Feynman theorem for the wave function $\Psi^\lambda[n]$ ⁶, we obtain

$$\frac{\partial E_c^\lambda[n]}{\partial \lambda} = \langle \Psi^\lambda[n] | \hat{W}_{ee} | \Psi^\lambda[n] \rangle - \langle \Phi[n] | \hat{W}_{ee} | \Phi[n] \rangle, \quad (2.25)$$

Integrating over λ from 0 to 1, and using $E_c^{\lambda=1}[n] = E_c[n]$ and $E_c^{\lambda=0}[n] = 0$, we arrive at the *adiabatic-connection formula* for the correlation energy functional of the physical system

$$E_c[n] = \int_0^1 d\lambda \langle \Psi^\lambda[n] | \hat{W}_{ee} | \Psi^\lambda[n] \rangle - \langle \Phi[n] | \hat{W}_{ee} | \Phi[n] \rangle. \quad (2.26)$$

⁶In this context, the Hellmann-Feynman theorem states that in the derivative $\frac{\partial F^\lambda[n]}{\partial \lambda} = \langle \frac{\partial \Psi^\lambda[n]}{\partial \lambda} | \hat{T} + \lambda \hat{W}_{ee} | \Psi^\lambda[n] \rangle + \langle \Psi^\lambda[n] | \hat{W}_{ee} | \frac{\partial \Psi^\lambda[n]}{\partial \lambda} \rangle + \langle \Psi^\lambda[n] | \hat{T} + \lambda \hat{W}_{ee} | \frac{\partial \Psi^\lambda[n]}{\partial \lambda} \rangle$ the first and third terms involving the derivative of $\Psi^\lambda[n]$ vanish. This is due to the fact that $\Psi^\lambda[n]$ is obtained via the minimization of Eq. (2.20) and thus any variation of $\Psi^\lambda[n]$ which keeps the density constant (which is the case for a variation with respect to λ) gives a vanishing variation of $F^\lambda[n]$.

By introducing the correlation hole $n_c^\lambda(\mathbf{r}_1, \mathbf{r}_2)$ associated to the wave function $\Psi^\lambda[n]$, the adiabatic-connection formula for the correlation energy can also be written as

$$E_c[n] = \frac{1}{2} \int_0^1 d\lambda \iint \frac{n(\mathbf{r}_1)n_c^\lambda(\mathbf{r}_1, \mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2, \quad (2.27)$$

or, noting that $n_c^\lambda(\mathbf{r}_1, \mathbf{r}_2)$ is the only quantity that depends on λ in Eq. (2.27), in a more compact way,

$$E_c[n] = \frac{1}{2} \iint \frac{n(\mathbf{r}_1)\bar{n}_c(\mathbf{r}_1, \mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2, \quad (2.28)$$

where $\bar{n}_c(\mathbf{r}_1, \mathbf{r}_2) = \int_0^1 d\lambda n_c^\lambda(\mathbf{r}_1, \mathbf{r}_2)$ is the coupling-constant-integrated correlation hole. It leads to the interpretation of E_c as the electrostatic interaction energy of an electron with its coupling-constant-integrated correlation hole. As for the exchange energy, the correlation energy functional can be written as

$$E_c[n] = \int n(\mathbf{r}_1)\varepsilon_c[n](\mathbf{r}_1)d\mathbf{r}_1, \quad (2.29)$$

where $\varepsilon_c[n](\mathbf{r}_1)$ is the correlation energy per particle

$$\varepsilon_c[n](\mathbf{r}_1) = \frac{1}{2} \int \frac{\bar{n}_c(\mathbf{r}_1, \mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_2, \quad (2.30)$$

which is a functional of the density that needs to be approximated.

2.2 Uniform coordinate scaling

We consider a norm-preserving uniform scaling of the spatial coordinates in the N -electron wave function along the adiabatic connection $\Psi^\lambda[n]$ [introduced in Eq. (2.20)] while leaving untouched the spin coordinates [30–32]

$$\Psi_\gamma^\lambda[n](\mathbf{r}_1, \sigma_1, \dots, \mathbf{r}_N, \sigma_N) = \gamma^{3N/2} \Psi^\lambda[n](\gamma\mathbf{r}_1, \sigma_1, \dots, \gamma\mathbf{r}_N, \sigma_N), \quad (2.31)$$

where $\gamma > 0$ is a scaling factor. The scaled wave function $\Psi_\gamma^\lambda[n]$ yields the scaled density

$$n_\gamma(\mathbf{r}) = \gamma^3 n(\gamma\mathbf{r}), \quad (2.32)$$

with $\int n_\gamma(\mathbf{r})d\mathbf{r} = \int n(\mathbf{r})d\mathbf{r} = N$ and minimizes $\langle \Psi | \hat{T} + \lambda\gamma\hat{W}_{ee} | \Psi \rangle$ since it can be shown that

$$\langle \Psi_\gamma^\lambda[n] | \hat{T} + \lambda\gamma\hat{W}_{ee} | \Psi_\gamma^\lambda[n] \rangle = \gamma^2 \langle \Psi^\lambda[n] | \hat{T} + \lambda\hat{W}_{ee} | \Psi^\lambda[n] \rangle. \quad (2.33)$$

Exercise 8 : Prove the identity of Eq. (2.33).

We thus conclude that the scaled wave function at the density n and coupling constant λ corresponds to the wave function at the scaled density n_γ and coupling constant $\lambda\gamma$

$$\Psi_\gamma^\lambda[n] = \Psi^{\lambda\gamma}[n_\gamma], \quad (2.34)$$

or, equivalently,

$$\Psi_\gamma^{\lambda/\gamma}[n] = \Psi^\lambda[n_\gamma], \quad (2.35)$$

and that the universal density functional satisfies the scaling relation

$$F^{\lambda\gamma}[n_\gamma] = \gamma^2 F^\lambda[n], \quad (2.36)$$

or, equivalently,

$$F^\lambda[n_\gamma] = \gamma^2 F^{\lambda/\gamma}[n]. \quad (2.37)$$

At $\lambda = 0$, we find the scaling relation of the KS wave function $\Phi[n]$

$$\Phi[n_\gamma] = \Phi_\gamma[n]. \quad (2.38)$$

This directly leads to the scaling relation for the non-interacting kinetic density functional [see Eq. (1.23)]

$$T_s[n_\gamma] = \gamma^2 T_s[n]. \quad (2.39)$$

Moreover, since the Hartree density functional [see Eq. (1.26)] scales linearly

$$E_H[n_\gamma] = \gamma E_H[n], \quad (2.40)$$

Eq. (2.38) implies that the exchange density functional [see Eq. (1.28)] also scales linearly

$$E_x[n_\gamma] = \gamma E_x[n]. \quad (2.41)$$

Exercise 9 : Prove the scaling relations satisfied by the Hartree and exchange density functionals in Eqs. (2.40) and (2.41).

However, the correlation density functional $E_c[n]$ has the more complicated scaling (as $F[n]$)

$$E_c^\lambda[n_\gamma] = \gamma^2 E_c^{\lambda/\gamma}[n], \quad (2.42)$$

and, in particular for $\lambda = 1$,

$$E_c[n_\gamma] = \gamma^2 E_c^{1/\gamma}[n]. \quad (2.43)$$

These scaling relations allow one to find the behavior of the density functionals in the high- and low-density limits. In the *high-density limit* ($\gamma \rightarrow \infty$), it can be shown from Eq. (2.43) that, for nondegenerate KS systems, the correlation functional $E_c[n]$ goes to a constant

$$\lim_{\gamma \rightarrow \infty} E_c[n_\gamma] = E_c^{\text{GL2}}[n], \quad (2.44)$$

where $E_c^{\text{GL2}}[n]$ is the second-order Görling-Levy (GL2) correlation energy [33, 34] (see Section 4.2). This is also called the *weak-correlation limit* since in this limit the correlation energy is negligible with respect to exchange energy which is itself negligible with respect to the non-interacting kinetic energy: $|E_c[n_\gamma]| = O(\gamma^0) \ll |E_x[n_\gamma]| = O(\gamma) \ll T_s[n_\gamma] = O(\gamma^2)$. Equation (2.44) is an important constraint since atomic and molecular correlation energies are often close to the high-density limit. For example, for the ground-state density of the helium atom, we have $E_c[n] = -0.0421$ hartree and $\lim_{\gamma \rightarrow \infty} E_c[n_\gamma] = -0.0467$ hartree [35].

In the *low-density limit* ($\gamma \rightarrow 0$), it can be shown from Eq. (2.37) that the Hartree-exchange-correlation $E_{\text{Hxc}}[n]$ goes to zero linearly in γ

$$E_{\text{Hxc}}[n_\gamma] \underset{\gamma \rightarrow 0}{\sim} \gamma W_{\text{ee}}^{\text{SCE}}[n], \quad (2.45)$$

where $W_{\text{ee}}^{\text{SCE}}[n] = \inf_{\Psi \in \mathcal{W}, \Psi \rightarrow n} \langle \Psi | \hat{W}_{\text{ee}} | \Psi \rangle$ is the strictly-correlated-electron (SCE) functional [36–39]. This is also called the *strong-interaction limit* since in this limit the Hartree-exchange-correlation energy dominates over the non-interacting kinetic energy: $E_{\text{Hxc}}[n_\gamma] = O(\gamma) \gg T_s[n_\gamma] = O(\gamma^2)$. In this limit, the electrons strictly localize relatively to each other. This corresponds to a Wigner crystallization.

2.3 One-orbital spatial regions and self-interaction

For systems composed of only one spin- \uparrow (or, symmetrically, one spin- \downarrow) electron (e.g., the hydrogen atom) with density $n_{1e}(\mathbf{r}) = |\varphi_{1\uparrow}(\mathbf{r})|^2$ where $\varphi_{1\uparrow}(\mathbf{r})$ is the unique occupied KS orbital, the exchange hole in Eq. (2.11) simplifies to $n_x(\mathbf{r}_1, \mathbf{r}_2) = -n(\mathbf{r}_2)$, and consequently the exchange energy cancels out the Hartree energy

$$E_x[n_{1e}] = -E_H[n_{1e}]. \quad (2.46)$$

Besides, the correlation energy vanishes

$$E_c[n_{1e}] = 0. \quad (2.47)$$

This is of course also true for the spin-dependent version of the functionals introduced in Section 1.3.4, i.e.

$$E_x[n_{1e}, 0] = -E_H[n_{1e}]. \quad (2.48)$$

and

$$E_c[n_{1e}, 0] = 0. \quad (2.49)$$

For systems composed of two opposite-spin electrons (e.g., the helium atom or the dihydrogen molecule) in a unique doubly occupied KS orbital $\varphi_1(\mathbf{r}) = \varphi_{1\uparrow}(\mathbf{r}) = \varphi_{1\downarrow}(\mathbf{r})$ with density $n_{2e}^{\uparrow\downarrow}(\mathbf{r}) = 2|\varphi_1(\mathbf{r})|^2$, the exchange hole simplifies to $n_x(\mathbf{r}_1, \mathbf{r}_2) = -n(\mathbf{r}_2)/2$, and consequently the exchange energy is equal to half the opposite of the Hartree energy

$$E_x[n_{2e}^{\uparrow\downarrow}] = -\frac{1}{2}E_H[n_{2e}^{\uparrow\downarrow}]. \quad (2.50)$$

These are constraints for the exchange and correlation density functionals in the special cases $N = 1$ and $N = 2$. For systems with more electrons, similar relations apply locally in spatial regions where, among the occupied KS orbitals, only one orbital is not zero (or, more generally, takes non-negligible values) (see Ref. [17]). This situation can be approximately realized in chemical systems. For example, this approximately corresponds to an unpaired electron in a radical, and to an electron pair in a single covalent bond, in a lone pair, or in a core orbital. If approximate exchange and correlation density functionals do not satisfy these constraints, we say that they introduce a *self-interaction error*.

2.4 Lieb-Oxford lower bound

Lieb and Oxford derived a lower bound for the indirect Coulomb energy (i.e., the two-particle Coulomb potential energy beyond the Hartree energy) [40], which, when expressed in terms of the exchange or exchange-correlation functional, takes the form [41]

$$E_x[n] \geq E_{xc}[n] \geq -C_{LO} \int n(\mathbf{r})^{4/3} d\mathbf{r}, \quad (2.51)$$

where the optimal (i.e., smallest) constant C_{LO} (independent of the electron number N) has been narrowed to $1.4442 \leq C_{LO} \leq 1.5765$. This bound is approached in the low-density limit. For one-electron densities and opposite-spin two-electron densities, specific tighter bounds (i.e., with smaller C_{LO}) are known.

2.5 Fractional electron numbers and frontier orbital energies

In 1982, Perdew, Parr, Levy, and Balduz [42] extended DFT to fractional electron numbers. Although systems with a noninteger number of electrons may appear as unphysical, such systems in fact naturally arise in quantum mechanics, e.g. as fragments from a molecular dissociation in entangled quantum states. One important result of this extension is that the frontier KS orbital energies can be seen as derivatives of the total energy with respect to the electron number.

2.5.1 Quantum mechanics with fractional electron numbers

The ground-state energy of a system with a fixed fractional number of electrons $\mathcal{N} = N - 1 + f$ (where N is an integer and $0 \leq f \leq 1$) can be defined in an ensemble formalism as

$$E_0^{N-1+f} = \min_{\hat{\Gamma} \in \mathcal{D}_{\text{DM}}^{N-1+f}} \text{Tr} \left[\left(\hat{T} + \hat{W}_{\text{ee}} + \hat{V}_{\text{ne}} \right) \hat{\Gamma} \right], \quad (2.52)$$

where Tr denotes the trace (in Fock space) and the minimization is over ensemble density matrices $\hat{\Gamma}$ in the set $\mathcal{D}_{\text{DM}}^{N-1+f}$

$$\mathcal{D}_{\text{DM}}^{N-1+f} = \left\{ \hat{\Gamma} = (1-f)|\Psi^{N-1}\rangle\langle\Psi^{N-1}| + f|\Psi^N\rangle\langle\Psi^N|, \Psi^{N-1} \in \mathcal{W}^{N-1}, \Psi^N \in \mathcal{W}^N \right\}, \quad (2.53)$$

where f is fixed, and Ψ^{N-1} and Ψ^N are arbitrary wave functions in the $(N-1)$ - and N -electron admissible wave-function sets \mathcal{W}^{N-1} and \mathcal{W}^N , respectively. This form of $\hat{\Gamma}$ in Eq. (2.53) ensures an average electron number of $N - 1 + f$.⁷ The minimizing ensemble density matrix is

$$\hat{\Gamma}_0 = (1-f)|\Psi_0^{N-1}\rangle\langle\Psi_0^{N-1}| + f|\Psi_0^N\rangle\langle\Psi_0^N|, \quad (2.54)$$

where Ψ_0^{N-1} and Ψ_0^N are ground-state wave functions of the $(N-1)$ - and N -electron systems, respectively.

The fact that the minimizing ensemble density matrix is linear in f implies that the ground-state energy is also linear in f between the integer electron numbers $N-1$ and N

$$\boxed{E_0^{N-1+f} = (1-f)E_0^{N-1} + fE_0^N}, \quad (2.55)$$

where E_0^{N-1} and E_0^N are the ground-state energies of the $(N-1)$ - and N -electron systems, respectively. Similarly, between the integer electron numbers N and $N+1$, we have

$$\boxed{E_0^{N+f} = (1-f)E_0^N + fE_0^{N+1}}. \quad (2.56)$$

Thus, the ground-state energy is a *continuous piecewise linear function* of the fractional electron number \mathcal{N} .

Exercise 10 : Prove that the minimizing ensemble density matrix is indeed given by Eq. (2.54) and the ground-state energy by Eq. (2.55).

⁷In fact, with the assumption that the ground-state energy for integer electron numbers in a fixed external potential is a convex function, $E^N \leq (E^{N+1} + E^{N-1})/2$, which is true for any known realistic system, the minimization in Eq. (2.52) can be done over ensemble density matrices $\hat{\Gamma}$ constructed from wave functions with any number of electrons with the only constraint that $\hat{\Gamma}$ gives an average electron number of $N - 1 + f$.

The derivative of $E_0^{\mathcal{N}}$ with respect to \mathcal{N} defines the *electronic chemical potential*⁸

$$\mu = \frac{\partial E_0^{\mathcal{N}}}{\partial \mathcal{N}}, \quad (2.57)$$

and is obtained as the derivative with respect to f of Eq. (2.55) for $N - 1 < \mathcal{N} < N$

$$\left(\frac{\partial E_0^{\mathcal{N}}}{\partial \mathcal{N}} \right)_{N-1 < \mathcal{N} < N} = E_0^N - E_0^{N-1} = -I_N, \quad (2.58)$$

where I_N is the ionization energy of the N electron system (I_N is always strictly positive), and similarly from Eq. (2.56) for $N < \mathcal{N} < N + 1$

$$\left(\frac{\partial E_0^{\mathcal{N}}}{\partial \mathcal{N}} \right)_{N < \mathcal{N} < N+1} = E_0^{N+1} - E_0^N = -A_N, \quad (2.59)$$

where A_N is the electron affinity of the N electron system (with this definition, A_N is strictly positive if the $(N + 1)$ -electron system is bound, otherwise it is zero). The electronic chemical potential μ has thus a discontinuity at the integer electron number N . So, the plot of $E_0^{\mathcal{N}}$ with respect to \mathcal{N} is made of a series of straight lines between integer electron numbers, with derivative discontinuities at each integer.

2.5.2 Density-functional theory with fractional electron numbers

The universal density functional $F[n]$ is extended to densities integrating to a fractional electron number, $\int n(\mathbf{r})d\mathbf{r} = \mathcal{N} = N - 1 + f$, as

$$F[n] = \min_{\substack{\hat{\Gamma} \in \mathcal{D}_{\text{DM}}^{N-1+f} \\ \hat{\Gamma} \rightarrow n}} \text{Tr} \left[\left(\hat{T} + \hat{W}_{\text{ee}} \right) \hat{\Gamma} \right], \quad (2.60)$$

where the minimization is performed over ensemble density matrices $\hat{\Gamma} \in \mathcal{D}_{\text{DM}}^{N-1+f}$ and yielding the density n . As usual, to set up a KS method, $F[n]$ is decomposed as

$$F[n] = T_{\text{s}}[n] + E_{\text{Hxc}}[n], \quad (2.61)$$

with a KS non-interacting kinetic-energy functional and a remaining Hartree-exchange-correlation functional $E_{\text{Hxc}}[n]$. Here, $T_{\text{s}}[n]$ is defined as

$$T_{\text{s}}[n] = \min_{\substack{\hat{\Gamma}_{\text{s}} \in \mathcal{D}_{\text{DM,s}}^{N-1+f} \\ \hat{\Gamma}_{\text{s}} \rightarrow n}} \text{Tr}[\hat{\Gamma}_{\text{s}} \hat{T}], \quad (2.62)$$

where the minimization is over ensemble non-interacting density matrices $\hat{\Gamma}_{\text{s}} \in \mathcal{D}_{\text{DM,s}}^{N-1+f}$ and yielding the density n . The set of non-interacting density matrices is where the minimization is over ensemble non-interacting density matrices $\hat{\Gamma}_{\text{s}}$ of the form

$$\mathcal{D}_{\text{DM,s}}^{N-1+f} = \left\{ \hat{\Gamma}_{\text{s}} = (1-f)|\Phi^{N-1,f}\rangle\langle\Phi^{N-1,f}| + f|\Phi^{N,f}\rangle\langle\Phi^{N,f}|, \Phi^{N-1,f} = \psi_1 \wedge \dots \wedge \psi_{N-1}, \right. \\ \left. \Phi^{N,f} = \psi_1 \wedge \dots \wedge \psi_N, \psi_i \in H^1(\mathbb{R}^3 \times \{\uparrow, \downarrow\}, \mathbb{C}), \langle\psi_i|\psi_j\rangle = \delta_{i,j} \right\}, \quad (2.63)$$

⁸The electronic chemical potential is an important quantity in the field of *conceptual DFT*. The electronegativity of a system can be defined as the opposite of the electronic chemical potential

i.e., $\Phi^{N-1,f}$ and $\Phi^{N,f}$ are $(N-1)$ - and N -electron single-determinant wave functions, respectively, constructed from a common set of spin orbitals $\{\psi_i\}$ depending on the fixed f . The exact ground-state energy can then be expressed as

$$E_0^{N-1+f} = \min_{\hat{\Gamma}_s \in \mathcal{D}_{\text{DM},s}^{N-1+f}} \left\{ \text{Tr} \left[\left(\hat{T} + \hat{V}_{\text{ne}} \right) \hat{\Gamma}_s \right] + E_{\text{Hxc}}[n_{\hat{\Gamma}_s}] \right\}, \quad (2.64)$$

In Eq. (2.64), E_{Hxc} is evaluated at $n_{\hat{\Gamma}_s}$, i.e. the density of $\hat{\Gamma}_s$. The total electronic energy can then be written in terms of these orbitals and occupation numbers n_i

$$E = \sum_{i=1}^N n_i \int \varphi_i^*(\mathbf{r}) \left(-\frac{1}{2} \nabla^2 + v_{\text{ne}}(\mathbf{r}) \right) \varphi_i(\mathbf{r}) \text{d}\mathbf{r} + E_{\text{Hxc}}[n], \quad (2.65)$$

with the density

$$n(\mathbf{r}) = \sum_{i=1}^N n_i |\varphi_i(\mathbf{r})|^2, \quad (2.66)$$

where $n_i = 1$ for $i \leq N-1$ and $n_N = f$ for the highest occupied molecular orbital (HOMO) which is assumed here to be nondegenerate for simplicity. As in the standard KS method, the variation of Eq. (2.65) with respect to the orthonormal orbitals but with fixed occupation numbers leads to the self-consistent KS equations

$$\left(-\frac{1}{2} \nabla^2 + v_s(\mathbf{r}) \right) \varphi_i(\mathbf{r}) = \varepsilon_i \varphi_i(\mathbf{r}), \quad (2.67)$$

where $v_s(\mathbf{r})$ is the KS potential

$$v_s(\mathbf{r}) = v_{\text{ne}}(\mathbf{r}) + \frac{\delta E_{\text{Hxc}}[n]}{\delta n(\mathbf{r})}. \quad (2.68)$$

Thus, the KS equations for fractional electron numbers look identical (with the caveat of a fractional occupation number for the HOMO) to the standard KS equations for integer electron numbers. There is however one important difference. In the standard KS equations, the functional derivative of $E_{\text{Hxc}}[n]$ is taken with respect to density variations $\delta n(\mathbf{r})$ keeping the number of electrons constant, i.e. $\int \delta n(\mathbf{r}) \text{d}\mathbf{r} = 0$. As a consequence, the functional derivative is defined only up to an additive constant, which can be seen from its definition

$$\delta E_{\text{Hxc}}[n] = \int \left(\frac{\delta E_{\text{Hxc}}[n]}{\delta n(\mathbf{r})} + \text{const} \right) \delta n(\mathbf{r}) \text{d}\mathbf{r}. \quad (2.69)$$

In the present extension to fractional electron numbers, since the functional $E_{\text{Hxc}}[n]$ is now defined for densities having any noninteger electron numbers, its functional derivative can now be generally taken with respect to density variations which can change the number of electrons, i.e. $\int \delta n(\mathbf{r}) \text{d}\mathbf{r} \neq 0$. Consequently, the constant in Eq. (2.69) is now determined. This unambiguously fixes the values of the KS orbital energies.

Eqs. (2.65)-(2.68) constitutes a general KS method with fractional occupation numbers. A useful result is that, after optimizing the orbitals with fixed occupation numbers, the derivative of the total energy with respect to the occupation number n_i of an occupied orbital equals the energy ε_i of this orbital,

$$\frac{\partial E}{\partial n_i} = \varepsilon_i, \quad (2.70)$$

which is known as *Janak's theorem* [43].

Exercise 11 : Prove Janak's theorem.

For clarity in the discussion, we will now explicitly indicate the dependence on the electron number \mathcal{N} in the density $n^{\mathcal{N}}(\mathbf{r})$, the KS potential $v_s^{\mathcal{N}}(\mathbf{r})$, the KS orbitals $\varphi_i^{\mathcal{N}}(\mathbf{r})$, and the KS orbital energies $\varepsilon_i^{\mathcal{N}}$.

2.5.3 The HOMO energy and the ionization energy

Applying Janak's theorem for the case of the HOMO for a fractional electron number $\mathcal{N} = N - \delta$ where $\delta \rightarrow 0^+$ (the limit will be always tacitly assumed) gives

$$\left(\frac{\partial E_0^{\mathcal{N}}}{\partial \mathcal{N}} \right)_{N-\delta} = \varepsilon_{\text{H}}^{N-\delta} = \varepsilon_{\text{H}}^N, \quad (2.71)$$

where ε_{H}^N is the HOMO energy of the N -electron system (here, the HOMO is the N^{th} orbital). Notice that, since $\partial E_0^{\mathcal{N}}/\partial \mathcal{N}$ has a discontinuity at the integer electron number N , it is important to specify that ε_{H}^N is defined as the limit when N is approached from the left (electron-deficient) side. Combining this result with Eq. (2.58) leads to

$$\boxed{\varepsilon_{\text{H}}^N = -I_N}, \quad (2.72)$$

i.e., the energy of the HOMO KS orbital is the opposite of the exact ionization energy.

This can be used, together with the asymptotic behavior of the density, to determine the constant in the KS potential. Indeed, for finite systems, it can be shown that the exact ground-state density of the N -electron system decays exponentially for $r = |\mathbf{r}| \rightarrow +\infty$ with an exponent related to the ionization energy I_N [44]

$$n^N(\mathbf{r}) \underset{r \rightarrow +\infty}{\propto} e^{-2\sqrt{2I_N} r}. \quad (2.73)$$

Besides, each occupied KS orbitals of the N -electron system has the following asymptotic behavior (see Exercise 12)

$$\varphi_i^N(\mathbf{r}) \underset{r \rightarrow +\infty}{\propto} e^{-\sqrt{-2(\varepsilon_i^N - v_s^N(\infty))} r}, \quad (2.74)$$

where $v_s^N(\infty) = v_s^{N-\delta}(\infty)$ is the asymptotic value of the KS potential (defined as the limit from the left side) which is for now unknown but fixed since the additive constant in the KS potential is determined now according to the discussion after Eq. (2.69). Consequently, the HOMO KS orbital $\varphi_{\text{H}}^N(\mathbf{r})$ is the slowest decaying orbital and dominates the asymptotic decay of the KS density [via Eq. (2.66)]

$$n^N(\mathbf{r}) \underset{r \rightarrow +\infty}{\propto} e^{-2\sqrt{-2(\varepsilon_{\text{H}}^N - v_s^N(\infty))} r}. \quad (2.75)$$

Since the KS density is the exact density, Eqs. (2.73) and (2.75) must agree, and since we have already shown that $\varepsilon_{\text{H}}^N = -I_N$, we thus find

$$v_s^N(\infty) = 0, \quad (2.76)$$

in accordance with the choice usually made in the standard KS method for integer electron numbers.

Exercise 12 : Check that the asymptotic decay of the KS orbitals is indeed given by Eq. (2.74). For this, use Eq. (2.67) and the asymptotic behavior of the KS potential $v_s^N(\mathbf{r}) \underset{r \rightarrow +\infty}{\sim} v_s^N(\infty) + (Q - 1)/r$ where Q is the total charge of the system.

2.5.4 The LUMO energy, the electron affinity, and the derivative discontinuity

Applying Janak's theorem for the HOMO but now for a fractional electron number $\mathcal{N} = N + \delta$ where $\delta \rightarrow 0^+$ gives

$$\left(\frac{\partial E_0^{\mathcal{N}}}{\partial \mathcal{N}} \right)_{N+\delta} = \varepsilon_{\text{H}}^{N+\delta}, \quad (2.77)$$

where $\varepsilon_{\text{H}}^{N+\delta}$ is the HOMO energy from the right (excess-electron) side of the discontinuity (here, the HOMO is the $(N+1)^{\text{th}}$ orbital). Since the derivative is in fact the same for all $N < \mathcal{N} < N+1$, it is also the HOMO energy of the $(N+1)$ -electron system, $\varepsilon_{\text{H}}^{N+\delta} = \varepsilon_{\text{H}}^{N+1-\delta} = \varepsilon_{\text{H}}^{N+1}$. Combining Eq. (2.77) with Eq. (2.59) leads to

$$\varepsilon_{\text{H}}^{N+\delta} = -A_N. \quad (2.78)$$

Naively, one may think that $\varepsilon_{\text{H}}^{N+\delta}$ is equal to the LUMO energy of the N -electron system ε_{L}^N (again defined as the limit from the left side, $\varepsilon_{\text{L}}^N = \varepsilon_{\text{L}}^{N-\delta}$, i.e. the $(N+1)^{\text{th}}$ orbital), and therefore that the LUMO KS energy equals to the opposite of the electron affinity. However, this is not as simple. Let us compare $\varepsilon_{\text{H}}^{N+\delta}$ that we can write as

$$\varepsilon_{\text{H}}^{N+\delta} = \int \varphi_{\text{H}}^{N+\delta}(\mathbf{r})^* \left(-\frac{1}{2} \nabla^2 + v_s^{N+\delta}(\mathbf{r}) \right) \varphi_{\text{H}}^{N+\delta}(\mathbf{r}) \text{d}\mathbf{r}, \quad (2.79)$$

with $\varepsilon_{\text{L}}^{N-\delta}$ that we can write as

$$\varepsilon_{\text{L}}^{N-\delta} = \int \varphi_{\text{L}}^{N-\delta}(\mathbf{r})^* \left(-\frac{1}{2} \nabla^2 + v_s^{N-\delta}(\mathbf{r}) \right) \varphi_{\text{L}}^{N-\delta}(\mathbf{r}) \text{d}\mathbf{r}. \quad (2.80)$$

The problem is that there is nothing preventing the KS potential to have a discontinuity $v_s^{N+\delta}(\mathbf{r}) \neq v_s^{N-\delta}(\mathbf{r})$. Indeed, the continuity of the density implies that $n^{N+\delta}(\mathbf{r}) = n^{N-\delta}(\mathbf{r})$,⁹ but this only imposes that $v_s^{N+\delta}(\mathbf{r})$ and $v_s^{N-\delta}(\mathbf{r})$ be equal up to an additive spatial constant (according to the Hohenberg-Kohn theorem). So we can have:

$$v_s^{N+\delta}(\mathbf{r}) - v_s^{N-\delta}(\mathbf{r}) = \Delta_{\text{xc}}^N, \quad (2.81)$$

where Δ_{xc}^N is independent from \mathbf{r} . Since the two potentials just differ by an additive constant, the orbitals are continuous at the integer N , and in particular $\varphi_{\text{H}}^{N+\delta}(\mathbf{r}) = \varphi_{\text{L}}^{N-\delta}(\mathbf{r})$. Using this fact and Eq. (2.81), we find

$$\begin{aligned} \varepsilon_{\text{H}}^{N+\delta} &= \int \varphi_{\text{L}}^{N-\delta}(\mathbf{r})^* \left(-\frac{1}{2} \nabla^2 + v_s^{N-\delta}(\mathbf{r}) \right) \varphi_{\text{L}}^{N-\delta}(\mathbf{r}) \text{d}\mathbf{r} + \Delta_{\text{xc}}^N \\ &= \varepsilon_{\text{L}}^{N-\delta} + \Delta_{\text{xc}}^N. \end{aligned} \quad (2.82)$$

In conclusion, the LUMO energy of the N -electron system is not the opposite of the exact electron affinity

$$\boxed{\varepsilon_{\text{L}}^N = -A_N - \Delta_{\text{xc}}^N}, \quad (2.83)$$

⁹From the linearity of the minimizing ensemble density matrix in Eq. (2.54), the exact ground-state density is a continuous piecewise linear function of \mathcal{N} : $n^{N-1+f}(\mathbf{r}) = (1-f)n^{N-1}(\mathbf{r}) + fn^N(\mathbf{r})$ and $n^{N+f}(\mathbf{r}) = (1-f)n^N(\mathbf{r}) + fn^{N+1}(\mathbf{r})$, just like the total ground-state energy.

due to the possible discontinuity Δ_{xc}^N in the KS potential.

Such a discontinuity can only come from the exchange-correlation part of the potential $v_{\text{xc}}^{\mathcal{N}}(\mathbf{r})$ since $v_{\text{ne}}(\mathbf{r})$ is independent from \mathcal{N} and the Hartree potential $v_{\text{H}}^{\mathcal{N}}(\mathbf{r}) = \int n^{\mathcal{N}}(\mathbf{r}')/|\mathbf{r} - \mathbf{r}'|d\mathbf{r}'$ is a continuous function of \mathcal{N} . So, we have

$$\Delta_{\text{xc}}^N = v_{\text{xc}}^{N+\delta}(\mathbf{r}) - v_{\text{xc}}^{N-\delta}(\mathbf{r}) = \left(\frac{\delta E_{\text{xc}}[n]}{\delta n(\mathbf{r})} \right)_{N+\delta} - \left(\frac{\delta E_{\text{xc}}[n]}{\delta n(\mathbf{r})} \right)_{N-\delta}, \quad (2.84)$$

i.e. Δ_{xc}^N is the *derivative discontinuity* in the exchange-correlation energy functional $E_{\text{xc}}[n]$. Theoretical and numerical examples show that this derivative discontinuity does exist [45–47].

2.5.5 Fundamental gap

The fundamental gap of the N -electron system is defined as

$$E_{\text{gap}}^N = I_N - A_N. \quad (2.85)$$

Using Eqs. (2.72) and (2.83), it can be expressed as

$$E_{\text{gap}}^N = \varepsilon_{\text{L}}^N - \varepsilon_{\text{H}}^N + \Delta_{\text{xc}}^N, \quad (2.86)$$

i.e., the difference between the LUMO and HOMO energies which defines the KS gap, $E_{\text{gap}}^{\text{KS},N} = \varepsilon_{\text{L}}^N - \varepsilon_{\text{H}}^N$, is not equal to the exact fundamental gap of the system. The difference comes from the derivative discontinuity Δ_{xc}^N . In practice, this last term can represent an important contribution to the fundamental gap. In the special case of open-shell systems, we have $\varepsilon_{\text{L}}^N = \varepsilon_{\text{H}}^N$, and thus if the fundamental gap of an open-shell system is not zero (Mott insulator), it is entirely given by Δ_{xc}^N .

3 Usual approximations for the exchange-correlation energy

We review here the main classes of usual approximations for the exchange-correlation energy. These classes are roughly ordered from the simplest to the most sophisticated ones. The discussion is focus on the contents of the approximations, not their performance in practical calculations. On average, more sophisticated approximations are usually more accurate than simpler ones, even though many exceptions can be found. For a review containing the explicit expressions of the main approximate functionals, see Ref. [17]. For extensive benchmarks of approximate functionals, see e.g. Refs. [16, 48]. The more complicated case of the so-called orbital-dependent exchange-correlation functionals such as exact exchange and the random-phase approximation, beyond usual hybrid or double-hybrid approximations, is treated separately in Section 4.

3.1 The local-density approximation

In the *local-density approximation* (LDA), introduced by Kohn and Sham [23], the exchange-correlation functional is approximated as

$$E_{xc}^{\text{LDA}}[n] = \int n(\mathbf{r}) \varepsilon_{xc}^{\text{UEG}}(n(\mathbf{r})) d\mathbf{r}, \quad (3.1)$$

where $\varepsilon_{xc}^{\text{UEG}}(n)$ is the exchange-correlation energy per particle of the infinite *uniform electron gas* (UEG) with the density n . The uniform electron gas represents a family of systems of interacting electrons with an arbitrary spatially constant density n that acts a parameter. Thus, in the LDA, the exchange-correlation energy per particle of an inhomogeneous system at a spatial point of density $n(\mathbf{r})$ is approximated as the exchange-correlation energy per particle of the uniform electron gas of the same density.

The function $\varepsilon_{xc}^{\text{UEG}}(n)$ is a sum of exchange and correlation contributions, $\varepsilon_{xc}^{\text{UEG}}(n) = \varepsilon_x^{\text{UEG}}(n) + \varepsilon_c^{\text{UEG}}(n)$. The exchange energy per particle of the uniform electron gas can be calculated analytically

$$\varepsilon_x^{\text{UEG}}(n) = C_x n^{1/3}, \quad (3.2)$$

where $C_x = -(3/4)(3/\pi)^{1/3}$. The LDA exchange functional is associated with the names of Dirac [49] and Slater [50].

Exercise 13 : Let us consider a local functional for the exchange energy of the form $E_x^{\text{local}}[n] = c \int n(\mathbf{r})^p d\mathbf{r}$ where c and p are constants. Show that requiring that $E_x^{\text{local}}[n]$ satisfies the scaling relation $E_x[n_\gamma] = \gamma E_x[n]$ [Eq. (2.41)] implies that $p = 4/3$, in accordance with the form of the LDA exchange functional.

Exercise 14 : The exchange energy of the uniform electron gas can be obtained by starting from the exact exchange energy expression for closed-shell systems [Eq. (3.18) or (4.1) with $\varphi_{i\uparrow}(\mathbf{r}) = \varphi_{i\downarrow}(\mathbf{r}) = \varphi_i(\mathbf{r})$]

$$E_x = - \sum_{i=1}^{N/2} \sum_{j=1}^{N/2} \iint \frac{\varphi_i^*(\mathbf{r}_1) \varphi_j(\mathbf{r}_1) \varphi_j^*(\mathbf{r}_2) \varphi_i(\mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2, \quad (3.3)$$

and by making the replacements $\varphi_i(\mathbf{r}) \rightarrow (1/V)^{1/2} e^{i\mathbf{k}\cdot\mathbf{r}}$ and $\sum_{i=1}^{N/2} \rightarrow V/(2\pi)^3 \int_{\Omega_{k_F}} d\mathbf{k}$ where Ω_{k_F} is the sphere of radius equal to the Fermi momentum $k_F = (3\pi^2 n)^{1/3}$ with the density

$n = N/V$, and doing the spatial integrations on a large box of volume V ($N \rightarrow \infty$ and $V \rightarrow \infty$ such that $n = N/V$ remains finite). Show that this gives

$$E_x^{\text{UEG}} = -\frac{V}{(2\pi)^6} \int_{\Omega_{k_F}} d\mathbf{k}_1 \int_{\Omega_{k_F}} d\mathbf{k}_2 \frac{4\pi}{|\mathbf{k}_1 - \mathbf{k}_2|^2}. \quad (3.4)$$

Calculate this integral to finally obtain the exchange energy per particle $\varepsilon_x^{\text{UEG}} = E_x^{\text{UEG}}/N$.

Exercise 15 : Show that the expression of the LDA exchange potential is (see Appendix B for an introduction to functional derivatives)

$$v_x^{\text{LDA}}(\mathbf{r}) = \frac{\delta E_x^{\text{LDA}}[n]}{\delta n(\mathbf{r})} = \frac{4}{3} C_x n(\mathbf{r})^{1/3}. \quad (3.5)$$

Does it satisfy the asymptotic behavior,

$$v_x(\mathbf{r}) \underset{r \rightarrow +\infty}{\sim} -\frac{1}{r}, \quad (3.6)$$

of the exact exchange potential?

The correlation energy per particle $\varepsilon_c^{\text{UEG}}(n)$ of the uniform electron gas cannot be calculated analytically. This quantity has been obtained numerically for a number of densities n using accurate quantum Monte Carlo calculations [51], and fitted to a parametrized function of n satisfying the known high- and low-density expansions. Expressed in terms of the Wigner-Seitz radius $r_s = (3/(4\pi n))^{1/3}$, the first terms of the high-density expansion ($r_s \rightarrow 0$) have the form

$$\varepsilon_c^{\text{UEG}} = A \ln r_s + B + C r_s \ln r_s + O(r_s), \quad (3.7)$$

and the first terms of the low-density expansion ($r_s \rightarrow +\infty$) have the form

$$\varepsilon_c^{\text{UEG}} = \frac{a}{r_s} + \frac{b}{r_s^{3/2}} + \frac{c}{r_s^2} + O\left(\frac{1}{r_s^{5/2}}\right), \quad (3.8)$$

where A , B , C , a , b , and c are constants. The two most used parametrizations are the ones of Vosko, Wilk, and Nusair (VWN) [52] and the one of Perdew and Wang (PW92) [53]. Their forms are too complicated to be given here. These parametrizations also include the generalization to spin densities, $\varepsilon_c^{\text{UEG}}(n_\uparrow, n_\downarrow)$ differing from $\varepsilon_c^{\text{UEG}}(n)$ for spin-polarized systems ($n_\uparrow \neq n_\downarrow$), which is sometimes specifically referred to as local-spin-density (LSD) approximation. For a recent review on the calculations of the exchange and correlation energies of the uniform electron gas, see Ref. [54].

Exercise 16 : The Wigner correlation functional is a simple functional derived from the low-density limit of the uniform electron gas

$$E_c^{\text{W}}[n] = \int n(\mathbf{r}) \frac{c}{d + r_s(\mathbf{r})} d\mathbf{r}, \quad (3.9)$$

where $r_s(\mathbf{r}) = (3/(4\pi n(\mathbf{r})))^{1/3}$, and c and d are two constants. Calculate the corresponding

correlation potential $v_c^W(\mathbf{r})$.

3.2 Generalized-gradient approximations

The next logical step beyond the LDA is the *gradient-expansion approximation* (GEA), initiated by Kohn and Sham [23]. One way of deriving the GEA is to start from the uniform electron gas, introduce a weak and slowly-varying external potential $v(\mathbf{r})$, and expand the exchange-correlation energy in terms of the gradients of the density. Alternatively, one can perform a semiclassical expansion of the exact $E_{xc}[n]$. At second order, one obtains a functional of the form

$$E_{xc}^{\text{GEA}}[n] = E_{xc}^{\text{LDA}}[n] + \int n(\mathbf{r})^{4/3} C_{xc}^{(2)}(n(\mathbf{r})) \left(\frac{|\nabla n(\mathbf{r})|}{n(\mathbf{r})^{4/3}} \right)^2 d\mathbf{r}, \quad (3.10)$$

where $C_{xc}^{(2)}(n) = C_x^{(2)} + C_c^{(2)}(n)$ is the sum of the exchange and correlation coefficients of the second-order gradient expansion. Note that the gradient expansion is most naturally written in terms of the reduced density gradient $|\nabla n|/n^{4/3}$ which is a dimensionless quantity. The GEA should improve over the LDA provided that the reduced density gradient is small. Unfortunately, for real systems, the reduced density gradient can be large in some regions of space, and the GEA turns out to be a worse approximation than the LDA.

The failure of the GEA led to the development of *generalized-gradient approximations* (GGAs), which really started in the 1980s, of the generic form

$$E_{xc}^{\text{GGA}}[n] = \int e_{xc}^{\text{GGA}}(n(\mathbf{r}), \nabla n(\mathbf{r})) d\mathbf{r}, \quad (3.11)$$

where e_{xc}^{GGA} is some function. The GGAs are *semilocal* approximations in the sense that e_{xc}^{GGA} does not only use the local value of the density $n(\mathbf{r})$ but also its gradient $\nabla n(\mathbf{r})$ ¹⁰. For simplicity, we consider here only the spin-independent form, but in practice GGA functionals are more generally formulated in terms of spin densities (n_\uparrow, n_\downarrow) and their gradients ($\nabla n_\uparrow, \nabla n_\downarrow$)¹¹.

Exercise 17 : Show that the potential of a GGA functional has the following form

$$v_{xc}^{\text{GGA}}(\mathbf{r}) = \frac{\delta E_{xc}^{\text{GGA}}[n]}{\delta n(\mathbf{r})} = \frac{\partial e_{xc}^{\text{GGA}}}{\partial n}(n(\mathbf{r}), \nabla n(\mathbf{r})) - \nabla \cdot \frac{\partial e_{xc}^{\text{GGA}}}{\partial \nabla n}(n(\mathbf{r}), \nabla n(\mathbf{r})), \quad (3.12)$$

which is a 3-dimensional generalization of Eq. (B.13). Give then a practical expression for the contribution $V_{xc,\mu\nu}^{\text{GGA}}$ of this potential to the KS Fock matrix [Eq. (1.51)].

Many GGA functionals have been proposed. We very briefly review here some of the most widely used ones.

B88 exchange functional

¹⁰For generality and simplicity, we consider here that the GGAs depend on the density gradient $\nabla n(\mathbf{r})$, but in practice GGAs depend only on the module of the density gradient $|\nabla n(\mathbf{r})|$, or equivalently on its square $(\nabla n(\mathbf{r}))^2$, and not on its direction.

¹¹Again, in practice, the spin-dependent GGAs do not actually depend on the gradients ∇n_\uparrow and ∇n_\downarrow but on the scalar quantities $(\nabla n_\uparrow)^2$, $(\nabla n_\downarrow)^2$, and $\nabla n_\uparrow \cdot \nabla n_\downarrow$.

The Becke 88 (B88 or B) [55] exchange functional is formulated as an additive correction to LDA. It consists in a compact function of n and $|\nabla n|/n^{4/3}$ chosen so as to satisfy the exact asymptotic behavior of the exchange energy per particle for finite systems (see Exercise 7), and with an empirical parameter fitted to Hartree-Fock exchange energies of rare-gas atoms.

LYP correlation functional

The Lee-Yang-Parr (LYP) [56] correlation functional is one of the rare functionals which have not been constructed starting from LDA. It originates from the Colle-Salvetti [57] correlation-energy approximation depending on the Hartree-Fock pair density and containing four parameters fitted to Helium data. By making a further reasonable approximation, LYP turned the Colle-Salvetti expression into a density functional depending on the density n , the square of the density gradient $(\nabla n)^2$, and the Laplacian of the density $\nabla^2 n$. The dependence on the Laplacian can be exactly eliminated by an integration by parts [58].

PW91 exchange-correlation functional

The Perdew-Wang 91 (PW91) (see Refs. [41, 59, 60]) exchange-correlation functional is based on a model of the exchange hole $n_x(\mathbf{r}_1, \mathbf{r}_2)$ in Eq. (2.15) and of the coupling-constant-integrated correlation hole in Eq. (2.30). The idea is to start from the GEA model of these holes given as gradient expansions and remove the unrealistic long-range parts of these holes to restore important conditions satisfied by the LDA. Specifically, the spurious positive parts of the GEA exchange hole are removed to enforce the negativity condition of Eq. (2.12) and a cutoff in $|\mathbf{r}_1 - \mathbf{r}_2|$ is applied to enforce the normalization condition of Eq. (2.10). Similarly, a cutoff is applied on the GEA correlation hole to enforce the condition that the hole integrates to zero [Eq. (2.17)]. The exchange and correlation energies per particle calculated from these numerical holes are then fitted to functions of n and $|\nabla n|$ chosen to satisfy a number of exact conditions.

PBE exchange-correlation functional

The Perdew-Burke-Ernzerhof (PBE) [61] exchange-correlation functional is a simplification of the PW91 functional. The exchange and correlation energies per particle are expressed as simpler functions of n and $|\nabla n|$ enforcing less exact conditions and with no fitted parameters. Specifically, the function used for correlation enforces the second-order small-gradient expansion in the high-density limit, the vanishing of correlation in the large-gradient limit, and removes the logarithm divergence of the LDA in the high-density limit [see Eq. (3.7)]. The function used for exchange is chosen to cancel out the second-order small-gradient expansion of correlation and enforces the Lieb-Oxford bound in the large-gradient limit.

3.3 Meta-generalized-gradient approximations

The *meta-generalized-gradient approximations* (meta-GGAs or mGGAs) are of the generic form

$$E_{xc}^{\text{mGGA}}[n, \tau] = \int e_{xc}^{\text{mGGA}}(n(\mathbf{r}), \nabla n(\mathbf{r}), \nabla^2 n(\mathbf{r}), \tau(\mathbf{r})) d\mathbf{r}, \quad (3.13)$$

i.e., they use more ingredients than the GGAs, namely the Laplacian of the density $\nabla^2 n(\mathbf{r})$ and/or the non-interacting positive kinetic energy density $\tau(\mathbf{r})$

$$\tau(\mathbf{r}) = \frac{1}{2} \sum_{i=1}^N |\nabla \varphi_i(\mathbf{r})|^2, \quad (3.14)$$

where $\varphi_i(\mathbf{r})$ are the orbitals composing a single-determinant wave function Φ .

A meta-GGA functional can be either considered as an implicit functional of the density in the KS method, i.e. $E_{xc}[n] = E_{xc}^{mGGA}[n, \tau_{\Phi}[n]]$, or more commonly as an explicit functional of a single-determinant Φ in the GKS method, i.e. $E_{xc}^S[\Phi] = E_{xc}^{mGGA}[n_{\Phi}, \tau_{\Phi}]$. In the GKS method, a meta-GGA functional generates a non-multiplicative potential (see, e.g., Ref. [17]). The meta-GGAs are considered part of the family of semilocal approximations, in the sense that $\tau(\mathbf{r})$ depends only the gradient of the orbitals at point \mathbf{r} . Again, we consider here only the spin-independent form, but meta-GGAs are more generally formulated in terms of spin-resolved quantities ($n_{\uparrow}, n_{\downarrow}, \nabla n_{\uparrow}, \nabla n_{\downarrow}, \nabla^2 n_{\uparrow}, \nabla^2 n_{\downarrow}, \tau_{\uparrow}, \tau_{\downarrow}$).

One motivation for the introduction of the variable $\tau(\mathbf{r})$ is that it appears in the expansion of the spherically average exchange hole $\tilde{n}_x(\mathbf{r}_1, r_{12})$ for small interelectronic distances r_{12} , which for the case of a closed-shell system is [62]

$$\tilde{n}_x(\mathbf{r}_1, r_{12}) = -\frac{n(\mathbf{r}_1)}{2} - \frac{1}{3} \left(\frac{1}{4} \nabla^2 n(\mathbf{r}_1) - 4\tau(\mathbf{r}_1) + \frac{|\nabla n(\mathbf{r}_1)|^2}{8n(\mathbf{r}_1)} \right) r_{12}^2 + O(r_{12}^4). \quad (3.15)$$

Thus $\tau(\mathbf{r})$ is needed to describe the curvature of the exchange hole. Another important motivation is that $\tau(\mathbf{r})$ can be used as an indicator of one-orbital spatial regions (regions containing one or two electrons in a single orbital). This is done by comparing $\tau(\mathbf{r})$ with the von Weizsäcker kinetic energy density

$$\tau^W(\mathbf{r}) = \frac{|\nabla n(\mathbf{r})|^2}{8n(\mathbf{r})}, \quad (3.16)$$

which is the exact $\tau(\mathbf{r})$ for one electron and for two electrons in a single orbital. In practice, $\tau(\mathbf{r})$ is often used through the variables $z(\mathbf{r}) = \tau^W(\mathbf{r})/\tau(\mathbf{r})$ or $\alpha(\mathbf{r}) = (\tau(\mathbf{r}) - \tau^W(\mathbf{r}))/\tau^{\text{UEG}}(\mathbf{r})$ where $\tau^{\text{UEG}}(\mathbf{r}) = c n(\mathbf{r})^{5/3}$.

Exercise 18 : Show that the von Weizsäcker kinetic energy density in Eq. (3.16) is indeed the exact (non-interacting) positive kinetic energy density $\tau(\mathbf{r})$ for one electron and for two electrons in a single orbital.

One the most used meta-GGA approximation is the Tao-Perdew-Staroverov-Scuseria (TPSS) [63] exchange-correlation functional. A more recently developed meta-GGA approximation is the SCAN exchange-correlation functional [64] which satisfies 17 known exact constraints and contains 7 parameters determined by fitting to a few simple systems.

3.4 Single-determinant hybrid approximations

3.4.1 Hybrid approximations

Based on arguments relying on the adiabatic-connection formalism, in 1993 Becke [65] proposed to mix a fraction of the exact or Hartree-Fock (HF) exchange energy E_x^{HF} with GGA functionals. In particular, he proposed a three-parameter hybrid (3H) approximation [66] of the form

$$E_{xc}^{3H}[\Phi] = a E_x^{\text{HF}}[\Phi] + b E_x^{\text{GGA}}[n_{\Phi}] + (1-a-b) E_x^{\text{LDA}}[n_{\Phi}] + c E_c^{\text{GGA}}[n_{\Phi}] + (1-c) E_c^{\text{LDA}}[n_{\Phi}], \quad (3.17)$$

where the three parameters a , b , and c are determined by fitting to experimental data. The functional $E_{xc}^{3H}[\Phi]$ is thought of as a functional of a single-determinant wave function $\Phi \in \mathcal{S}$

since $E_x^{\text{HF}}[\Phi]$ is itself a functional of Φ

$$\begin{aligned} E_x^{\text{HF}}[\Phi] &= \langle \Phi | \hat{W}_{\text{ee}} | \Phi \rangle - E_{\text{H}}[n_{\Phi}] \\ &= -\frac{1}{2} \sum_{\sigma \in \{\uparrow, \downarrow\}} \sum_{i=1}^{N_{\sigma}} \sum_{j=1}^{N_{\sigma}} \iint \frac{\varphi_{i\sigma}^*(\mathbf{r}_1) \varphi_{j\sigma}(\mathbf{r}_1) \varphi_{j\sigma}^*(\mathbf{r}_2) \varphi_{i\sigma}(\mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2, \end{aligned} \quad (3.18)$$

where $\{\varphi_{i\sigma}\}_{i=1, \dots, N_{\sigma}}$ are the orbitals occupied in Φ . The HF exchange energy has exactly the same form as the exact-exchange energy given in Eq. (4.1). The notation ‘‘HF’’ is there to specify that it is not viewed as a functional of the density but instead as a functional of Φ . These hybrids are thus approximations within the GKS method, with a term $S[\Phi] = aE_x^{\text{HF}}[\Phi]$ which generates a nonlocal HF exchange potential $av_{x,\sigma}^{\text{HF}}(\mathbf{r}, \mathbf{r}')$ instead of a local one (see Exercise 19). Having a nonlocal potential is perfectly allowed in the GKS method.¹² The most famous and widely used three-parameter hybrid approximation is B3LYP [67], which uses the B88 exchange functional and the LYP correlation functional, and the parameters $a = 0.20$, $b = 0.72$, and $c = 0.81$.

In 1996, Becke proposed a simpler one-parameter hybrid (1H) approximation [68]

$$E_{\text{xc}}^{1\text{H}}[\Phi] = a E_x^{\text{HF}}[\Phi] + (1 - a) E_x^{\text{DFA}}[n_{\Phi}] + E_c^{\text{DFA}}[n_{\Phi}], \quad (3.19)$$

where $E_x^{\text{DFA}}[n]$ and $E_c^{\text{DFA}}[n]$ can be any (semilocal) density-functional approximations (DFA), and the fraction a of HF exchange has to be determined.

The main benefit of adding a fraction a of HF exchange is to decrease the *self-interaction error* in the exchange functional which tends to favor too much delocalized electron densities over localized electron densities (leading to problems with dissociation of charged fragments, reaction barriers, radicals,...). However, a too large a tends to increase the *static-correlation error* (leading to problems with stretched chemical bonds, transition metal elements, ...). The fraction of HF exchange should thus be small enough to keep the compensation of errors usually occurring between the approximate semilocal exchange functional $E_x^{\text{DFA}}[n]$ and the approximate semilocal correlation functional $E_c^{\text{DFA}}[n]$. Fits to experimental data often give an optimal parameter a around 0.25. A rationale has also been proposed in favor of this value [69]. For example, PBE0 [70, 71] is a popular one-parameter hybrid approximation which uses $a = 0.25$ and the PBE exchange and correlation functionals.

A strategy that has been sometimes used to construct approximations of the form of Eq. (3.19) is to employ parameterized flexible functions for $E_x^{\text{DFA}}[n]$ and $E_c^{\text{DFA}}[n]$, and systematically optimize all the parameters (including the fraction a of HF exchange) on large sets of physicochemical properties of molecular systems. For example, the Becke 97 (B97) exchange-correlation functional [72] is a hybrid GGA approximation containing 13 parameters optimized on atomic exchange and correlation energies, atomization energies, ionization potentials, and proton affinities. Another example is the so-called family of ‘‘Minnesota’’ functionals, and in particular the M06 exchange-correlation functional [73] which is a hybrid meta-GGA approximation containing 36 parameters optimized on a very large set of diverse physicochemical properties concerning main-group thermochemistry, reaction barrier heights, noncovalent interactions, electronic spectroscopy, and transition metal bonding.

¹²In fact, the possibility of combining a nonlocal HF potential with a local correlation potential was mentioned already in 1965 in the paper by Kohn and Sham [23].

Exercise 19 : Show that the functional derivative of E_x^{HF} with respect to $\varphi_{i\sigma}^*(\mathbf{r})$ gives

$$\frac{\delta E_x^{\text{HF}}}{\delta \varphi_{i\sigma}^*(\mathbf{r})} = \int v_{x,\sigma}^{\text{HF}}(\mathbf{r}, \mathbf{r}') \varphi_{i\sigma}(\mathbf{r}') d\mathbf{r}', \quad (3.20)$$

where $v_{x,\sigma}^{\text{HF}}(\mathbf{r}, \mathbf{r}')$ is the nonlocal HF exchange potential

$$v_{x,\sigma}^{\text{HF}}(\mathbf{r}, \mathbf{r}') = - \sum_{j=1}^{N_\sigma} \frac{\varphi_{j\sigma}(\mathbf{r}) \varphi_{j\sigma}^*(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|}. \quad (3.21)$$

Show also that Eq. (3.20) can be reformulated in terms of a local HF exchange potential $v_{x,i\sigma}^{\text{HF}}(\mathbf{r})$ depending on each orbital it acts on.

3.4.2 Range-separated hybrid approximations

Based on earlier ideas of Savin [74], in 2001, Iikura, Tsuneda, Yanai, and Hirao [75] proposed a long-range correction (LC) scheme

$$E_{xc}^{\text{LC}}[\Phi] = E_x^{\text{lr,HF}}[\Phi] + E_x^{\text{sr,DFA}}[n_\Phi] + E_c^{\text{DFA}}[n_\Phi], \quad (3.22)$$

where $E_x^{\text{lr,HF}}[\Phi]$ is the HF exchange energy for a long-range electron-electron interaction $w_{ee}^{\text{lr}}(r_{12}) = \text{erf}(\mu r_{12})/r_{12}$ (where $r_{12} = |\mathbf{r}_1 - \mathbf{r}_2|$ is the interelectronic distance and erf is the error function, see Exercise 20)

$$E_x^{\text{lr,HF}}[\Phi] = -\frac{1}{2} \sum_{\sigma \in \{\uparrow, \downarrow\}} \sum_{i=1}^{N_\sigma} \sum_{j=1}^{N_\sigma} \iint \varphi_{i\sigma}^*(\mathbf{r}_1) \varphi_{j\sigma}(\mathbf{r}_1) \varphi_{j\sigma}^*(\mathbf{r}_2) \varphi_{i\sigma}(\mathbf{r}_2) w_{ee}^{\text{lr}}(r_{12}) d\mathbf{r}_1 d\mathbf{r}_2, \quad (3.23)$$

and $E_x^{\text{sr,DFA}}[n]$ is the DFA exchange energy for the complementary short-range interaction $w_{ee}^{\text{sr}}(r_{12}) = 1/r_{12} - w_{ee}^{\text{lr}}(r_{12})$. Similarly to the hybrid approximations of Section 3.4.1, the introduction of a fraction of long-range HF exchange reduces the self-interaction error (see, e.g., Ref. [76]). In addition, the short-range exchange part is easier to approximate with semilocal DFA. In particular, the $-1/r$ asymptotic behavior of the exchange potential [Eq. (3.6)], which is difficult to satisfy with DFAs, does not apply anymore to the short-range exchange potential. The parameter μ (also sometimes denoted as ω) in the error function controls the range of the separation and must be chosen, e.g. by fitting to experimental data. In practice, a value around $\mu \approx 0.3-0.5 \text{ bohr}^{-1}$ is often found to be optimal. A popular example of such LC approximations is LC- ω PBE [77] which uses a short-range version of the PBE exchange functional, as well as the standard PBE correlation functional. Note that the LC scheme has also been referred to as the range-separated hybrid exchange (RSHX) scheme [78].

Exercise 20 : Using the definition of the error function

$$\text{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt, \quad (3.24)$$

show that $\text{erf}(x) \underset{x \rightarrow 0}{\sim} (2/\sqrt{\pi})x$ and $\text{erf}(x) \xrightarrow{x \rightarrow +\infty} 1$. Draw then the form of the long-range

interaction $\text{erf}(\mu r)/r$ as a function r for different values of the range-separation parameter: $\mu = 0$, $\mu = 1$, and $\mu \rightarrow +\infty$.

In 2004, Yanai, Tew, and Handy [79], introduced a more flexible scheme called the Coulomb-attenuating method (CAM) [79] in which fractions of HF exchange are added at both short and long range

$$E_{xc}^{\text{CAM}}[\Phi] = a E_x^{\text{sr,HF}}[\Phi] + b E_x^{\text{lr,HF}}[\Phi] + (1-a) E_x^{\text{sr,DFA}}[n_\Phi] + (1-b) E_x^{\text{lr,DFA}}[n_\Phi] + E_c^{\text{DFA}}[n_\Phi], \quad (3.25)$$

where $E_x^{\text{sr,HF}}[\Phi]$ is the HF exchange energy for the short-range interaction $w_{ee}^{\text{sr}}(r_{12})$ and $E_x^{\text{lr,DFA}}[n]$ is the DFA exchange energy for the long-range interaction $w_{ee}^{\text{lr}}(r_{12})$. The reintroduction of HF exchange at short range improves thermodynamic properties such as atomization energies. According to this scheme, the authors proposed the CAM-B3LYP approximation which uses short- and long-range versions of the B88 exchange functional, the same correlation functional used in B3LYP (i.e., $0.81 E_c^{\text{LYP}}[n] + 0.19 E_c^{\text{LDA}}[n]$), and optimized parameters $a = 0.19$, $b = 0.65$, and $\mu = 0.33 \text{ bohr}^{-1}$. Another example in this class of approximations is the ω B97X exchange-correlation functional [80] which is based on the B97 exchange-correlation functional with reoptimized parameters, and uses $a = 0.16$, $b = 1$, and $\mu = 0.3 \text{ bohr}^{-1}$. Another functional that can be considered as part of this class of approximations is the Heyd-Scuseria-Ernzerhof (HSE) exchange-correlation functional [81], which uses the parameters $a = 0.25$, $b = 0$ (i.e. no long-range HF exchange), and $\mu = 0.15 \text{ bohr}^{-1}$, with a long-range version of the PBE exchange correlation functional and the standard PBE correlation functional. The absence of HF exchange at very long range makes this approximation particularly useful for solids.

3.5 Multideterminant hybrid approximations

3.5.1 Double-hybrid approximations

In 2006, Grimme [82] introduced a two-parameter double-hybrid (2DH) approximation

$$E_{xc}^{2\text{DH}} = a_x E_x^{\text{HF}}[\Phi] + (1 - a_x) E_x^{\text{DFA}}[n_\Phi] + (1 - a_c) E_c^{\text{DFA}}[n_\Phi] + a_c E_c^{\text{MP2}}, \quad (3.26)$$

mixing a fraction a_x of the HF exchange energy with a semilocal exchange DFA, and a fraction a_c of the second-order Møller-Plesset (MP2) correlation energy E_c^{MP2} with a semilocal correlation DFA. In Eq. (3.26), the first three terms are first calculated in self-consistent manner, and then the last term E_c^{MP2} is added perturbatively using the orbitals determined in the first step. The expression of E_c^{MP2} is

$$E_c^{\text{MP2}} = -\frac{1}{4} \sum_{i=1}^N \sum_{j=1}^N \sum_{a \geq N+1} \sum_{b \geq N+1} \frac{|\langle \psi_i \psi_j | \psi_a \psi_b \rangle|^2}{\varepsilon_a + \varepsilon_b - \varepsilon_i - \varepsilon_j}, \quad (3.27)$$

where i, j and a, b run over occupied and virtual spin orbitals, respectively, ε_k are spin orbital energies, and $\langle \psi_i \psi_j | \psi_a \psi_b \rangle = \langle \psi_i \psi_j | \psi_a \psi_b \rangle - \langle \psi_i \psi_j | \psi_b \psi_a \rangle$ are antisymmetrized two-electron integrals with (in physicists' notation)

$$\langle \psi_p \psi_q | \psi_r \psi_s \rangle = \iint \frac{\psi_p^*(\mathbf{x}_1) \psi_q^*(\mathbf{x}_2) \psi_r(\mathbf{x}_1) \psi_s(\mathbf{x}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{x}_1 d\mathbf{x}_2. \quad (3.28)$$

Note that the notation in Eq. (3.27) assumes that the one-electron wave-function space is spanned by a discrete set of spin orbitals. In the exact theory, the continuum limit of the set of virtual spin orbitals is implied.

The presence of nonlocal MP2 correlation allows one to use a larger fraction of nonlocal HF exchange. For example, Grimme proposed the B2-PLYP approximation which uses the B88 exchange and LYP correlation functionals, and the parameters $a_x = 0.53$ and $a_c = 0.27$ optimized on experimental heats of formation of a set of molecules.

In 2011, Sharkas, Toulouse, and Savin [83] showed that double hybrids can be understood as approximations of a *multideterminant extension of the KS method* based on the adiabatic-connection formalism in which the exact ground-state energy is written as

$$E_0 = \min_{\Psi \in \mathcal{W}} \left\{ \langle \Psi | \hat{T} + \hat{V}_{\text{ne}} + \lambda \hat{W}_{\text{ee}} | \Psi \rangle + \bar{E}_{\text{Hxc}}^\lambda[n_\Psi] \right\}, \quad (3.29)$$

where $\bar{E}_{\text{Hxc}}^\lambda[n] = (1 - \lambda)E_{\text{H}}[n] + (1 - \lambda)E_{\text{x}}[n] + \bar{E}_{\text{c}}^\lambda[n]$ and $\bar{E}_{\text{c}}^\lambda[n] = E_{\text{c}}[n] - \lambda^2 E_{\text{c}}[n_{1/\lambda}]$. At second order of a non-linear Møller-Plesset-like perturbation theory, and using $E_{\text{c}}[n_{1/\lambda}] \approx E_{\text{c}}[n]$, we obtain a one-parameter double-hybrid (1DH) approximation

$$E_{\text{xc}}^{\text{1DH}} = \lambda E_{\text{x}}^{\text{HF}}[\Phi] + (1 - \lambda) E_{\text{x}}^{\text{DFA}}[n_\Phi] + (1 - \lambda^2) E_{\text{c}}^{\text{DFA}}[n_\Phi] + \lambda^2 E_{\text{c}}^{\text{MP2}}, \quad (3.30)$$

where the fraction HF exchange $a_x = \lambda$ is now connected to the fraction of MP2 correlation $a_c = \lambda^2$. It turns out that using $a_x = \lambda = 0.53$ nearly reproduces the parameter $a_c = \lambda^2 = 0.28 \approx 0.27$ independently optimized in Eq. (3.26). Likewise, Fromager [84] proposed a rigorous formulation of the two-parameter double-hybrid approximations. The multideterminant extension of the KS method can also be used to rigorously combine wave-function methods such as MCSCF with DFT [85].

The double-hybrid approximations are examples of correlation functionals depending on virtual orbitals. Another example of a correlation functional depending on virtual orbitals is the random-phase approximation (RPA), which goes beyond second order and has been the subject of intensive developments since the 2000s.

3.5.2 Range-separated double-hybrid approximations

In 1996, Savin [74] introduced the *range-separated multideterminant extension of the KS scheme* in which the exact ground-state energy is written as

$$E_0 = \min_{\Psi \in \mathcal{W}} \left\{ \langle \Psi | \hat{T} + \hat{V}_{\text{ne}} + \hat{W}_{\text{ee}}^{\text{lr}} | \Psi \rangle + \bar{E}_{\text{Hxc}}^{\text{sr}}[n_\Psi] \right\}, \quad (3.31)$$

where $\hat{W}_{\text{ee}}^{\text{lr}}$ is the long-range electron-electron operator for the pair potential $w_{\text{ee}}^{\text{lr}}(r_{12})$ and $\bar{E}_{\text{Hxc}}^{\text{sr}}[n]$ is the complementary short-range density functional. This approach have been used to rigorously combine many wave-function methods with DFT (see references in Ref. [17]).

In 2005, starting from Eq. (3.31) and using a a non-linear long-range second-order Møller-Plesset-like perturbation theory, Ángyán, Gerber, Savin, and Toulouse [86] introduced a *range-separated double-hybrid (RSDH)* approximation (also called RSH+MP2)

$$E_{\text{xc}}^{\text{RSDH}} = E_{\text{x}}^{\text{lr, HF}}[\Phi] + E_{\text{x}}^{\text{sr, DFA}}[n_\Phi] + E_{\text{c}}^{\text{sr, DFA}}[n_\Phi] + E_{\text{c}}^{\text{lr, MP2}}, \quad (3.32)$$

where $E_{\text{c}}^{\text{lr, MP2}}$ is the MP2 correlation energy calculated with long-range two-electron integrals and the previously calculated orbitals. The RSDH scheme is thus similar to the double-hybrid approximations of Section 3.5.1 but with range separation. One of main advantages of using a long-range MP2 correlation energy is the correct qualitative description of London dispersion interaction energies, while displaying a fast convergence with the basis size [87]. For more accurate results, one can go beyond second order by using long-range coupled-cluster [88] or random-phase approximations [89–91]. Extensions of this scheme to a more flexible CAM decomposition have also been proposed [92–95].

3.6 Dispersion corrections

3.6.1 Semiempirical dispersion corrections

To explicitly account for London dispersion interactions, it has been proposed in the 2000s to add to the standard approximate functionals a *semiempirical dispersion correction* of the form [96–98]

$$E_{\text{disp}} = -s \sum_{\alpha < \beta} f(R_{\alpha\beta}) \frac{C_6^{\alpha\beta}}{R_{\alpha\beta}^6}, \quad (3.33)$$

where $R_{\alpha\beta}$ is the distance between each pair of atoms, $C_6^{\alpha\beta}$ is the dispersion coefficient between these atoms, $f(R_{\alpha\beta})$ is a parametrized damping function which tends to 1 at large $R_{\alpha\beta}$ and tends to zero at small $R_{\alpha\beta}$, and s is a possible scaling parameter that can be adjusted for each approximate functional. The dispersion coefficient $C_6^{\alpha\beta}$ for any pair of atoms is empirically calculated from tabulated same-atom dispersion coefficients $C_6^{\alpha\alpha}$ and/or atomic polarizabilities. The most recent versions include $C_8^{\alpha\beta}$ two-body terms and $C_9^{\alpha\beta\gamma}$ three-body terms [99]. This approach was named “DFT-D” by Grimme [98]. Examples of DFT-D functionals are: PBE-D [100], B97-D [100], B3LYP-D [100], ω B97X-D [101], B2PLYP-D [102]. There have also been various proposals to make the determination of dispersion coefficients less empirical, such as the scheme of Becke and Johnson [103] based on the exchange-hole dipole moment, the scheme of Tkatchenko and Scheffler [104] based on a Hirshfeld atomic partitioning, or the scheme of Sato and Nakai [105] based on the local-response approximation [106].

3.6.2 Nonlocal van der Waals density functionals

Another approach to describe dispersion interactions is to add to the standard approximate functionals a *nonlocal van der Waals density functional* of the form

$$E_c^{\text{nl}}[n] = \frac{1}{2} \iint n(\mathbf{r}_1)n(\mathbf{r}_2)\phi(\mathbf{r}_1, \mathbf{r}_2)d\mathbf{r}_1d\mathbf{r}_2, \quad (3.34)$$

where $\phi(\mathbf{r}_1, \mathbf{r}_2)$ is a correlation kernel. Two main families of such nonlocal correlation functionals exist: the “van der Waals density functionals” (vdW-DF) of Langreth, Lundqvist and coworkers [107, 108] and the Vydrov-Van Voorhis (VV) functionals [109–111].

For example, the VV10 nonlocal correlation functional [111] (2010) uses a theory-based kernel $\phi(\mathbf{r}_1, \mathbf{r}_2)$ with two adjustable parameters. Nonlocal van der Waals density functionals are less empirical but more computationally expensive than semiempirical dispersion corrections. Examples of functionals using VV10 are: ω B97X-V [112] and ω B97M-V [113].

4 Some less usual orbital-dependent exchange-correlation functionals

We discuss here some exchange-correlation energy functionals explicitly depending on the KS orbitals. Since the KS orbitals are themselves functionals of the density, these exchange-correlation expressions are thus *implicit* functionals of the density (for notational simplicity, this dependence on the density will not be explicitly indicated). In fact, the (range-separated) hybrid and double-hybrid approximations of Sections 3.4 and 3.5 are sometimes considered to belong to this family, but they are more commonly considered within the GKS method, i.e. the orbitals are obtained with a *nonlocal* potential. In this Section, we are concerned with orbital-dependent exchange-correlation energy functionals within the KS method, i.e. with orbitals obtained with a local potential. These approximations tend to be more computationally involved than the approximations of the previous Section and are thus much less used.

4.1 Exact exchange

Being defined as an expectation value over a single-determinant wave function [see Eq. (1.28)], the exact exchange (EXX) energy functional can be expressed in terms of the orbitals

$$E_x = -\frac{1}{2} \sum_{\sigma \in \{\uparrow, \downarrow\}} \sum_{i=1}^{N_\sigma} \sum_{j=1}^{N_\sigma} \iint \frac{\varphi_{i\sigma}^*(\mathbf{r}_1) \varphi_{j\sigma}(\mathbf{r}_1) \varphi_{j\sigma}^*(\mathbf{r}_2) \varphi_{i\sigma}(\mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2, \quad (4.1)$$

where i and j run over spatial occupied orbitals. The exchange energy in Eq. (4.1) has exactly the same form as the HF exchange [Eq. (3.18)], but the orbitals used in this expression are in general different.

Since the exact exchange energy in Eq. (4.1) is not an explicit functional of the density, the corresponding exchange potential $v_x(\mathbf{r}) = \delta E_x / \delta n(\mathbf{r})$ cannot be calculated directly. We can however find an workable equation for $v_x(\mathbf{r})$ by first considering the functional derivative of E_x with respect to the KS potential $v_s(\mathbf{r})$ and then applying the chain rule

$$\frac{\delta E_x}{\delta v_s(\mathbf{r})} = \int \frac{\delta E_x}{\delta n(\mathbf{r}')} \frac{\delta n(\mathbf{r}')}{\delta v_s(\mathbf{r})} d\mathbf{r}'. \quad (4.2)$$

Introducing the non-interacting KS static linear-response function $\chi_0(\mathbf{r}', \mathbf{r}) = \delta n(\mathbf{r}') / \delta v_s(\mathbf{r})$, we can rewrite Eq. (4.2) as

$$\int v_x(\mathbf{r}') \chi_0(\mathbf{r}', \mathbf{r}) d\mathbf{r}' = \frac{\delta E_x}{\delta v_s(\mathbf{r})}, \quad (4.3)$$

which is known as the *optimized-effective-potential* (OEP) equation for the exact-exchange potential. Explicit expressions in terms of the orbitals can be derived for $\delta E_x / \delta v_s(\mathbf{r})$ and $\chi_0(\mathbf{r}', \mathbf{r})$.

Exercise 21 : Using first-order perturbation theory on the KS system, show that

$$\frac{\delta \varphi_{i\sigma}(\mathbf{r}')}{\delta v_s(\mathbf{r})} = - \sum_{\substack{p \geq 1 \\ p \neq i}} \frac{\varphi_{p\sigma}^*(\mathbf{r}) \varphi_{i\sigma}(\mathbf{r})}{\varepsilon_{p\sigma} - \varepsilon_{i\sigma}} \varphi_{p\sigma}(\mathbf{r}'). \quad (4.4)$$

where the sum is over all spatial orbitals p different from orbital i but of the same spin.

Show then that the expression of $\chi_0(\mathbf{r}', \mathbf{r})$ is

$$\chi_0(\mathbf{r}', \mathbf{r}) = - \sum_{\sigma \in \{\uparrow, \downarrow\}} \sum_{i=1}^{N_\sigma} \sum_{a \geq N_\sigma + 1} \frac{\varphi_{i\sigma}^*(\mathbf{r}') \varphi_{a\sigma}^*(\mathbf{r}) \varphi_{i\sigma}(\mathbf{r}) \varphi_{a\sigma}(\mathbf{r}')}{\varepsilon_{a\sigma} - \varepsilon_{i\sigma}} + \text{c.c.} , \quad (4.5)$$

where a runs over virtual spatial orbitals and c.c. stands for the complex conjugate. Finally, by using the definition of E_x in terms of the orbitals, show that

$$\frac{\delta E_x}{\delta v_s(\mathbf{r})} = \sum_{\sigma \in \{\uparrow, \downarrow\}} \sum_{i=1}^{N_\sigma} \sum_{j=1}^{N_\sigma} \sum_{a \geq N_\sigma + 1} (\varphi_{a\sigma} \varphi_{j\sigma} | \varphi_{j\sigma} \varphi_{i\sigma}) \frac{\varphi_{a\sigma}(\mathbf{r}) \varphi_{i\sigma}^*(\mathbf{r})}{\varepsilon_{a\sigma} - \varepsilon_{i\sigma}} + \text{c.c.} , \quad (4.6)$$

where $(\varphi_{a\sigma} \varphi_{j\sigma} | \varphi_{j\sigma} \varphi_{i\sigma})$ are the two-electron integrals [see Eq. (1.50) for the definition] in the basis of the KS spatial orbitals.

Applying this OEP method with the EXX energy (and no correlation energy functional) is an old idea [114, 115], but reasonable efficient calculations for molecules have been possible only relatively recently [116, 117]. The EXX occupied orbitals turn out to be very similar to the HF occupied orbitals, and thus the EXX ground-state properties are also similar to the HF ones. However, the EXX virtual orbitals tend to be much less diffuse than the HF virtual orbitals, and may be more adapted for calculating excited-state properties.

4.2 Second-order Görling-Levy perturbation theory

In 1993, Görling and Levy [33, 34] developed a perturbation theory in terms of the coupling constant λ of the adiabatic connection (Section 2.1.2) which provides an explicit orbital-dependent second-order approximation for the correlation energy functional. The Hamiltonian along the adiabatic connection [Eq. (2.19)] can be written as

$$\begin{aligned} \hat{H}^\lambda &= \hat{T} + \lambda \hat{W}_{ee} + \hat{V}^\lambda \\ &= \hat{H}_s + \lambda(\hat{W}_{ee} - \hat{V}_{Hx}) - \hat{V}_c^\lambda, \end{aligned} \quad (4.7)$$

where $\hat{H}_s = \hat{H}^{\lambda=0} = \hat{T} + \hat{V}_s$ is the KS non-interacting reference Hamiltonian (which will be assumed to have a nondegenerate ground state). Equation (4.7) was obtained by decomposing the potential operator keeping the density constant as $\hat{V}^\lambda = \hat{V}_s - \lambda \hat{V}_{Hx} - \hat{V}_c^\lambda$ where $\hat{V}_s = \hat{V}^{\lambda=0}$ is the KS potential operator, $\lambda \hat{V}_{Hx}$ is the Hartree-exchange potential operator which is linear in λ , and \hat{V}_c^λ is the correlation potential which starts at second order in λ , i.e. $\hat{V}_c^\lambda = \lambda^2 \hat{V}_c^{(2)} + \dots$. Using a complete set of orthonormal eigenfunctions Φ_n and eigenvalues \mathcal{E}_n of the KS Hamiltonian, $\hat{H}_s |\Phi_n\rangle = \mathcal{E}_n |\Phi_n\rangle$, the normalized ground-state wave function of the Hamiltonian \hat{H}^λ can be expanded as $\Psi^\lambda = \Phi + \lambda \Psi^{(1)} + \dots$ where $\Phi = \Phi_0$ is the ground-state KS single-determinant wave function and $\Psi^{(1)}$ is its first-order correction given by

$$|\Psi^{(1)}\rangle = - \sum_{n \neq 0} \frac{\langle \Phi_n | \hat{W}_{ee} - \hat{V}_{Hx} | \Phi \rangle}{\mathcal{E}_n - \mathcal{E}_0} |\Phi_n\rangle. \quad (4.8)$$

Using the expression in Eq. (2.24), the correlation energy functional can also be expanded in powers of λ

$$\begin{aligned} E_c^\lambda &= \langle \Psi^\lambda | \hat{T} + \lambda \hat{W}_{ee} | \Psi^\lambda \rangle - \langle \Phi | \hat{T} + \lambda \hat{W}_{ee} | \Phi \rangle \\ &= E_c^{(0)} + \lambda E_c^{(1)} + \lambda^2 E_c^{(2)} + \dots \end{aligned} \quad (4.9)$$

Since $\Psi^{\lambda=0} = \Phi$, the zeroth-order term vanishes: $E_c^{(0)} = 0$. Using the expression of the first-order derivative of E_c^λ with respect to λ in Eq. (2.25), i.e. $\partial E_c^\lambda / \partial \lambda = \langle \Psi^\lambda | \hat{W}_{ee} | \Psi^\lambda \rangle - \langle \Phi | \hat{W}_{ee} | \Phi \rangle$, we find that the first-order term vanishes as well: $E_c^{(1)} = 0$. The second-order term corresponds to the second-order Görling-Levy (GL2) correlation energy and is given by

$$E_c^{(2)} = E_c^{\text{GL2}} = \langle \Phi | \hat{W}_{ee} | \Psi^{(1)} \rangle = \langle \Phi | \hat{W}_{ee} - \hat{V}_{\text{Hx}} | \Psi^{(1)} \rangle, \quad (4.10)$$

where the second equality comes from the fact that $\langle \Phi | \hat{V}_{\text{Hx}} | \Psi^{(1)} \rangle = 0$ since it is the derivative with respect to λ at $\lambda = 0$ of $\langle \Psi^\lambda | \hat{V}_{\text{Hx}} | \Psi^\lambda \rangle = \int v_{\text{Hx}}(\mathbf{r}) n(\mathbf{r}) d\mathbf{r}$ which does not depend on λ by virtue of the fact the density $n(\mathbf{r})$ is constant along the adiabatic connection. Using the last expression in Eq. (4.10) allows one to express the GL2 correlation energy as

$$E_c^{\text{GL2}} = - \sum_{n \neq 0} \frac{|\langle \Phi | \hat{W}_{ee} - \hat{V}_{\text{Hx}} | \Phi_n \rangle|^2}{\mathcal{E}_n - \mathcal{E}_0}. \quad (4.11)$$

It is instructive to decompose the GL2 correlation energy as

$$E_c^{\text{GL2}} = E_c^{\text{MP2}} + E_c^{\text{S}}, \quad (4.12)$$

where E_c^{MP2} is a MP2-like correlation energy evaluated with KS spin orbitals

$$E_c^{\text{MP2}} = -\frac{1}{4} \sum_{i=1}^N \sum_{j=1}^N \sum_{a \geq N+1} \sum_{b \geq N+1} \frac{|\langle \psi_i \psi_j | \psi_a \psi_b \rangle|^2}{\varepsilon_a + \varepsilon_b - \varepsilon_i - \varepsilon_j}, \quad (4.13)$$

and E_c^{S} is the contribution coming from the single excitations (which does not vanish here, contrary to HF-based MP2 perturbation theory)

$$E_c^{\text{S}} = - \sum_{i=1}^N \sum_{a \geq N+1} \frac{|\langle \psi_i | \hat{V}_x^{\text{HF}} - \hat{V}_x | \psi_a \rangle|^2}{\varepsilon_a - \varepsilon_i}, \quad (4.14)$$

involving the difference between the integrals over the nonlocal HF exchange potential $\langle \psi_i | \hat{V}_x^{\text{HF}} | \psi_a \rangle = - \sum_{j=1}^N \langle \psi_i \psi_j | \psi_j \psi_a \rangle$ and over the local KS exchange potential $\langle \psi_i | \hat{V}_x | \psi_a \rangle = \int \psi_i^*(\mathbf{x}) v_x(\mathbf{r}) \psi_a(\mathbf{x}) d\mathbf{x}$.

Exercise 22 : Derive Eqs. (4.12)-(4.14).

Calculations of the GL2 correlation energy using either a non-self-consistent post-EXX implementation or a more complicated OEP self-consistent procedure have been tested (see, e.g., Refs. [118–120]) but the results are often disappointing. It is preferable to go beyond second order with the random-phase approximation in the adiabatic-connection fluctuation-dissipation approach.

4.3 Adiabatic-connection fluctuation-dissipation approach

4.3.1 Exact adiabatic-connection fluctuation-dissipation expression

Using the *adiabatic-connection* formula of Eq. (2.26), the correlation energy functional can be written as

$$\begin{aligned} E_c &= \int_0^1 d\lambda \langle \Psi^\lambda | \hat{W}_{ee} | \Psi^\lambda \rangle - \langle \Phi | \hat{W}_{ee} | \Phi \rangle \\ &= \frac{1}{2} \int_0^1 d\lambda \iint \frac{n_{2,c}^\lambda(\mathbf{r}_1, \mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2, \end{aligned} \quad (4.15)$$

where $n_{2,c}^\lambda(\mathbf{r}_1, \mathbf{r}_2) = n_2^\lambda(\mathbf{r}_1, \mathbf{r}_2) - n_{2,\text{KS}}(\mathbf{r}_1, \mathbf{r}_2)$ is the correlation part of the pair density along the adiabatic connection. The pair density $n_2^\lambda(\mathbf{r}_1, \mathbf{r}_2)$ can be expressed with the pair-density operator $\hat{n}_2(\mathbf{r}_1, \mathbf{r}_2)$ [Eq. (A.10)]

$$n_2^\lambda(\mathbf{r}_1, \mathbf{r}_2) = \langle \Psi^\lambda | \hat{n}_2(\mathbf{r}_1, \mathbf{r}_2) | \Psi^\lambda \rangle = \langle \Psi^\lambda | \hat{n}(\mathbf{r}_2) \hat{n}(\mathbf{r}_1) | \Psi^\lambda \rangle - \delta(\mathbf{r}_1 - \mathbf{r}_2) \langle \Psi^\lambda | \hat{n}(\mathbf{r}_1) | \Psi^\lambda \rangle, \quad (4.16)$$

where $\hat{n}(\mathbf{r})$ is the density operator [Eq. (A.8)], and the KS pair density $n_{2,\text{KS}}(\mathbf{r}_1, \mathbf{r}_2)$ simply corresponds to the case $\lambda = 0$

$$n_{2,\text{KS}}(\mathbf{r}_1, \mathbf{r}_2) = n_2^{\lambda=0}(\mathbf{r}_1, \mathbf{r}_2) = \langle \Phi | \hat{n}(\mathbf{r}_2) \hat{n}(\mathbf{r}_1) | \Phi \rangle - \delta(\mathbf{r}_1 - \mathbf{r}_2) \langle \Phi | \hat{n}(\mathbf{r}_1) | \Phi \rangle, \quad (4.17)$$

Since the density does not change with λ , i.e. $\langle \Psi^\lambda | \hat{n}(\mathbf{r}) | \Psi^\lambda \rangle = \langle \Phi | \hat{n}(\mathbf{r}) | \Phi \rangle = n(\mathbf{r})$, the correlation pair density needed in Eq. (4.15) can thus be expressed as

$$n_{2,c}^\lambda(\mathbf{r}_1, \mathbf{r}_2) = \langle \Psi^\lambda | \hat{n}(\mathbf{r}_2) \hat{n}(\mathbf{r}_1) | \Psi^\lambda \rangle - \langle \Phi | \hat{n}(\mathbf{r}_2) \hat{n}(\mathbf{r}_1) | \Phi \rangle. \quad (4.18)$$

We would like to calculate $n_{2,c}^\lambda(\mathbf{r}_1, \mathbf{r}_2)$ without having to calculate the complicated many-body wave function Ψ^λ . For this, we consider the (time-ordered) linear-response function along the adiabatic connection¹³

$$\begin{aligned} i\chi_\lambda(\mathbf{r}_1 t_1, \mathbf{r}_2 t_2) &= \langle \Psi^\lambda | T[\hat{n}^\lambda(\mathbf{r}_1 t_1) \hat{n}^\lambda(\mathbf{r}_2 t_2)] | \Psi^\lambda \rangle - \langle \Psi^\lambda | \hat{n}^\lambda(\mathbf{r}_1 t_1) | \Psi^\lambda \rangle \langle \Psi^\lambda | \hat{n}^\lambda(\mathbf{r}_2 t_2) | \Psi^\lambda \rangle \\ &= \langle \Psi^\lambda | T[\hat{n}^\lambda(\mathbf{r}_1 t_1) \hat{n}^\lambda(\mathbf{r}_2 t_2)] | \Psi^\lambda \rangle - \langle \Psi^\lambda | \hat{n}(\mathbf{r}_1) | \Psi^\lambda \rangle \langle \Psi^\lambda | \hat{n}(\mathbf{r}_2) | \Psi^\lambda \rangle, \end{aligned} \quad (4.19)$$

where $\hat{n}^\lambda(\mathbf{r}t) = e^{i\hat{H}^\lambda t} \hat{n}(\mathbf{r}) e^{-i\hat{H}^\lambda t}$ is the density operator in the Heisenberg picture, and T is the Wick time-ordering operator which orders the operators with larger times on the left, i.e. $T[\hat{n}^\lambda(\mathbf{r}_1 t_1) \hat{n}^\lambda(\mathbf{r}_2 t_2)] = \theta(t_1 - t_2) \hat{n}^\lambda(\mathbf{r}_1 t_1) \hat{n}^\lambda(\mathbf{r}_2 t_2) + \theta(t_2 - t_1) \hat{n}^\lambda(\mathbf{r}_2 t_2) \hat{n}^\lambda(\mathbf{r}_1 t_1)$ where θ is the Heaviside step function. Due to time translation invariance, the linear-response function depends in fact only on $\tau = t_1 - t_2$. If we set $t_2 = t_1 + 0^+$ where 0^+ is an infinitesimal positive shift, i.e. $\tau = 0^-$, we get

$$i\chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \tau = 0^-) = \langle \Psi^\lambda | \hat{n}(\mathbf{r}_2) \hat{n}(\mathbf{r}_1) | \Psi^\lambda \rangle - \langle \Psi^\lambda | \hat{n}(\mathbf{r}_1) | \Psi^\lambda \rangle \langle \Psi^\lambda | \hat{n}(\mathbf{r}_2) | \Psi^\lambda \rangle, \quad (4.20)$$

and, similarly, for the non-interacting KS linear-response function corresponding to the case $\lambda = 0$

$$i\chi_0(\mathbf{r}_1, \mathbf{r}_2; \tau = 0^-) = \langle \Phi | \hat{n}(\mathbf{r}_2) \hat{n}(\mathbf{r}_1) | \Phi \rangle - \langle \Phi | \hat{n}(\mathbf{r}_1) | \Phi \rangle \langle \Phi | \hat{n}(\mathbf{r}_2) | \Phi \rangle. \quad (4.21)$$

We can thus express $n_{2,c}^\lambda(\mathbf{r}_1, \mathbf{r}_2)$ as the difference between the zero-time linear-response functions

$$n_{2,c}^\lambda(\mathbf{r}_1, \mathbf{r}_2) = i[\chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \tau = 0^-) - \chi_0(\mathbf{r}_1, \mathbf{r}_2; \tau = 0^-)]. \quad (4.22)$$

Alternatively, Eq (4.22) can be rewritten in terms of the Fourier transforms of the linear-response functions, using the definition $\chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \tau) = \int_{-\infty}^{+\infty} d\omega / (2\pi) \chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \omega) e^{-i\omega\tau}$,

$$\boxed{n_{2,c}^\lambda(\mathbf{r}_1, \mathbf{r}_2) = - \int_{-\infty}^{+\infty} \frac{d\omega}{2\pi i} e^{i\omega 0^+} [\chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \omega) - \chi_0(\mathbf{r}_1, \mathbf{r}_2; \omega)],} \quad (4.23)$$

which is known as the (zero-temperature) *fluctuation-dissipation theorem*. This theorem relates ground-state correlations in the time-independent system, $n_{2,c}^\lambda(\mathbf{r}_1, \mathbf{r}_2)$, to the linear response of the system due to a time-dependent external perturbation, $\chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \omega)$.

¹³We choose to work with the *time-ordered* linear-response function (or polarization propagator) which is used in many-body Green function theory. In linear-response time-dependent DFT, one normally uses the *retarded* linear-response function $i\chi_\lambda^R(\mathbf{r}_1 t_1, \mathbf{r}_2 t_2) = \theta(t_1 - t_2) \langle \Psi^\lambda | [\hat{n}^\lambda(\mathbf{r}_1 t_1), \hat{n}^\lambda(\mathbf{r}_2 t_2)] | \Psi^\lambda \rangle - \langle \Psi^\lambda | \hat{n}^\lambda(\mathbf{r}_1 t_1) | \Psi^\lambda \rangle \langle \Psi^\lambda | \hat{n}^\lambda(\mathbf{r}_2 t_2) | \Psi^\lambda \rangle$. Their Fourier transforms are related by $\chi_\lambda^R(\mathbf{r}_1, \mathbf{r}_2; \omega) = \chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \omega)$ for $\omega \geq 0$, and $\chi_\lambda^R(\mathbf{r}_1, \mathbf{r}_2; \omega) = \chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \omega)^*$ for $\omega < 0$. One should be careful to be consistent when switching between these related quantities. For a presentation based on the retarded linear-response function, see e.g. Ref. [17].

Exercise 23 : Show that the linear-response function can be rewritten as

$$i\chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \tau) = \theta(\tau) \sum_{n \neq 0} \langle \Psi^\lambda | \hat{n}(\mathbf{r}_1) | \Psi_n^\lambda \rangle \langle \Psi_n^\lambda | \hat{n}(\mathbf{r}_2) | \Psi^\lambda \rangle e^{-i\omega_n^\lambda \tau} \\ + \theta(-\tau) \sum_{n \neq 0} \langle \Psi^\lambda | \hat{n}(\mathbf{r}_2) | \Psi_n^\lambda \rangle \langle \Psi_n^\lambda | \hat{n}(\mathbf{r}_1) | \Psi^\lambda \rangle e^{i\omega_n^\lambda \tau}, \quad (4.24)$$

where the sums are over all eigenstates Ψ_n^λ of the Hamiltonian \hat{H}^λ , i.e. $\hat{H}^\lambda | \Psi_n^\lambda \rangle = E_n^\lambda | \Psi_n^\lambda \rangle$, except the ground state $\Psi^\lambda = \Psi_0^\lambda$, and $\omega_n^\lambda = E_n^\lambda - E_0^\lambda$ are the corresponding excitation energies. Show then that the Fourier transform of $\chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \tau)$ is

$$\chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \omega) = \sum_{n \neq 0} \frac{\langle \Psi^\lambda | \hat{n}(\mathbf{r}_1) | \Psi_n^\lambda \rangle \langle \Psi_n^\lambda | \hat{n}(\mathbf{r}_2) | \Psi^\lambda \rangle}{\omega - \omega_n^\lambda + i0^+} - \frac{\langle \Psi^\lambda | \hat{n}(\mathbf{r}_2) | \Psi_n^\lambda \rangle \langle \Psi_n^\lambda | \hat{n}(\mathbf{r}_1) | \Psi^\lambda \rangle}{\omega + \omega_n^\lambda - i0^+}, \quad (4.25)$$

which is called the Lehmann representation of χ_λ . Finally, check directly Eq. (4.23) by performing the integration over ω using the residue theorem.

Combining Eqs. (4.15) and (4.23), we finally arrive the exact *adiabatic-connection fluctuation-dissipation* (ACFD) formula for the correlation energy

$$E_c = -\frac{1}{2} \int_0^1 d\lambda \int_{-\infty}^{+\infty} \frac{d\omega}{2\pi i} e^{i\omega 0^+} \iint \frac{\chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \omega) - \chi_0(\mathbf{r}_1, \mathbf{r}_2; \omega)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2. \quad (4.26)$$

The usefulness of the ACFD formula is due to the fact there are practical ways of directly calculating $\chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \omega)$ without having to calculate the many-body wave function Ψ^λ . In linear-response time-dependent DFT, one can find a Dyson-like equation for $\chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \omega)$

$$\chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \omega) = \chi_0(\mathbf{r}_1, \mathbf{r}_2; \omega) + \iint \chi_0(\mathbf{r}_1, \mathbf{r}_3; \omega) f_{\text{Hxc}}^\lambda(\mathbf{r}_3, \mathbf{r}_4; \omega) \chi_\lambda(\mathbf{r}_4, \mathbf{r}_2; \omega) d\mathbf{r}_3 d\mathbf{r}_4, \quad (4.27)$$

where $f_{\text{Hxc}}^\lambda(\mathbf{r}_3, \mathbf{r}_4; \omega)$ is the *Hartree-exchange-correlation kernel* associated to the Hamiltonian \hat{H}^λ . Here, Eq. (4.27) will be considered as the definition for f_{Hxc}^λ . In principle, the exact correlation energy can be obtained with Eqs. (4.26) and (4.27). In practice, however, we need to use an approximation for f_{Hxc}^λ .

4.3.2 Direct random-phase approximation

In the *direct random-phase approximation* (dRPA, also just referred to as RPA, or sometimes as time-dependent Hartree), only the Hartree part of the kernel, which is linear in λ and independent from ω , is retained

$$f_{\text{Hxc}}^\lambda(\mathbf{r}_1, \mathbf{r}_2; \omega) \approx f_{\text{H}}^\lambda(\mathbf{r}_1, \mathbf{r}_2) = \lambda w_{\text{ee}}(\mathbf{r}_1, \mathbf{r}_2), \quad (4.28)$$

where $w_{\text{ee}}(\mathbf{r}_1, \mathbf{r}_2) = 1/|\mathbf{r}_1 - \mathbf{r}_2|$ is the Coulomb interaction, and the corresponding dRPA linear-response function then satisfies the equation

$$\chi_\lambda^{\text{dRPA}}(\mathbf{r}_1, \mathbf{r}_2; \omega) = \chi_0(\mathbf{r}_1, \mathbf{r}_2; \omega) + \lambda \iint \chi_0(\mathbf{r}_1, \mathbf{r}_3; \omega) w_{\text{ee}}(\mathbf{r}_3, \mathbf{r}_4) \chi_\lambda^{\text{dRPA}}(\mathbf{r}_4, \mathbf{r}_2; \omega) d\mathbf{r}_3 d\mathbf{r}_4. \quad (4.29)$$

The physical contents of this approximation can be seen by iterating Eq. (4.29) which generates an infinite series

$$\begin{aligned} \chi_\lambda^{\text{dRPA}}(\mathbf{r}_1, \mathbf{r}_2; \omega) &= \chi_0(\mathbf{r}_1, \mathbf{r}_2; \omega) + \lambda \iint \chi_0(\mathbf{r}_1, \mathbf{r}_3; \omega) w_{ee}(\mathbf{r}_3, \mathbf{r}_4) \chi_0(\mathbf{r}_4, \mathbf{r}_2; \omega) d\mathbf{r}_3 d\mathbf{r}_4 \\ &+ \lambda^2 \iiint \chi_0(\mathbf{r}_1, \mathbf{r}_3; \omega) w_{ee}(\mathbf{r}_3, \mathbf{r}_4) \chi_0(\mathbf{r}_4, \mathbf{r}_5; \omega) w_{ee}(\mathbf{r}_5, \mathbf{r}_6) \chi_0(\mathbf{r}_6, \mathbf{r}_2; \omega) d\mathbf{r}_3 d\mathbf{r}_4 d\mathbf{r}_5 d\mathbf{r}_6 + \dots, \end{aligned} \quad (4.30)$$

which, after plugging it into Eq. (4.26), leads to the dRPA correlation energy as the following perturbation expansion

$$\begin{aligned} E_c^{\text{dRPA}} &= -\frac{1}{2} \int_0^1 d\lambda \int_{-\infty}^{+\infty} \frac{d\omega}{2\pi i} e^{i\omega 0^+} \left[\lambda \iiint \frac{\chi_0(\mathbf{r}_1, \mathbf{r}_3; \omega) \chi_0(\mathbf{r}_4, \mathbf{r}_2; \omega)}{|\mathbf{r}_1 - \mathbf{r}_2| |\mathbf{r}_3 - \mathbf{r}_4|} d\mathbf{r}_1 d\mathbf{r}_2 d\mathbf{r}_3 d\mathbf{r}_4 \right. \\ &\left. + \lambda^2 \iiint \frac{\chi_0(\mathbf{r}_1, \mathbf{r}_3; \omega) \chi_0(\mathbf{r}_4, \mathbf{r}_5; \omega) \chi_0(\mathbf{r}_6, \mathbf{r}_2; \omega)}{|\mathbf{r}_1 - \mathbf{r}_2| |\mathbf{r}_3 - \mathbf{r}_4| |\mathbf{r}_5 - \mathbf{r}_6|} d\mathbf{r}_1 d\mathbf{r}_2 d\mathbf{r}_3 d\mathbf{r}_4 d\mathbf{r}_5 d\mathbf{r}_6 + \dots \right]. \end{aligned} \quad (4.31)$$

Using now the Lehmann representation [Eq. (4.25)] of the KS linear-response function in terms of the KS orbitals and their energies

$$\chi_0(\mathbf{r}_1, \mathbf{r}_2; \omega) = \sum_{\sigma \in \{\uparrow, \downarrow\}} \sum_{i=1}^{N_\sigma} \sum_{a \geq N_\sigma + 1} \left[\frac{\varphi_{i\sigma}^*(\mathbf{r}_1) \varphi_{a\sigma}(\mathbf{r}_1) \varphi_{a\sigma}^*(\mathbf{r}_2) \varphi_{i\sigma}(\mathbf{r}_2)}{\omega - (\varepsilon_a - \varepsilon_i) + i0^+} - \frac{\varphi_{i\sigma}^*(\mathbf{r}_2) \varphi_{a\sigma}(\mathbf{r}_2) \varphi_{a\sigma}^*(\mathbf{r}_1) \varphi_{i\sigma}(\mathbf{r}_1)}{\omega + (\varepsilon_a - \varepsilon_i) - i0^+} \right], \quad (4.32)$$

one can obtain, after quite some work,

$$\begin{aligned} E_c^{\text{dRPA}} &= -\frac{1}{2} \sum_{i=1}^N \sum_{j=1}^N \sum_{a \geq N+1} \sum_{b \geq N+1} \frac{|\langle \psi_i \psi_j | \psi_a \psi_b \rangle|^2}{\varepsilon_a + \varepsilon_b - \varepsilon_i - \varepsilon_j} \\ &+ \sum_{i=1}^N \sum_{j=1}^N \sum_{k=1}^N \sum_{a \geq N+1} \sum_{b \geq N+1} \sum_{c \geq N+1} \frac{\langle \psi_i \psi_j | \psi_a \psi_b \rangle \langle \psi_j \psi_k | \psi_b \psi_c \rangle \langle \psi_k \psi_i | \psi_c \psi_a \rangle}{(\varepsilon_a + \varepsilon_b - \varepsilon_i - \varepsilon_j)(\varepsilon_a + \varepsilon_c - \varepsilon_i - \varepsilon_k)} + \dots \end{aligned} \quad (4.33)$$

The dRPA correlation energy is the sum of all the direct terms (i.e., no exchange terms) of the Møller-Plesset or Görling-Levy perturbation expansion. Of course, Eq. (4.33) is not the way to calculate the dRPA correlation energy in practice. To do this, we need to solve the Dyson equation [Eq. (4.29)] without explicitly expanding in powers of λ .

4.3.3 Practical calculation in a spin orbital basis

For solving Eq. (4.29) in a spin orbital basis, it is more convenient to introduce the four-point (time-ordered) linear-response function

$$\begin{aligned} i\chi_\lambda(\mathbf{x}_1, \mathbf{x}_2; \mathbf{x}'_1, \mathbf{x}'_2; \tau = t_1 - t_2) &= \langle \Psi^\lambda | T[\hat{n}_1^\lambda(\mathbf{x}_1, \mathbf{x}'_1; t_1) \hat{n}_1^\lambda(\mathbf{x}_2, \mathbf{x}'_2; t_2)] | \Psi^\lambda \rangle \\ &- \langle \Psi^\lambda | \hat{n}_1^\lambda(\mathbf{x}_1, \mathbf{x}'_1; t_1) | \Psi^\lambda \rangle \langle \Psi^\lambda | \hat{n}_1^\lambda(\mathbf{x}_2, \mathbf{x}'_2; t_2) | \Psi^\lambda \rangle, \end{aligned} \quad (4.34)$$

where $\hat{n}_1^\lambda(\mathbf{x}, \mathbf{x}'; t) = e^{i\hat{H}^\lambda t} \hat{n}_1(\mathbf{x}, \mathbf{x}') e^{-i\hat{H}^\lambda t}$ is the (spin-dependent) one-electron density matrix operator in the Heisenberg picture. The Fourier transform of the linear-response function

$\chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \omega)$ involved in Eq. (4.26) is just the spin-summed diagonal part of the Fourier transform of the four-point linear-response function

$$\chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \omega) = \sum_{\sigma_1 \in \{\uparrow, \downarrow\}} \sum_{\sigma_2 \in \{\uparrow, \downarrow\}} \chi_\lambda(\mathbf{r}_1 \sigma_1, \mathbf{r}_2 \sigma_2; \mathbf{r}_1 \sigma_1, \mathbf{r}_2 \sigma_2; \omega). \quad (4.35)$$

In the dRPA approximation, the four-point linear-response function is given by

$$\begin{aligned} \chi_\lambda^{\text{dRPA}}(\mathbf{x}_1, \mathbf{x}_2; \mathbf{x}'_1, \mathbf{x}'_2; \omega) &= \chi_0(\mathbf{x}_1, \mathbf{x}_2; \mathbf{x}'_1, \mathbf{x}'_2; \omega) \\ &+ \iiint \! \! \! \int d\mathbf{x}_3 d\mathbf{x}_4 d\mathbf{x}_5 d\mathbf{x}_6 \chi_0(\mathbf{x}_1, \mathbf{x}_4; \mathbf{x}'_1, \mathbf{x}_3; \omega) f_{\text{H}}^\lambda(\mathbf{x}_3, \mathbf{x}_6; \mathbf{x}_4, \mathbf{x}_5) \chi_\lambda^{\text{dRPA}}(\mathbf{x}_5, \mathbf{x}_2; \mathbf{x}_6, \mathbf{x}'_2; \omega), \end{aligned} \quad (4.36)$$

with the four-point Hartree kernel $f_{\text{H}}^\lambda(\mathbf{x}_1, \mathbf{x}_2; \mathbf{x}'_1, \mathbf{x}'_2) = \lambda w_{\text{ee}}(|\mathbf{r}_1 - \mathbf{r}_2|) \delta(\mathbf{x}_1 - \mathbf{x}'_1) \delta(\mathbf{x}_2 - \mathbf{x}'_2)$, or equivalently in terms of inverses

$$(\chi_\lambda^{\text{dRPA}})^{-1}(\mathbf{x}_1, \mathbf{x}_2; \mathbf{x}'_1, \mathbf{x}'_2; \omega) = \chi_0^{-1}(\mathbf{x}_1, \mathbf{x}_2; \mathbf{x}'_1, \mathbf{x}'_2; \omega) - f_{\text{H}}^\lambda(\mathbf{x}_1, \mathbf{x}_2; \mathbf{x}'_1, \mathbf{x}'_2). \quad (4.37)$$

The non-interacting four-point linear-response function can be explicitly written in terms of the KS spin orbitals and their energies

$$\chi_0(\mathbf{x}_1, \mathbf{x}_2; \mathbf{x}'_1, \mathbf{x}'_2; \omega) = \sum_{i=1}^N \sum_{a \geq N+1} \left[\frac{\psi_i^*(\mathbf{x}'_1) \psi_a(\mathbf{x}_1) \psi_a^*(\mathbf{x}'_2) \psi_i(\mathbf{x}_2)}{\omega - (\varepsilon_a - \varepsilon_i) + i0^+} - \frac{\psi_i^*(\mathbf{x}'_2) \psi_a(\mathbf{x}_2) \psi_a^*(\mathbf{x}'_1) \psi_i(\mathbf{x}_1)}{\omega + (\varepsilon_a - \varepsilon_i) - i0^+} \right], \quad (4.38)$$

which can be seen as expanded in a basis, $\chi_0(\mathbf{x}_1, \mathbf{x}_2; \mathbf{x}'_1, \mathbf{x}'_2; \omega) = \sum_{p,q} [\chi_0(\omega)]_{p,q} f_p(\mathbf{x}_1, \mathbf{x}'_1) f_q^*(\mathbf{x}_2, \mathbf{x}'_2)$, where the basis functions are tensor products of two occupied/virtual (ov) $f_{ia}(\mathbf{x}_1, \mathbf{x}'_1) = \psi_i^*(\mathbf{x}'_1) \psi_a(\mathbf{x}_1)$ or two virtual/occupied (vo) spin orbitals $f_{ai}(\mathbf{x}_1, \mathbf{x}'_1) = \psi_a^*(\mathbf{x}'_1) \psi_i(\mathbf{x}_1)$, and the matrix of $\chi_0(\omega)$ is in fact diagonal: $[\chi_0(\omega)]_{ia,jb} = \delta_{ij} \delta_{ab} / (\omega - (\varepsilon_a - \varepsilon_i) + i0^+)$, $[\chi_0(\omega)]_{ai,bj} = \delta_{ij} \delta_{ab} / (\omega + (\varepsilon_a - \varepsilon_i) - i0^+)$, and $[\chi_0(\omega)]_{ia,bj} = [\chi_0(\omega)]_{ai,jb} = 0$. The inverse of $\chi_0(\omega)$ in this basis is, written with ov/ov, ov/vo, vo/ov, ov/ov block matrices,

$$\chi_0^{-1}(\omega) = - \left[\begin{pmatrix} \Delta\varepsilon & \mathbf{0} \\ \mathbf{0} & \Delta\varepsilon \end{pmatrix} - \omega \begin{pmatrix} \mathbf{1} & \mathbf{0} \\ \mathbf{0} & -\mathbf{1} \end{pmatrix} \right], \quad (4.39)$$

with the diagonal matrix $\Delta\varepsilon_{ia,jb} = (\varepsilon_a - \varepsilon_i) \delta_{ij} \delta_{ab}$. Using Eq. (4.37), the matrix representation of the inverse of $\chi_\lambda^{\text{dRPA}}(\omega)$ can be easily find

$$(\chi_\lambda^{\text{dRPA}})^{-1}(\omega) = - \left[\begin{pmatrix} \mathbf{A}_\lambda & \mathbf{B}_\lambda \\ \mathbf{B}_\lambda^* & \mathbf{A}_\lambda^* \end{pmatrix} - \omega \begin{pmatrix} \mathbf{1} & \mathbf{0} \\ \mathbf{0} & -\mathbf{1} \end{pmatrix} \right], \quad (4.40)$$

with the matrices \mathbf{A}_λ and \mathbf{B}_λ defined by

$$(A_\lambda)_{ia,jb} = \Delta\varepsilon_{ia,jb} + \lambda \langle \psi_a \psi_j | \psi_i \psi_b \rangle, \quad (4.41a)$$

$$(B_\lambda)_{ia,jb} = \lambda \langle \psi_a \psi_b | \psi_i \psi_j \rangle, \quad (4.41b)$$

and it can be checked that the matrix \mathbf{A}_λ is Hermitian [i.e., $(A_\lambda)_{jb,ia} = (A_\lambda)_{ia,jb}^*$] and the matrix \mathbf{B}_λ is symmetric [i.e., $(B_\lambda)_{jb,ia} = (B_\lambda)_{ia,jb}$]. To calculate the inverse of the matrix in the right-hand-side of Eq. (4.40), we then consider the following generalized eigenvalue equation

$$\boxed{\begin{pmatrix} \mathbf{A}_\lambda & \mathbf{B}_\lambda \\ \mathbf{B}_\lambda^* & \mathbf{A}_\lambda^* \end{pmatrix} \begin{pmatrix} \mathbf{X}_{n,\lambda} \\ \mathbf{Y}_{n,\lambda} \end{pmatrix} = \omega_n^\lambda \begin{pmatrix} \mathbf{1} & \mathbf{0} \\ \mathbf{0} & -\mathbf{1} \end{pmatrix} \begin{pmatrix} \mathbf{X}_{n,\lambda} \\ \mathbf{Y}_{n,\lambda} \end{pmatrix},} \quad (4.42)$$

whose solutions come in pairs: if $(\mathbf{X}_{n,\lambda}, \mathbf{Y}_{n,\lambda})$ is an eigenvector with eigenvalue ω_n^λ , then it is easy to check that $(\mathbf{Y}_{n,\lambda}^*, \mathbf{X}_{n,\lambda}^*)$ is also an eigenvector with opposite eigenvalue $-\omega_n^\lambda$. Choosing the normalization of the eigenvectors so that $\mathbf{X}_{n,\lambda}^\dagger \mathbf{X}_{m,\lambda} - \mathbf{Y}_{n,\lambda}^\dagger \mathbf{Y}_{m,\lambda} = \delta_{nm}$, the matrix $\chi_\lambda^{\text{dRPA}}(\omega)$ can be expressed as the following spectral representation

$$\chi_\lambda^{\text{dRPA}}(\omega) = \sum_n \left[\frac{1}{\omega - \omega_n^\lambda + i0^+} \begin{pmatrix} \mathbf{X}_{n,\lambda} \\ \mathbf{Y}_{n,\lambda} \end{pmatrix} \begin{pmatrix} \mathbf{X}_{n,\lambda}^\dagger & \mathbf{Y}_{n,\lambda}^\dagger \end{pmatrix} - \frac{1}{\omega + \omega_n^\lambda - i0^+} \begin{pmatrix} \mathbf{Y}_{n,\lambda}^* \\ \mathbf{X}_{n,\lambda}^* \end{pmatrix} \begin{pmatrix} \mathbf{Y}_{n,\lambda}^{*\dagger} & \mathbf{X}_{n,\lambda}^{*\dagger} \end{pmatrix} \right], \quad (4.43)$$

where the sum is over eigenvectors with positive eigenvalues. The fluctuation-dissipation theorem [Eq. (4.23)] leads to the matrix representation of the correlation part of the dRPA two-particle density matrix $\mathbf{n}_{2,c}^{\lambda, \text{dRPA}}$ (using contour integration in the upper half of the complex plane)

$$\begin{aligned} \mathbf{n}_{2,c}^{\lambda, \text{dRPA}} &= - \int_{-\infty}^{+\infty} \frac{d\omega}{2\pi i} e^{i\omega 0^+} [\chi_\lambda(\omega) - \chi_0(\omega)] \\ &= \sum_n \begin{pmatrix} \mathbf{Y}_{n,\lambda}^* & \mathbf{Y}_{n,\lambda}^{*\dagger} & \mathbf{Y}_{n,\lambda}^* & \mathbf{X}_{n,\lambda}^{*\dagger} \\ \mathbf{X}_{n,\lambda}^* & \mathbf{Y}_{n,\lambda}^{*\dagger} & \mathbf{X}_{n,\lambda}^* & \mathbf{X}_{n,\lambda}^{*\dagger} \end{pmatrix} - \begin{pmatrix} \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{1} \end{pmatrix}, \end{aligned} \quad (4.44)$$

the simple contribution coming from $\chi_0(\omega)$ resulting from its diagonal form, and the dRPA correlation energy has then the following expression

$$\begin{aligned} E_c^{\text{dRPA}} &= \frac{1}{2} \int_0^1 d\lambda \sum_{i=1}^N \sum_{j=1}^N \sum_{a \geq N+1} \sum_{b \geq N+1} \sum_n \left\{ \langle \psi_i \psi_b | \psi_a \psi_j \rangle (Y_{n,\lambda})_{ia}^* (Y_{n,\lambda})_{jb} \right. \\ &\quad + \langle \psi_i \psi_j | \psi_a \psi_b \rangle (Y_{n,\lambda})_{ia}^* (X_{n,\lambda})_{jb} + \langle \psi_a \psi_b | \psi_i \psi_j \rangle (X_{n,\lambda})_{ia}^* (Y_{n,\lambda})_{jb} \\ &\quad \left. + \langle \psi_a \psi_j | \psi_i \psi_b \rangle [(X_{n,\lambda})_{ia}^* (X_{n,\lambda})_{jb} - \delta_{ij} \delta_{ab}] \right\}. \end{aligned} \quad (4.45)$$

For real-valued spin orbitals, the correlation energy can be simplified to

$$E_c^{\text{dRPA}} = \frac{1}{2} \int_0^1 d\lambda \sum_{i=1}^N \sum_{j=1}^N \sum_{a \geq N+1} \sum_{b \geq N+1} \langle \psi_i \psi_b | \psi_a \psi_j \rangle (P_{c,\lambda})_{ia,jb}, \quad (4.46)$$

where

$$(P_{c,\lambda})_{ia,jb} = \sum_n (X_{n,\lambda} + Y_{n,\lambda})_{ia} (X_{n,\lambda} + Y_{n,\lambda})_{jb} - \delta_{ij} \delta_{ab}, \quad (4.47)$$

or, in matrix form,

$$\mathbf{P}_{c,\lambda} = \sum_n (\mathbf{X}_{n,\lambda} + \mathbf{Y}_{n,\lambda}) (\mathbf{X}_{n,\lambda} + \mathbf{Y}_{n,\lambda})^\top - \mathbf{1}. \quad (4.48)$$

Using the fact that, if $\mathbf{A}_\lambda + \mathbf{B}_\lambda$ and $\mathbf{A}_\lambda - \mathbf{B}_\lambda$ are positive definite, the non-Hermitian eigenvalue equation (4.42) with real spin orbitals can be transformed into the following half-size symmetric eigenvalue equation

$$\mathbf{M}_\lambda \mathbf{Z}_{n,\lambda} = (\omega_n^\lambda)^2 \mathbf{Z}_{n,\lambda}, \quad (4.49)$$

where $\mathbf{M}_\lambda = (\mathbf{A}_\lambda - \mathbf{B}_\lambda)^{1/2} (\mathbf{A}_\lambda + \mathbf{B}_\lambda) (\mathbf{A}_\lambda - \mathbf{B}_\lambda)^{1/2}$ and with eigenvectors $\mathbf{Z}_{n,\lambda} = \sqrt{\omega_n^\lambda} (\mathbf{A}_\lambda - \mathbf{B}_\lambda)^{-1/2} (\mathbf{X}_{n,\lambda} + \mathbf{Y}_{n,\lambda})$, and using the spectral decomposition $\mathbf{M}_\lambda^{-1/2} =$

$\sum_n (\omega_n^\lambda)^{-1} \mathbf{Z}_{n,\lambda} \mathbf{Z}_{n,\lambda}^\top$, the correlation two-particle density matrix $\mathbf{P}_{c,\lambda}$ can be finally expressed as

$$\mathbf{P}_{c,\lambda} = (\mathbf{A}_\lambda - \mathbf{B}_\lambda)^{1/2} \mathbf{M}_\lambda^{-1/2} (\mathbf{A}_\lambda - \mathbf{B}_\lambda)^{1/2} - \mathbf{1}. \quad (4.50)$$

In practice, the integration over λ in Eq. (4.46) is done by numerical integration.

This is the *adiabatic-connection formulation* of dRPA. There are also other equivalent formulations of dRPA in which the integration over λ is done analytically: the *plasmon-formula formulation*, the *ring coupled-cluster formulation*, and the *dielectric-matrix formulation*. For more details about all the different formulations, see e.g. Refs. [91, 121–126].

Most dRPA correlation energy (combined with the EXX energy) calculations are done in a non-self-consistent way but self-consistent OEP dRPA calculations have also recently been performed [127, 128]. One of the main advantages of dRPA is that it accounts for van der Waals dispersion interactions, and it can somewhat deal with systems with static correlation. However, it shows large self-interaction errors. To overcome the latter drawback, the exchange contribution to the kernel needs to be included.

4.3.4 Random-phase approximation with exchange and beyond

The next logical improvement over dRPA is then to include exchange terms by adding the EXX kernel which is also linear in λ but depends on the frequency

$$f_{\text{Hxc}}^\lambda(\mathbf{r}_1, \mathbf{r}_2; \omega) \approx f_{\text{Hx}}^\lambda(\mathbf{r}_1, \mathbf{r}_2; \omega) = \lambda w_{\text{ee}}(\mathbf{r}_1, \mathbf{r}_2) + \lambda f_{\text{x}}(\mathbf{r}_1, \mathbf{r}_2; \omega), \quad (4.51)$$

and the corresponding RPax(EXX) [129–132] linear-response function satisfies the Dyson equation

$$\chi_\lambda^{\text{RPax(EXX)}}(\mathbf{r}_1, \mathbf{r}_2; \omega) = \chi_0(\mathbf{r}_1, \mathbf{r}_2; \omega) + \iint \chi_0(\mathbf{r}_1, \mathbf{r}_3; \omega) f_{\text{Hx}}^\lambda(\mathbf{r}_3, \mathbf{r}_4; \omega) \chi_\lambda^{\text{RPax(EXX)}}(\mathbf{r}_4, \mathbf{r}_2; \omega) d\mathbf{r}_3 d\mathbf{r}_4. \quad (4.52)$$

Alternatively, one can define a RPax(HF) [89, 123] approximation at the level of the four-point linear-response function

$$\begin{aligned} \chi_\lambda^{\text{RPax(HF)}}(\mathbf{x}_1, \mathbf{x}_2; \mathbf{x}'_1, \mathbf{x}'_2; \omega) &= \chi_0(\mathbf{x}_1, \mathbf{x}_2; \mathbf{x}'_1, \mathbf{x}'_2; \omega) \\ &+ \iiint \int d\mathbf{x}_3 d\mathbf{x}_4 d\mathbf{x}_5 d\mathbf{x}_6 \chi_0(\mathbf{x}_1, \mathbf{x}_4; \mathbf{x}'_1, \mathbf{x}'_3; \omega) f_{\text{Hx}}^{\lambda, \text{HF}}(\mathbf{x}_3, \mathbf{x}_6; \mathbf{x}_4, \mathbf{x}_5) \chi_\lambda^{\text{RPax(HF)}}(\mathbf{x}_5, \mathbf{x}_2; \mathbf{x}_6, \mathbf{x}'_2; \omega), \end{aligned} \quad (4.53)$$

with the four-point Hartree-Fock kernel $f_{\text{Hx}}^{\lambda, \text{HF}}(\mathbf{x}_1, \mathbf{x}_2; \mathbf{x}'_1, \mathbf{x}'_2) = \lambda w_{\text{ee}}(|\mathbf{r}_1 - \mathbf{r}_2|) [\delta(\mathbf{x}_1 - \mathbf{x}'_1) \delta(\mathbf{x}_2 - \mathbf{x}'_2) - \delta(\mathbf{x}_1 - \mathbf{x}'_2) \delta(\mathbf{x}_2 - \mathbf{x}'_1)]$ which is linear in λ but independent from the frequency ω . However, contrary to the dRPA case, Eqs. (4.52) and (4.53) do not lead to the same correlation energies. Moreover, several non-equivalent correlation energies can be extracted from Eq. (4.53) [124].

In practice, these diverse RPax variants do not always improve over dRPA. Going beyond RPax by also including a correlation kernel is an active area of research (see e.g. Ref. [133]).

Appendices

A The Hamiltonian in second quantization

It is convenient to express the Hamiltonian operator in second quantization. In this formalism, the operators are independent from the number of electrons (i.e., we work in Fock space).

In real-space second quantization, the electronic Hamiltonian operator is written as

$$\hat{H} = \hat{T} + \hat{W}_{ee} + \hat{V}_{ne}, \quad (\text{A.1})$$

where \hat{T} is the kinetic-energy operator

$$\hat{T} = -\frac{1}{2} \sum_{\sigma \in \{\uparrow, \downarrow\}} \int \hat{\psi}_{\sigma}^{\dagger}(\mathbf{r}) \nabla^2 \hat{\psi}_{\sigma}(\mathbf{r}) d\mathbf{r}, \quad (\text{A.2})$$

\hat{W}_{ee} is the electron-electron interaction operator

$$\hat{W}_{ee} = \frac{1}{2} \sum_{\sigma_1 \in \{\uparrow, \downarrow\}} \sum_{\sigma_2 \in \{\uparrow, \downarrow\}} \iint \hat{\psi}_{\sigma_2}^{\dagger}(\mathbf{r}_2) \hat{\psi}_{\sigma_1}^{\dagger}(\mathbf{r}_1) w_{ee}(\mathbf{r}_1, \mathbf{r}_2) \hat{\psi}_{\sigma_1}(\mathbf{r}_1) \hat{\psi}_{\sigma_2}(\mathbf{r}_2) d\mathbf{r}_1 d\mathbf{r}_2, \quad (\text{A.3})$$

with $w_{ee}(\mathbf{r}_1, \mathbf{r}_2) = 1/|\mathbf{r}_1 - \mathbf{r}_2|$, and \hat{V}_{ne} is the nuclei-electron interaction operator

$$\hat{V}_{ne} = \sum_{\sigma \in \{\uparrow, \downarrow\}} \int \hat{\psi}_{\sigma}^{\dagger}(\mathbf{r}) v_{ne}(\mathbf{r}) \hat{\psi}_{\sigma}(\mathbf{r}) d\mathbf{r}. \quad (\text{A.4})$$

In these expressions, $\hat{\psi}_{\sigma}^{\dagger}(\mathbf{r})$ and $\hat{\psi}_{\sigma}(\mathbf{r})$ are the creation and annihilation field operators, respectively, which obey Fermionic anticommutation rules

$$\left\{ \hat{\psi}_{\sigma}^{\dagger}(\mathbf{r}), \hat{\psi}_{\sigma'}^{\dagger}(\mathbf{r}') \right\} = 0, \quad (\text{A.5})$$

$$\left\{ \hat{\psi}_{\sigma}(\mathbf{r}), \hat{\psi}_{\sigma'}(\mathbf{r}') \right\} = 0, \quad (\text{A.6})$$

$$\left\{ \hat{\psi}_{\sigma}^{\dagger}(\mathbf{r}), \hat{\psi}_{\sigma'}(\mathbf{r}') \right\} = \delta(\mathbf{r} - \mathbf{r}') \delta_{\sigma\sigma'}. \quad (\text{A.7})$$

It is also convenient to define the density operator

$$\hat{n}(\mathbf{r}) = \sum_{\sigma \in \{\uparrow, \downarrow\}} \hat{\psi}_{\sigma}^{\dagger}(\mathbf{r}) \hat{\psi}_{\sigma}(\mathbf{r}), \quad (\text{A.8})$$

the one-particle density-matrix operator

$$\hat{n}_1(\mathbf{r}, \mathbf{r}') = \sum_{\sigma \in \{\uparrow, \downarrow\}} \hat{\psi}_{\sigma}^{\dagger}(\mathbf{r}') \hat{\psi}_{\sigma}(\mathbf{r}), \quad (\text{A.9})$$

and the pair-density operator

$$\begin{aligned} \hat{n}_2(\mathbf{r}_1, \mathbf{r}_2) &= \sum_{\sigma_1 \in \{\uparrow, \downarrow\}} \sum_{\sigma_2 \in \{\uparrow, \downarrow\}} \hat{\psi}_{\sigma_2}^{\dagger}(\mathbf{r}_2) \hat{\psi}_{\sigma_1}^{\dagger}(\mathbf{r}_1) \hat{\psi}_{\sigma_1}(\mathbf{r}_1) \hat{\psi}_{\sigma_2}(\mathbf{r}_2) \\ &= \hat{n}(\mathbf{r}_2) \hat{n}(\mathbf{r}_1) - \hat{n}(\mathbf{r}_1) \delta(\mathbf{r}_1 - \mathbf{r}_2), \\ &= \hat{n}(\mathbf{r}_1) \hat{n}(\mathbf{r}_2) - \hat{n}(\mathbf{r}_1) \delta(\mathbf{r}_1 - \mathbf{r}_2), \end{aligned} \quad (\text{A.10})$$

so that Eqs. (A.2), (A.3), and (A.4) can be rewritten in a more compact way

$$\hat{T} = -\frac{1}{2} \int [\nabla_{\mathbf{r}}^2 \hat{n}_1(\mathbf{r}, \mathbf{r}')]_{\mathbf{r}'=\mathbf{r}} d\mathbf{r}, \quad (\text{A.11})$$

$$\hat{W}_{ee} = \frac{1}{2} \iint w_{ee}(\mathbf{r}_1, \mathbf{r}_2) \hat{n}_2(\mathbf{r}_1, \mathbf{r}_2) d\mathbf{r}_1 d\mathbf{r}_2, \quad (\text{A.12})$$

$$\hat{V}_{ne} = \int v_{ne}(\mathbf{r}) \hat{n}(\mathbf{r}) d\mathbf{r}. \quad (\text{A.13})$$

We can also use the second-quantization formalism in an orthonormal spin-orbital basis $\{\psi_p(\mathbf{x})\}$ where $\mathbf{x} = (\mathbf{r}, \sigma)$. For this, we expand the field operators as

$$\hat{\psi}_{\sigma}^{\dagger}(\mathbf{r}) = \sum_p \psi_p^*(\mathbf{x}) \hat{a}_p^{\dagger}, \quad (\text{A.14})$$

and

$$\hat{\psi}_{\sigma}(\mathbf{r}) = \sum_p \psi_p(\mathbf{x}) \hat{a}_p, \quad (\text{A.15})$$

where \hat{a}_p^{\dagger} and \hat{a}_p are the creation and annihilation operators in this basis, which still obey anticommutation rules: $\{\hat{a}_p^{\dagger}, \hat{a}_q^{\dagger}\} = \{\hat{a}_p, \hat{a}_q\} = 0$ and $\{\hat{a}_p^{\dagger}, \hat{a}_q\} = \delta_{pq}$. The expressions of the operators are then

$$\hat{T} = \sum_{pq} t_{pq} \hat{a}_p^{\dagger} \hat{a}_q, \quad (\text{A.16})$$

$$\hat{W}_{ee} = \frac{1}{2} \sum_{pqrs} \langle \psi_p \psi_q | \psi_r \psi_s \rangle \hat{a}_p^{\dagger} \hat{a}_q^{\dagger} \hat{a}_s \hat{a}_r, \quad (\text{A.17})$$

$$\hat{V}_{ne} = \sum_{pq} v_{ne,pq} \hat{a}_p^{\dagger} \hat{a}_q, \quad (\text{A.18})$$

where t_{pq} and $v_{ne,pq}$ are the one-electron kinetic and nuclei-electron integrals, respectively, and $\langle \psi_p \psi_q | \psi_r \psi_s \rangle$ are the two-electron integrals.

B A brief introduction to functional derivatives

This section is inspired from Appendix A of Ref. [1] to which the reader is referred for more details.

A function f is a mapping from a number x to another number $f(x)$, i.e. $x \rightarrow f(x)$. Similarly, a *functional* F is a mapping from a function f to a number $F[f]$, i.e. $f \rightarrow F[f]$. In other words, a functional is a function of a function.

The differential of a functional $F[f]$ is $\delta F[f] = F[f + \delta f] - F[f]$, where δf is a infinitesimal variation of f . It represents the infinitesimal variation of $F[f]$ due to an infinitesimal variation of f . The infinitesimal variation $\delta F[f]$ is linear in $\delta f(x)$ at any point x

$$\delta F[f] = \int \frac{\delta F[f]}{\delta f(x)} \delta f(x) dx, \quad (\text{B.1})$$

which defines the *functional derivative* of $F[f]$ with respect $f(x)$, denoted by $\delta F[f]/\delta f(x)$. The functional derivative generalizes the concept of partial derivative. Indeed, if we consider a function $F(f_1, f_2, \dots)$ of several variables f_1, f_2, \dots , then the differential of F is $dF = \sum_i \partial F/\partial f_i df_i$, which is the analog of Eq. (B.1). Thus, $\delta F[f]/\delta f(x)$ is the analog of $\partial F/\partial f_i$ for the case of an infinitely continuous number of variables.

Functional derivatives shares most of the properties of ordinary derivatives. The functional derivative of a linear combination of functionals $c_1 F[f] + c_2 G[f]$ is

$$\frac{\delta}{\delta f(x)} (c_1 F[f] + c_2 G[f]) = c_1 \frac{\delta F[f]}{\delta f(x)} + c_2 \frac{\delta G[f]}{\delta f(x)}. \quad (\text{B.2})$$

The functional derivative of a product of two functionals $F[f]G[f]$ is

$$\frac{\delta}{\delta f(x)} (F[f]G[f]) = \frac{\delta F[f]}{\delta f(x)} G[f] + F[f] \frac{\delta G[f]}{\delta f(x)}. \quad (\text{B.3})$$

A functional $F[f]$ of a function $f[g](x)$ which is itself a functional of a function $g(x)$ has a functional derivative with respect to $g(x)$ given by the chain rule

$$\frac{\delta F[f]}{\delta g(x)} = \int \frac{\delta F[f]}{\delta f(x')} \frac{\delta f(x')}{\delta g(x)} dx'. \quad (\text{B.4})$$

It is the analog of the chain rule for a function $F(f_1, f_2, \dots)$ of several variables $f_i(g_1, g_2, \dots)$ which are themselves functions of other variables g_1, g_2, \dots , i.e. $\partial F/\partial g_i = \sum_j (\partial F/\partial f_j)(\partial f_j/\partial g_i)$.

An important special case is when the functional $F(f(x))$ is just an ordinary function of $f(x)$. The functional derivative of $F(f(x))$ with respect to $f(x')$ is

$$\frac{\delta F(f(x))}{\delta f(x')} = \frac{dF(f(x))}{df} \delta(x - x'), \quad (\text{B.5})$$

where dF/df is the ordinary derivative of the function F and $\delta(x - x')$ is the Dirac delta function. In particular, if $F(f(x)) = f(x)$, we have

$$\frac{\delta f(x)}{\delta f(x')} = \delta(x - x'). \quad (\text{B.6})$$

Also, if $f(x)$ is a functional of $g(x)$, the chain rule (B.4) and Eq. (B.6) give

$$\frac{\delta f(x)}{\delta f(x')} = \int \frac{\delta f(x)}{\delta g(x'')} \frac{\delta g(x'')}{\delta f(x')} dx'' = \delta(x - x'), \quad (\text{B.7})$$

which permits one to interpret $\delta g/\delta f$ as the inverse of $\delta f/\delta g$. Eq. (B.7) is analogous to the matrix relation: $\sum_k (A)_{ik} (A^{-1})_{kj} = \delta_{ij}$.

Higher-order functional derivatives can also be defined. For example, the second-order functional derivative $\delta^2 F[f]/\delta f(x)\delta f(x')$ is defined as the first-order functional derivative of $\delta F[f]/\delta f(x)$ with respect to $f(x')$. The order of differentiation is usually irrelevant

$$\frac{\delta^2 F[f]}{\delta f(x)\delta f(x')} = \frac{\delta^2 F[f]}{\delta f(x')\delta f(x)}. \quad (\text{B.8})$$

These functional derivatives can be used to expand a functional in a Taylor series

$$F[f + \Delta f] = F[f] + \int \frac{\delta F[f]}{\delta f(x)} \Delta f(x) dx + \frac{1}{2} \iint \frac{\delta^2 F[f]}{\delta f(x)\delta f(x')} \Delta f(x) \Delta f(x') dx dx' + \dots, \quad (\text{B.9})$$

where Δf is a finite change in f .

Finally, consider the following frequently occurring semilocal form for the functional $F[f]$ depending on a function $f(x)$ and its first-order derivative $f'(x) = df(x)/dx$

$$F[f] = \int h(f(x), f'(x)) dx, \quad (\text{B.10})$$

where h is some function and $f(x)$ vanishes at the boundary of x . The differential of $F[f]$ can be written as

$$\delta F[f] = \int \delta h(f(x), f'(x)) dx = \int \left[\frac{\partial h(f(x), f'(x))}{\partial f} \delta f(x) + \frac{\partial h(f(x), f'(x))}{\partial f'} \delta f'(x) \right] dx. \quad (\text{B.11})$$

Using now $\delta f'(x) = \delta[f(x + \varepsilon) - f(x)]/\varepsilon = [\delta f(x + \varepsilon) - \delta f(x)]/\varepsilon = (\delta f(x))'$ (with $\varepsilon \rightarrow 0$), and integrating by parts the second term in Eq. (B.11) gives

$$\delta F[f] = \int \left[\frac{\partial h(f(x), f'(x))}{\partial f} \delta f(x) - \frac{d}{dx} \left(\frac{\partial h(f(x), f'(x))}{\partial f'} \right) \delta f(x) \right] dx, \quad (\text{B.12})$$

where we have used that $\delta f(x)$ must vanish on the boundary so that the boundary term of the integration by parts vanishes. Comparing Eq. (B.12) with Eq. (B.1) shows that the functional derivative of $F[f]$ is

$$\frac{\delta F[f]}{\delta f(x)} = \frac{\partial h(f(x), f'(x))}{\partial f} - \frac{d}{dx} \left(\frac{\partial h(f(x), f'(x))}{\partial f'} \right). \quad (\text{B.13})$$

Exercise 24 : Generalize Eq. (B.13) to a semilocal functional F depending up to on n -order derivatives of f

$$F[f] = \int h(f(x), f'(x), f''(x), \dots, f^{(n)}(x)) dx, \quad (\text{B.14})$$

where $f^{(n)}(x) = d^n f(x)/dx^n$.



Solutions to the exercises

Exercise 1

For the special case of Coulombic potentials of the form $v_{\text{ne}}(\mathbf{r}) = -\sum_{\alpha} Z_{\alpha}/|\mathbf{r} - \mathbf{R}_{\alpha}|$, it is easy to see that the ground-state density $n_0(\mathbf{r})$ determines the parameters of the potential. Indeed, the locations of the local maxima of $n_0(\mathbf{r})$ give the positions of the nuclei $\mathbf{r}_{\text{max}} = \mathbf{R}_{\alpha}$, and the electron-nucleus cusp condition at each nucleus, $(\partial \bar{n}_0(r_{\alpha})/\partial r_{\alpha})_{r_{\alpha}=0} = -2Z_{\alpha} \bar{n}_0(0)$ where $\bar{n}_0(r_{\alpha})$ is the spherical average of the density around the nucleus α ($r_{\alpha} = |\mathbf{r} - \mathbf{R}_{\alpha}|$), gives the nucleus charge Z_{α} . Moreover, the integral of the density gives the number of electrons, $\int n_0(\mathbf{r}) d\mathbf{r} = N$. It is important to realize however that the Hohenberg-Kohn theorem is much more general than that since it states that the ground-state density determines the potential *for any form of potential* (that bounds N electrons).

Exercise 2

Let n_0 be a ground-state density associated with the potential $v_{\text{ne}}(\mathbf{r})$ and let $\Psi[n_0]$ be a corresponding ground-state wave function. Let n be a ground-state density associated with another potential $v(\mathbf{r})$ and let $\Psi[n]$ be a corresponding ground-state wave function. The variational property of the Hohenberg-Kohn density functional $E[n]$ directly follows from the variational theorem on wave functions:

$$\begin{aligned}
 E_0 = E[n_0] &= F[n_0] + \int v_{\text{ne}}(\mathbf{r}) n_0(\mathbf{r}) d\mathbf{r} \\
 &= \langle \Psi[n_0] | \hat{T} + \hat{W}_{\text{ee}} + \hat{V}_{\text{ne}} | \Psi[n_0] \rangle \\
 &\leq \langle \Psi[n] | \hat{T} + \hat{W}_{\text{ee}} + \hat{V}_{\text{ne}} | \Psi[n] \rangle \\
 &= F[n] + \int v_{\text{ne}}(\mathbf{r}) n(\mathbf{r}) d\mathbf{r} = E[n],
 \end{aligned} \tag{S.1}$$

establishing Eq. (1.16).

Exercise 3

Let us consider a variation of the density δn , the induced variation of the Hartree energy functional [Eq. (1.26)] is:

$$\begin{aligned}
 \delta E_{\text{H}}[n] &= \frac{1}{2} \iint \frac{\delta n(\mathbf{r}_1) n(\mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2 + \frac{1}{2} \iint \frac{n(\mathbf{r}_1) \delta n(\mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2 \\
 &= \frac{1}{2} \int \left[\int \frac{n(\mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_2 \right] \delta n(\mathbf{r}_1) d\mathbf{r}_1 + \frac{1}{2} \int \left[\int \frac{n(\mathbf{r}_1)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 \right] \delta n(\mathbf{r}_2) d\mathbf{r}_2 \\
 &= \int \left[\int \frac{n(\mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_2 \right] \delta n(\mathbf{r}_1) d\mathbf{r}_1,
 \end{aligned} \tag{S.2}$$

which, according to Eq. (B.1), allows us to identify the functional derivative of $E_{\text{H}}[n]$

$$v_{\text{H}}(\mathbf{r}_1) = \frac{\delta E_{\text{H}}[n]}{\delta n(\mathbf{r}_1)} = \int \frac{n(\mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_2. \tag{S.3}$$

Exercise 4

We start from the expression of the density in Eq. (1.31) and insert the expansion of $\varphi_i(\mathbf{r})$ in terms of basis functions [Eq. (1.43)]

$$\begin{aligned}
 n(\mathbf{r}) &= \sum_{i=1}^N |\varphi_i(\mathbf{r})|^2 = \sum_{i=1}^N \varphi_i(\mathbf{r}) \varphi_i^*(\mathbf{r}) \\
 &= \sum_{i=1}^N \left(\sum_{\gamma=1}^M c_{\gamma i} \chi_{\gamma}(\mathbf{r}) \right) \left(\sum_{\lambda=1}^M c_{\lambda i}^* \chi_{\lambda}^*(\mathbf{r}) \right) \\
 &= \sum_{\gamma=1}^M \sum_{\lambda=1}^M P_{\gamma\lambda} \chi_{\gamma}(\mathbf{r}) \chi_{\lambda}^*(\mathbf{r}),
 \end{aligned} \tag{S.4}$$

where $P_{\gamma\lambda} = \sum_{i=1}^N c_{\gamma i} c_{\lambda i}^*$. The second equality in Eq. (1.48) is then found by inserting the expression of the Hartree potential v_H [Eq. S.3] and using Eq. (S.4)

$$\begin{aligned}
J_{\mu\nu} &= \int \chi_{\mu}^*(\mathbf{r}_1) v_H(\mathbf{r}_1) \chi_{\nu}(\mathbf{r}_1) d\mathbf{r}_1 \\
&= \iint \chi_{\mu}^*(\mathbf{r}_1) \frac{n(\mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} \chi_{\nu}(\mathbf{r}_1) d\mathbf{r}_1 d\mathbf{r}_2 \\
&= \sum_{\gamma=1}^M \sum_{\lambda=1}^M P_{\gamma\lambda} \iint \frac{\chi_{\mu}^*(\mathbf{r}_1) \chi_{\gamma}(\mathbf{r}_2) \chi_{\lambda}^*(\mathbf{r}_2) \chi_{\nu}(\mathbf{r}_1)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2 \\
&= \sum_{\lambda=1}^M \sum_{\gamma=1}^M P_{\gamma\lambda} (\chi_{\mu} \chi_{\nu} | \chi_{\lambda} \chi_{\gamma}),
\end{aligned} \tag{S.5}$$

where $(\chi_{\mu} \chi_{\nu} | \chi_{\lambda} \chi_{\gamma})$ are the two-electron integrals defined in Eq. (1.50).

Exercise 5

Since the \uparrow - and \downarrow -spin electrons are uncoupled in the exchange energy (in the non-relativistic approximation), the spin-dependent exchange functional can be decomposed as

$$E_x[n_{\uparrow}, n_{\downarrow}] = E_x[n_{\uparrow}, 0] + E_x[0, n_{\downarrow}]. \tag{S.6}$$

Applying this equation with $n_{\uparrow} = n_{\downarrow} = n/2$ gives

$$E_x[n/2, n/2] = E_x[n/2, 0] + E_x[0, n/2] = 2E_x[n/2, 0] = 2E_x[0, n/2], \tag{S.7}$$

where it has been used that the functional is symmetric in its two arguments. Now, since $E_x[n/2, n/2] = E_x[n]$ where $E_x[n]$ is the spin-independent exchange functional, we thus have

$$2E_x[n/2, 0] = 2E_x[0, n/2] = E_x[n], \tag{S.8}$$

or, equivalently,

$$E_x[n, 0] = E_x[0, n] = \frac{1}{2} E_x[2n]. \tag{S.9}$$

Since this is true for any density n , it can be applied with arbitrary spin-resolved densities n_{\uparrow} and n_{\downarrow} , leading to the spin-scaling relation

$$E_x[n_{\uparrow}, n_{\downarrow}] = \frac{1}{2} (E_x[2n_{\uparrow}] + E_x[2n_{\downarrow}]). \tag{S.10}$$

Exercise 6

Using Eqs. (2.3) and (2.4), we can write the exchange-correlation hole as

$$n_{xc}(\mathbf{r}_1, \mathbf{r}_2) = \frac{n_2(\mathbf{r}_1, \mathbf{r}_2)}{n(\mathbf{r}_1)} - n(\mathbf{r}_2), \tag{S.11}$$

which, after integrating over \mathbf{r}_2 , leads to

$$\int n_{xc}(\mathbf{r}_1, \mathbf{r}_2) d\mathbf{r}_2 = \frac{\int n_2(\mathbf{r}_1, \mathbf{r}_2) d\mathbf{r}_2}{n(\mathbf{r}_1)} - \int n(\mathbf{r}_2) d\mathbf{r}_2 = \frac{(N-1)n(\mathbf{r}_1)}{n(\mathbf{r}_1)} - N = -1, \tag{S.12}$$

where the relation $\int n_2(\mathbf{r}_1, \mathbf{r}_2) d\mathbf{r}_2 = (N-1)n(\mathbf{r}_1)$ has been used, stemming directly from the definition of $n_2(\mathbf{r}_1, \mathbf{r}_2)$ in Eq. (2.1).

Exercise 7

We start from the definition of the exchange energy per particle of Eq. (2.15) and separate the domain of integration over \mathbf{r}_2 into two subdomains

$$\begin{aligned}\varepsilon_x[n](\mathbf{r}_1) &= \frac{1}{2} \int \frac{n_x(\mathbf{r}_1, \mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_2 \\ &= \frac{1}{2} \int_{\Omega(r_1)} \frac{n_x(\mathbf{r}_1, \mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_2 + \frac{1}{2} \int_{\bar{\Omega}(r_1)} \frac{n_x(\mathbf{r}_1, \mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_2,\end{aligned}\quad (\text{S.13})$$

where $\Omega(r_1)$ is the ball centered at $\mathbf{0}$ of radius r_1 and $\bar{\Omega}(r_1)$ is the complement subdomain. In the first subdomain $\Omega(r_1)$, we have $r_2 < r_1$ so we can make a convergent multipole expansion of the Coulomb interaction $1/|\mathbf{r}_1 - \mathbf{r}_2|$ around $\mathbf{r}_2 = \mathbf{0}$

$$\begin{aligned}\frac{1}{2} \int_{\Omega(r_1)} \frac{n_x(\mathbf{r}_1, \mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_2 &= \frac{1}{2} \int_{\Omega(r_1)} n_x(\mathbf{r}_1, \mathbf{r}_2) \left(\frac{1}{r_1} + \frac{\mathbf{r}_1 \cdot \mathbf{r}_2}{r_1^3} + \dots \right) d\mathbf{r}_2 \\ &= \frac{1}{2r_1} \int_{\Omega(r_1)} n_x(\mathbf{r}_1, \mathbf{r}_2) d\mathbf{r}_2 + \frac{\mathbf{r}_1}{2r_1^3} \cdot \int_{\Omega(r_1)} n_x(\mathbf{r}_1, \mathbf{r}_2) \mathbf{r}_2 d\mathbf{r}_2 + \dots \\ &\underset{r_1 \rightarrow +\infty}{\sim} \frac{1}{2r_1} \int n_x(\mathbf{r}_1, \mathbf{r}_2) d\mathbf{r}_2 = -\frac{1}{2r_1},\end{aligned}\quad (\text{S.14})$$

where we have used the sum rule on the exchange hole [Eq. (2.10)] and the fact that all the moments of the exchange hole exist for finite systems. As regards now the second contribution to the integral in Eq. (S.13), since for finite systems the exchange hole decays exponentially, $n_x(\mathbf{r}_1, \mathbf{r}_2) \underset{|\mathbf{r}_1 - \mathbf{r}_2| \rightarrow 0}{\propto} e^{-\alpha|\mathbf{r}_1 - \mathbf{r}_2|}$, this remaining contribution vanishes exponentially for $r_1 \rightarrow +\infty$

$$\frac{1}{2} \int_{\bar{\Omega}(r_1)} \frac{n_x(\mathbf{r}_1, \mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_2 \underset{r_1 \rightarrow +\infty}{=} O(e^{-\alpha r_1}). \quad (\text{S.15})$$

This thus proves that

$$\varepsilon_x[n](\mathbf{r}_1) \underset{r \rightarrow +\infty}{\sim} -\frac{1}{2r_1}. \quad (\text{S.16})$$

Exercise 8

We use the definition of the scaled wave function $\Psi_\gamma^\lambda[n]$ in Eq. (2.31) and perform the change of variables $\tilde{\mathbf{r}}_i = \gamma \mathbf{r}_i$ in the integrals to obtain

$$\begin{aligned}\langle \Psi_\gamma^\lambda[n] | \hat{T} + \lambda \gamma \hat{W}_{\text{ee}} | \Psi_\gamma^\lambda[n] \rangle &= \gamma^{3N} \int \dots \int \Psi^\lambda[n](\gamma \mathbf{r}_1, \sigma_1, \dots, \gamma \mathbf{r}_N, \sigma_N)^* \left[-\frac{1}{2} \sum_{i=1}^N \nabla_{\mathbf{r}_i}^2 + \sum_{1 \leq i < j \leq N} \frac{\lambda \gamma}{|\mathbf{r}_i - \mathbf{r}_j|} \right] \\ &\quad \times \Psi^\lambda[n](\gamma \mathbf{r}_1, \sigma_1, \dots, \gamma \mathbf{r}_N, \sigma_N) d\mathbf{r}_1 d\sigma_1 \dots d\mathbf{r}_N d\sigma_N \\ &= \int \dots \int \Psi^\lambda[n](\tilde{\mathbf{r}}_1, \sigma_1, \dots, \tilde{\mathbf{r}}_N, \sigma_N)^* \left[-\frac{\gamma^2}{2} \sum_{i=1}^N \nabla_{\tilde{\mathbf{r}}_i}^2 + \sum_{1 \leq i < j \leq N} \frac{\lambda \gamma^2}{|\tilde{\mathbf{r}}_i - \tilde{\mathbf{r}}_j|} \right] \\ &\quad \times \Psi^\lambda[n](\tilde{\mathbf{r}}_1, \sigma_1, \dots, \tilde{\mathbf{r}}_N, \sigma_N) d\tilde{\mathbf{r}}_1 d\sigma_1 \dots d\tilde{\mathbf{r}}_N d\sigma_N \\ &= \gamma^2 \langle \Psi^\lambda[n] | \hat{T} + \lambda \hat{W}_{\text{ee}} | \Psi^\lambda[n] \rangle,\end{aligned}\quad (\text{S.17})$$

which proves the identity of Eq. (2.33).

Exercise 9

The scaling relation satisfied by the Hartree density functional is directly found by using the definition

of the scaled density $n_\gamma(\mathbf{r}) = \gamma^3 n(\gamma\mathbf{r})$ and performing the change of variables $\tilde{\mathbf{r}} = \gamma\mathbf{r}$

$$\begin{aligned}
E_{\text{H}}[n_\gamma] &= \frac{1}{2} \iint \frac{n_\gamma(\mathbf{r}_1)n_\gamma(\mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2 \\
&= \frac{\gamma^6}{2} \iint \frac{n(\gamma\mathbf{r}_1)n(\gamma\mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2 \\
&= \frac{\gamma}{2} \iint \frac{n(\tilde{\mathbf{r}}_1)n(\tilde{\mathbf{r}}_2)}{|\tilde{\mathbf{r}}_1 - \tilde{\mathbf{r}}_2|} d\tilde{\mathbf{r}}_1 d\tilde{\mathbf{r}}_2 \\
&= \gamma E_{\text{H}}[n].
\end{aligned} \tag{S.18}$$

Using then the scaling of the KS wave function, $\Phi[n_\gamma] = \Phi_\gamma[n]$ [Eq. (2.38)] and the fact that $\langle \Phi_\gamma[n] | \hat{W}_{\text{ee}} | \Phi_\gamma[n] \rangle = \gamma \langle \Phi[n] | \hat{W}_{\text{ee}} | \Phi[n] \rangle$ [same proof as in Eq. (S.17)], we find the scaling relation satisfied by the exchange density functional

$$\begin{aligned}
E_{\text{x}}[n_\gamma] &= \langle \Phi[n_\gamma] | \hat{W}_{\text{ee}} | \Phi[n_\gamma] \rangle - E_{\text{H}}[n_\gamma] \\
&= \langle \Phi_\gamma[n] | \hat{W}_{\text{ee}} | \Phi_\gamma[n] \rangle - \gamma E_{\text{H}}[n] \\
&= \gamma \langle \Phi[n] | \hat{W}_{\text{ee}} | \Phi[n] \rangle - \gamma E_{\text{H}}[n] \\
&= \gamma E_{\text{x}}[n].
\end{aligned} \tag{S.19}$$

Exercise 10

We search the minimizing ensemble density matrices in the form

$$\hat{\Gamma} = (1-f) \sum_n w_n^{N-1} |\Psi_n^{N-1}\rangle \langle \Psi_n^{N-1}| + f \sum_m w_m^N |\Psi_m^N\rangle \langle \Psi_m^N|, \tag{S.20}$$

where $\{\Psi_n^{N-1}\}$ and $\{\Psi_m^N\}$ are complete orthonormal basis of fixed eigenstates of the $(N-1)$ - and N -electron systems, respectively, and w_n^{N-1} and w_m^N are weights to be optimized with the constraints $0 \leq w_n^{N-1} \leq 1$ and $\sum_n w_n^{N-1} = 1$, and similarly for w_m^N . The energy corresponding to the ensemble density matrix of Eq. (S.20) is

$$\begin{aligned}
E^{N-1+f} &= \text{Tr} [\hat{\Gamma} \hat{H}] \\
&= (1-f) \sum_n w_n^{N-1} \text{Tr} [|\Psi_n^{N-1}\rangle \langle \Psi_n^{N-1}| \hat{H}] + f \sum_m w_m^N \text{Tr} [|\Psi_m^N\rangle \langle \Psi_m^N| \hat{H}] \\
&= (1-f) \sum_{n,n'} w_n^{N-1} \langle \Psi_{n'}^{N-1} | \Psi_n^{N-1} \rangle \langle \Psi_n^{N-1} | \hat{H} | \Psi_{n'}^{N-1} \rangle + f \sum_{m,m'} w_m^N \langle \Psi_{m'}^N | \Psi_m^N \rangle \langle \Psi_m^N | \hat{H} | \Psi_{m'}^N \rangle \\
&= (1-f) \sum_n w_n^{N-1} \langle \Psi_n^{N-1} | \hat{H} | \Psi_n^{N-1} \rangle + f \sum_m w_m^N \langle \Psi_m^N | \hat{H} | \Psi_m^N \rangle \\
&= (1-f) \sum_n w_n^{N-1} E_n^{N-1} + f \sum_m w_m^N E_m^N,
\end{aligned} \tag{S.21}$$

where $E_n^{N-1} = \langle \Psi_n^{N-1} | \hat{H} | \Psi_n^{N-1} \rangle$ and $E_m^N = \langle \Psi_m^N | \hat{H} | \Psi_m^N \rangle$ are the corresponding eigenenergies for the $(N-1)$ - and N -electron systems, respectively. Clearly, the minimum of E^{N-1+f} in Eq. (S.21) is reached when only the ground-state energies E_0^{N-1} and E_0^N of the $(N-1)$ - and N -electron systems remain in the sums, i.e. for the weights $w_0^{N-1} = 1$ and $w_{n>0}^{N-1} = 0$, and $w_0^N = 1$ and $w_{m>0}^N = 0$. The minimizing ensemble density matrix is thus indeed given by Eq. (2.54) and the corresponding ground-state energy by Eq. (2.55).

Exercise 11

In the energy expression of Eq. (2.65), the orbitals $\{\varphi_j\}$ are optimized in the presence of the orbital occupation numbers $\{n_i\}$. Therefore, the derivative of the energy with respect to n_i contains two contributions, a term corresponding to the explicit dependence on n_i at fixed orbitals $\{\varphi_j\}$ and a term coming

from the implicit dependence on n_i via the orbitals $\{\varphi_j\}$,

$$\frac{\partial E}{\partial n_i} = \left(\frac{\partial E}{\partial n_i} \right)_{\{\varphi_j\}} + \left[\sum_{j=1}^N \int \frac{\delta E}{\delta \varphi_j^*(\mathbf{r})} \frac{\partial \varphi_j^*(\mathbf{r})}{\partial n_i} d\mathbf{r} + \text{c.c.} \right]. \quad (\text{S.22})$$

The first term gives

$$\begin{aligned} \left(\frac{\partial E}{\partial n_i} \right)_{\{\varphi_j\}} &= \int \varphi_i^*(\mathbf{r}) \left(-\frac{1}{2} \nabla^2 + v_{\text{ne}}(\mathbf{r}) \right) \varphi_i(\mathbf{r}) d\mathbf{r} + \left(\frac{\partial E_{\text{Hxc}}[n]}{\partial n_i} \right)_{\{\varphi_j\}} \\ &= \int \varphi_i^*(\mathbf{r}) \left(-\frac{1}{2} \nabla^2 + v_{\text{ne}}(\mathbf{r}) \right) \varphi_i(\mathbf{r}) d\mathbf{r} + \int \frac{\delta E_{\text{Hxc}}[n]}{\delta n(\mathbf{r})} \left(\frac{\partial n(\mathbf{r})}{\partial n_i} \right)_{\{\varphi_j\}} d\mathbf{r} \\ &= \int \varphi_i^*(\mathbf{r}) \left(-\frac{1}{2} \nabla^2 + v_{\text{ne}}(\mathbf{r}) + v_{\text{Hxc}}(\mathbf{r}) \right) \varphi_i(\mathbf{r}) d\mathbf{r} \\ &= \varepsilon_i, \end{aligned} \quad (\text{S.23})$$

where the expression of the density in terms of n_i in Eq. (2.66) has been used, and ε_i is the orbital energy introduced in Eq. (2.67). It can be shown that the second term in Eq. (S.22) vanishes

$$\begin{aligned} \sum_{j=1}^N \int \frac{\delta E}{\delta \varphi_j^*(\mathbf{r})} \frac{\partial \varphi_j^*(\mathbf{r})}{\partial n_i} d\mathbf{r} + \text{c.c.} &= \sum_{j=1}^N \int \left[n_j \left(-\frac{1}{2} \nabla^2 + v_{\text{ne}}(\mathbf{r}) \right) \varphi_j(\mathbf{r}) + \frac{\delta E_{\text{Hxc}}[n]}{\delta \varphi_j^*(\mathbf{r})} \right] \frac{\partial \varphi_j^*(\mathbf{r})}{\partial n_i} d\mathbf{r} + \text{c.c.} \\ &= \sum_{j=1}^N \int \left[n_j \left(-\frac{1}{2} \nabla^2 + v_{\text{ne}}(\mathbf{r}) + v_{\text{Hxc}}(\mathbf{r}) \right) \varphi_j(\mathbf{r}) \right] \frac{\partial \varphi_j^*(\mathbf{r})}{\partial n_i} d\mathbf{r} + \text{c.c.} \\ &= \sum_{j=1}^N \int n_j \varepsilon_j \varphi_j(\mathbf{r}) \frac{\partial \varphi_j^*(\mathbf{r})}{\partial n_i} d\mathbf{r} + \text{c.c.} \\ &= \sum_{j=1}^N n_j \varepsilon_j \frac{\partial}{\partial n_i} \int |\varphi_j(\mathbf{r})|^2 d\mathbf{r} = 0, \end{aligned} \quad (\text{S.24})$$

since the orbitals are normalized, $\int |\varphi_j(\mathbf{r})|^2 d\mathbf{r} = 1$. This proves Janak's theorem [Eq. (2.70)].

Exercise 12

For large r , the KS potential $v_s^N(\mathbf{r}) \underset{r \rightarrow +\infty}{\sim} v_s^N(\infty) + (Q-1)/r$ has spherical symmetry, and the KS equations (2.67) for the asymptotic orbitals $\varphi_{i,\text{asympt}}^N(r)$ can thus be written as

$$\begin{aligned} -\frac{1}{2} \left(\frac{d^2 \varphi_{i,\text{asympt}}^N(r)}{dr^2} + \frac{2}{r} \frac{d\varphi_{i,\text{asympt}}^N(r)}{dr} - \frac{\ell(\ell+1)}{r^2} \varphi_{i,\text{asympt}}^N(r) \right) + \left(v_s^N(\infty) + \frac{Q-1}{r} \right) \varphi_{i,\text{asympt}}^N(r) \\ = \varepsilon_i^N \varphi_{i,\text{asympt}}^N(r), \end{aligned} \quad (\text{S.25})$$

where ℓ is the angular momentum. For $r \rightarrow +\infty$, the terms in $1/r$ or $1/r^2$ are negligible, so it remains

$$-\frac{1}{2} \frac{d^2 \varphi_{i,\text{asympt}}^N(r)}{dr^2} + v_s^N(\infty) \varphi_{i,\text{asympt}}^N(r) = \varepsilon_i^N \varphi_{i,\text{asympt}}^N(r), \quad (\text{S.26})$$

which has the general solutions

$$\varphi_{i,\text{asympt}}^N(r) = c_1 e^{-\sqrt{-2(\varepsilon_i^N - v_s^N(\infty))} r} + c_2 e^{+\sqrt{-2(\varepsilon_i^N - v_s^N(\infty))} r}, \quad (\text{S.27})$$

for bound states, i.e. $\varepsilon_i^N < v_s^N(\infty)$. Since the second term on the right-hand-side of Eq. (S.27) diverges for $r \rightarrow +\infty$, we must have $c_2 = 0$ and therefore the asymptotic behavior of the KS orbitals is

$$\varphi_i^N(r) \underset{r \rightarrow +\infty}{\propto} e^{-\sqrt{-2(\varepsilon_i^N - v_s^N(\infty))} r}. \quad (\text{S.28})$$

Exercise 13

Let us impose the scaling relation $E_x[n_\gamma] = \gamma E_x[n]$ [Eq. (2.41)] to the functional $E_x^{\text{local}}[n] = c \int n(\mathbf{r})^p d\mathbf{r}$, we get

$$\begin{aligned}
E_x^{\text{local}}[n_\gamma] &= \gamma E_x^{\text{local}}[n] \\
\Leftrightarrow c \int n_\gamma(\mathbf{r})^p d\mathbf{r} &= \gamma c \int n(\mathbf{r})^p d\mathbf{r} \\
\Leftrightarrow c \gamma^{3p} \int n(\gamma \mathbf{r})^p d\mathbf{r} &= \gamma c \int n(\mathbf{r})^p d\mathbf{r} \\
\Leftrightarrow c \gamma^{3p-3} \int n(\mathbf{r}')^p d\mathbf{r}' &= \gamma c \int n(\mathbf{r})^p d\mathbf{r},
\end{aligned} \tag{S.29}$$

giving $3p - 3 = 1$ or $p = 4/3$. So, the simple uniform-coordinate scaling relation entirely determines the exponent of the density in the LDA exchange functional. Of course, it does not determine the prefactor c .

Exercise 14

By making the indicated replacements in Eq. (3.3), we arrive at

$$E_x^{\text{UEG}} = \frac{1}{(2\pi)^6} \int_{\Omega_{k_F}} d\mathbf{k}_1 \int_{\Omega_{k_F}} d\mathbf{k}_2 \iint \frac{1}{|\mathbf{r}_1 - \mathbf{r}_2|} e^{i(\mathbf{k}_1 - \mathbf{k}_2) \cdot (\mathbf{r}_2 - \mathbf{r}_1)} d\mathbf{r}_1 d\mathbf{r}_2, \tag{S.30}$$

where the spatial integrations are on a box of volume $V \rightarrow \infty$. Performing the change of variables $(\mathbf{r}_1, \mathbf{r}_2) \rightarrow (\mathbf{r}_1, \mathbf{r}_{12} = \mathbf{r}_2 - \mathbf{r}_1)$, we get

$$\begin{aligned}
E_x^{\text{UEG}} &= \frac{1}{(2\pi)^6} \int_{\Omega_{k_F}} d\mathbf{k}_1 \int_{\Omega_{k_F}} d\mathbf{k}_2 \iint \frac{1}{|\mathbf{r}_{12}|} e^{i(\mathbf{k}_1 - \mathbf{k}_2) \cdot \mathbf{r}_{12}} d\mathbf{r}_1 d\mathbf{r}_{12} \\
&= -\frac{V}{(2\pi)^6} \int_{\Omega_{k_F}} d\mathbf{k}_1 \int_{\Omega_{k_F}} d\mathbf{k}_2 \frac{4\pi}{|\mathbf{k}_1 - \mathbf{k}_2|^2},
\end{aligned} \tag{S.31}$$

where it was used that the integration over \mathbf{r}_{12} gives the Fourier transform of the Coulomb interaction $\int (1/|\mathbf{r}_{12}|) e^{i(\mathbf{k}_1 - \mathbf{k}_2) \cdot \mathbf{r}_{12}} d\mathbf{r}_{12} = 4\pi/|\mathbf{k}_1 - \mathbf{k}_2|^2$ and the integration over \mathbf{r}_1 gives the volume of the box $\int d\mathbf{r}_1 = V$. We can rewrite the integrals over \mathbf{k}_1 and \mathbf{k}_2 in spherical coordinates choosing as the polar angle θ of \mathbf{k}_2 the angle between \mathbf{k}_1 and \mathbf{k}_2

$$E_x^{\text{UEG}} = -\frac{V}{(2\pi)^6} \int_0^{k_F} dk_1 4\pi k_1^2 \int_0^{k_F} dk_2 2\pi k_2^2 \int_0^\pi d\theta \sin \theta \frac{4\pi}{k_1^2 + k_2^2 - 2k_1 k_2 \cos \theta}. \tag{S.32}$$

The integral over θ is easy to calculate

$$\begin{aligned}
\int_0^\pi d\theta \sin \theta \frac{4\pi}{k_1^2 + k_2^2 - 2k_1 k_2 \cos \theta} &= \int_{-1}^1 dx \frac{4\pi}{k_1^2 + k_2^2 - 2k_1 k_2 x} \\
&= \frac{-2\pi}{k_1 k_2} [\ln((k_1 - k_2)^2) - \ln((k_1 + k_2)^2)],
\end{aligned} \tag{S.33}$$

and we are left with

$$\begin{aligned}
E_x^{\text{UEG}} &= \frac{2V}{(2\pi)^3} \int_0^{k_F} dk_1 \int_0^{k_F} dk_2 k_1 k_2 [\ln((k_1 - k_2)^2) - \ln((k_1 + k_2)^2)] \\
&= \frac{2V k_F^4}{(2\pi)^3} \int_0^1 dx_1 \int_0^1 dx_2 x_1 x_2 [\ln((x_1 - x_2)^2) - \ln((x_1 + x_2)^2)],
\end{aligned} \tag{S.34}$$

where we have introduced $x_1 = k_1/k_F$ and $x_2 = k_2/k_F$. It can be shown that the last double integral over x_1 and x_2 equals -1 , so we get

$$E_x^{\text{UEG}} = -\frac{2V k_F^4}{(2\pi)^3}. \tag{S.35}$$

We finally obtain the exchange energy per particle

$$\varepsilon_x^{\text{UEG}} = \frac{E_x^{\text{UEG}}}{N} = -\frac{2k_F^4}{(2\pi)^3 n} = C_x n^{1/3}, \quad (\text{S.36})$$

where the definition of $k_F = (3\pi^2 n)^{1/3}$ was used and the constant $C_x = -(3/4)(3/\pi)^{1/3}$ was introduced. For an alternative way of calculating the integral in Eq. (S.31) see Fetter and Walecka, page 28 [134].

Exercise 15

The LDA exchange energy functional is

$$E_x^{\text{LDA}}[n] = \int n(\mathbf{r}) \varepsilon_x^{\text{UEG}}(n(\mathbf{r})) d\mathbf{r} = \int f(n(\mathbf{r})) d\mathbf{r}, \quad (\text{S.37})$$

where $f(n) = C_x n^{4/3}$. The variation of $E_x^{\text{LDA}}[n]$ due to a variation of the density can be written as

$$\delta E_x^{\text{LDA}}[n] = \int \delta f(n(\mathbf{r})) d\mathbf{r} = \int \left(\frac{df(n)}{dn} \right)_{n=n(\mathbf{r})} \delta n(\mathbf{r}) d\mathbf{r}, \quad (\text{S.38})$$

which, according to Eq. (B.1), means that the functional derivative is

$$v_x^{\text{LDA}}(\mathbf{r}) = \frac{\delta E_x^{\text{LDA}}[n]}{\delta n(\mathbf{r})} = \left(\frac{df(n)}{dn} \right)_{n=n(\mathbf{r})} = \frac{4}{3} C_x n(\mathbf{r})^{1/3}. \quad (\text{S.39})$$

Since the density $n(\mathbf{r})$ decays exponentially for $r \rightarrow +\infty$ [Eq. (2.73)], the LDA exchange potential $v_x^{\text{LDA}}(\mathbf{r})$ also decays exponentially for $r \rightarrow +\infty$, i.e. much too fast in comparison to the asymptotic behavior of the exact exchange potential $v_x(\mathbf{r}) \underset{r \rightarrow +\infty}{\sim} -1/r$.

Exercise 16

The Wigner correlation energy functional can be written as

$$E_c^{\text{W}}[n] = \int f(n(\mathbf{r})) d\mathbf{r}, \quad (\text{S.40})$$

with $f(n) = cn/(d+kn^{-1/3})$ where $k = (3/(4\pi))^{1/3}$. Similarly to Exercise 15, the corresponding potential is calculated as

$$\begin{aligned} v_c^{\text{W}}(\mathbf{r}) &= \frac{\delta E_c^{\text{W}}[n]}{\delta n(\mathbf{r})} = \left(\frac{df(n)}{dn} \right)_{n=n(\mathbf{r})} = \frac{c(d+kn(\mathbf{r})^{-1/3}) - cn(\mathbf{r})(-kn(\mathbf{r})^{-4/3}/3)}{(d+kn(\mathbf{r})^{-1/3})^2} \\ &= \frac{cd + (4/3)ckn(\mathbf{r})^{-1/3}}{(d+kn(\mathbf{r})^{-1/3})^2}. \end{aligned} \quad (\text{S.41})$$

Exercise 17

The derivation of the potential of a GGA functional is just a 3D generalization of the 1D case of Eqs.(B.10)-(B.13). The variation of $E_{xc}^{\text{GGA}}[n]$ induced by a variation of the density $\delta n(\mathbf{r})$ is

$$\begin{aligned} \delta E_{xc}^{\text{GGA}}[n] &= \int \delta e_{xc}^{\text{GGA}}(n(\mathbf{r}), \nabla n(\mathbf{r})) d\mathbf{r} \\ &= \int \left[\frac{\partial e_{xc}^{\text{GGA}}(n(\mathbf{r}), \nabla n(\mathbf{r}))}{\partial n} \delta n(\mathbf{r}) + \frac{\partial e_{xc}^{\text{GGA}}(n(\mathbf{r}), \nabla n(\mathbf{r}))}{\partial \nabla n} \cdot \delta \nabla n(\mathbf{r}) \right] d\mathbf{r}, \end{aligned} \quad (\text{S.42})$$

where the notation $\partial e_{xc}^{\text{GGA}}/\partial \nabla n$ means the vector containing the derivatives of e_{xc}^{GGA} with respect to the different components of ∇n . Using $\delta \nabla n(\mathbf{r}) = \nabla \delta n(\mathbf{r})$ (since ∇ is a linear operator) and integrating by parts the second term in Eq. (S.42)

$$\delta E_{xc}^{\text{GGA}}[n] = \int \left[\frac{\partial e_{xc}^{\text{GGA}}(n(\mathbf{r}), \nabla n(\mathbf{r}))}{\partial n} \delta n(\mathbf{r}) - \nabla \cdot \frac{\partial e_{xc}^{\text{GGA}}(n(\mathbf{r}), \nabla n(\mathbf{r}))}{\partial \nabla n} \delta n(\mathbf{r}) \right] d\mathbf{r}, \quad (\text{S.43})$$

where we have used that $\delta n(\mathbf{r})$ must vanish at infinity so that the boundary term of the integration by parts vanishes. Using the definition of a functional derivative, the potential associated with $E_{xc}^{\text{GGA}}[n]$ is thus

$$v_{xc}^{\text{GGA}}(\mathbf{r}) = \frac{\delta E_{xc}^{\text{GGA}}[n]}{\delta n(\mathbf{r})} = \frac{\partial e_{xc}^{\text{GGA}}}{\partial n}(n(\mathbf{r}), \nabla n(\mathbf{r})) - \nabla \cdot \frac{\partial e_{xc}^{\text{GGA}}}{\partial \nabla n}(n(\mathbf{r}), \nabla n(\mathbf{r})). \quad (\text{S.44})$$

The contribution of $v_{xc}^{\text{GGA}}(\mathbf{r})$ to the KS Fock matrix [Eq. (1.51)] can then be expressed as

$$\begin{aligned} V_{xc,\mu\nu}^{\text{GGA}} &= \int \chi_{\mu}^*(\mathbf{r}) v_{xc}^{\text{GGA}}(\mathbf{r}) \chi_{\nu}(\mathbf{r}) d\mathbf{r} \\ &= \int \chi_{\mu}^*(\mathbf{r}) \left[\frac{\partial e_{xc}^{\text{GGA}}}{\partial n}(n(\mathbf{r}), \nabla n(\mathbf{r})) - \nabla \cdot \frac{\partial e_{xc}^{\text{GGA}}}{\partial \nabla n}(n(\mathbf{r}), \nabla n(\mathbf{r})) \right] \chi_{\nu}(\mathbf{r}) d\mathbf{r} \\ &= \int \frac{\partial e_{xc}^{\text{GGA}}}{\partial n}(n(\mathbf{r}), \nabla n(\mathbf{r})) \chi_{\mu}^*(\mathbf{r}) \chi_{\nu}(\mathbf{r}) d\mathbf{r} + \int \frac{\partial e_{xc}^{\text{GGA}}}{\partial \nabla n}(n(\mathbf{r}), \nabla n(\mathbf{r})) \cdot \nabla (\chi_{\mu}^*(\mathbf{r}) \chi_{\nu}(\mathbf{r})) d\mathbf{r}, \end{aligned} \quad (\text{S.45})$$

where an integration by parts has been performed in the second term. Note that, since in practice e_{xc}^{GGA} depends on $(\nabla n)^2$, the derivative $\partial e_{xc}^{\text{GGA}}/\partial \nabla n$ is calculated as

$$\frac{\partial e_{xc}^{\text{GGA}}}{\partial \nabla n} = \frac{\partial e_{xc}^{\text{GGA}}}{\partial (\nabla n)^2} \frac{d(\nabla n)^2}{d\nabla n} = 2 \frac{\partial e_{xc}^{\text{GGA}}}{\partial (\nabla n)^2} \nabla n. \quad (\text{S.46})$$

Exercise 18

For a one-electron system, there is only one occupied orbital, which can be explicitly written in terms of the density, $\varphi(\mathbf{r}) = \sqrt{n(\mathbf{r})}$, up to a unimportant phase factor. It is then immediate to check that the kinetic energy density of such a system is the von Weizsäcker kinetic energy density:

$$\tau_{1e}(\mathbf{r}) = \frac{1}{2} |\nabla \varphi(\mathbf{r})|^2 = \frac{1}{2} \left(\nabla \sqrt{n(\mathbf{r})} \right)^2 = \frac{1}{2} \left(\frac{\nabla n(\mathbf{r})}{2\sqrt{n(\mathbf{r})}} \right)^2 = \frac{|\nabla n(\mathbf{r})|^2}{8n(\mathbf{r})} = \tau^{\text{W}}(\mathbf{r}). \quad (\text{S.47})$$

Similarly, for two electrons in a single spatial occupied, we have $\varphi(\mathbf{r}) = \sqrt{n(\mathbf{r})/2}$, and it is easy to show that the *non-interacting* kinetic energy density is again the von Weizsäcker kinetic energy density.

Exercise 19

The variation of the HF exchange energy [Eq. (3.18)] due to a variation of $\varphi_{i\sigma}^*(\mathbf{r})$ is

$$\begin{aligned} \delta E_x^{\text{HF}} &= -\frac{1}{2} \sum_{j=1}^{N_{\sigma}} \iint \frac{\delta \varphi_{i\sigma}^*(\mathbf{r}_1) \varphi_{j\sigma}(\mathbf{r}_1) \varphi_{j\sigma}^*(\mathbf{r}_2) \varphi_{i\sigma}(\mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2 \\ &\quad -\frac{1}{2} \sum_{j=1}^{N_{\sigma}} \iint \frac{\varphi_{j\sigma}^*(\mathbf{r}_1) \varphi_{i\sigma}(\mathbf{r}_1) \delta \varphi_{i\sigma}^*(\mathbf{r}_2) \varphi_{j\sigma}(\mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2 \\ &= -\sum_{j=1}^{N_{\sigma}} \iint \frac{\delta \varphi_{i\sigma}^*(\mathbf{r}_1) \varphi_{j\sigma}(\mathbf{r}_1) \varphi_{j\sigma}^*(\mathbf{r}_2) \varphi_{i\sigma}(\mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2, \end{aligned} \quad (\text{S.48})$$

and the functional derivative is thus

$$\frac{\delta E_x^{\text{HF}}}{\delta \varphi_{i\sigma}^*(\mathbf{r}_1)} = -\sum_{j=1}^{N_{\sigma}} \int \frac{\varphi_{j\sigma}(\mathbf{r}_1) \varphi_{j\sigma}^*(\mathbf{r}_2) \varphi_{i\sigma}(\mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_2 = \int v_{x,\sigma}^{\text{HF}}(\mathbf{r}_1, \mathbf{r}_2) \varphi_{i\sigma}(\mathbf{r}_2) d\mathbf{r}_2, \quad (\text{S.49})$$

where $v_{x,\sigma}^{\text{HF}}(\mathbf{r}_1, \mathbf{r}_2)$ is the nonlocal HF exchange potential

$$v_{x,\sigma}^{\text{HF}}(\mathbf{r}_1, \mathbf{r}_2) = -\sum_{j=1}^{N_{\sigma}} \frac{\varphi_{j\sigma}(\mathbf{r}_1) \varphi_{j\sigma}^*(\mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|}. \quad (\text{S.50})$$

Alternatively, the functional derivative of Eq. (S.49) can be reformulated as $\delta E_x^{\text{HF}} / \delta \varphi_{i\sigma}^*(\mathbf{r}_1) = v_{x,i\sigma}^{\text{HF}}(\mathbf{r}_1) \varphi_{i\sigma}(\mathbf{r}_1)$ where $v_{x,i\sigma}^{\text{HF}}(\mathbf{r})$ is a local HF exchange potential depending on each orbital it acts on

$$v_{x,i\sigma}^{\text{HF}}(\mathbf{r}_1) = \frac{1}{\varphi_{i\sigma}(\mathbf{r}_1)} \int v_{x,\sigma}^{\text{HF}}(\mathbf{r}_1, \mathbf{r}_2) \varphi_{i\sigma}(\mathbf{r}_2) d\mathbf{r}_2. \quad (\text{S.51})$$

Exercise 20

The error function is written as $\text{erf}(x) = (2/\sqrt{\pi})F(x)$ with $F(x) = \int_0^x e^{-t^2} dt$. Its Taylor expansion for $x \rightarrow 0$ is

$$\text{erf}(x) = \frac{2}{\sqrt{\pi}} (F(0) + F'(0)x + \dots). \quad (\text{S.52})$$

We have $F(0) = 0$, and since $F'(x) = e^{-x^2}$, we have $F'(0) = 1$, leading to $\text{erf}(x) \underset{x \rightarrow 0}{\sim} (2/\sqrt{\pi})x$. The limit $x \rightarrow +\infty$ is easily obtained by using the Gaussian integral $\int_0^{+\infty} e^{-t^2} dt = \sqrt{\pi}/2$, leading to $\text{erf}(x) \xrightarrow{x \rightarrow +\infty} 1$.

Exercise 21

According to standard first-order perturbation theory, if a perturbation $v_s(\mathbf{r}) \rightarrow v_s(\mathbf{r}) + \delta v_s(\mathbf{r})$ is applied on the KS system, the first-order variation of the orbital $\varphi_{i\sigma}(\mathbf{r}')$ (assumed to be nondegenerate) is

$$\delta \varphi_{i\sigma}(\mathbf{r}') = - \sum_{\substack{p \geq 1 \\ p \neq i}} \frac{\int \varphi_{p\sigma}^*(\mathbf{r}) \delta v_s(\mathbf{r}) \varphi_{i\sigma}(\mathbf{r}) d\mathbf{r}}{\varepsilon_{p\sigma} - \varepsilon_{i\sigma}} \varphi_{p\sigma}(\mathbf{r}'), \quad (\text{S.53})$$

where the sum is over all spatial orbitals p different from orbital i but of the same spin. The functional derivative of $\varphi_{i\sigma}(\mathbf{r}')$ with respect to $v_s(\mathbf{r})$ is thus

$$\frac{\delta \varphi_{i\sigma}(\mathbf{r}')}{\delta v_s(\mathbf{r})} = - \sum_{\substack{p \geq 1 \\ p \neq i}} \frac{\varphi_{p\sigma}^*(\mathbf{r}) \varphi_{i\sigma}(\mathbf{r})}{\varepsilon_{p\sigma} - \varepsilon_{i\sigma}} \varphi_{p\sigma}(\mathbf{r}'). \quad (\text{S.54})$$

This result can be used to calculate the expression of $\chi_0(\mathbf{r}', \mathbf{r})$

$$\begin{aligned} \chi_0(\mathbf{r}', \mathbf{r}) = \frac{\delta n(\mathbf{r}')}{\delta v_s(\mathbf{r})} &= \sum_{\sigma \in \{\uparrow, \downarrow\}} \sum_{i=1}^{N_\sigma} \left[\varphi_{i\sigma}^*(\mathbf{r}') \frac{\delta \varphi_{i\sigma}(\mathbf{r}')}{\delta v_s(\mathbf{r})} + \text{c.c.} \right] \\ &= - \sum_{\sigma \in \{\uparrow, \downarrow\}} \sum_{i=1}^{N_\sigma} \sum_{\substack{p \geq 1 \\ p \neq i}} \left[\varphi_{i\sigma}^*(\mathbf{r}') \frac{\varphi_{p\sigma}^*(\mathbf{r}) \varphi_{i\sigma}(\mathbf{r})}{\varepsilon_{p\sigma} - \varepsilon_{i\sigma}} \varphi_{p\sigma}(\mathbf{r}') + \text{c.c.} \right]. \end{aligned} \quad (\text{S.55})$$

The last sum can be simplified by decomposing it as $\sum_{i=1}^{N_\sigma} \sum_{p \geq 1, p \neq i} = \sum_{i=1}^{N_\sigma} \sum_{p=1, p \neq i}^{N_\sigma} + \sum_{i=1}^{N_\sigma} \sum_{a \geq N_\sigma + 1}$ and realizing that the double sum $\sum_{i=1}^{N_\sigma} \sum_{p=1, p \neq i}^{N_\sigma}$ is zero because the summand inside the square bracket is antisymmetric with respect to the exchange of i and p . We thus arrive at the expected expression

$$\chi_0(\mathbf{r}', \mathbf{r}) = - \sum_{\sigma \in \{\uparrow, \downarrow\}} \sum_{i=1}^{N_\sigma} \sum_{a \geq N_\sigma + 1} \frac{\varphi_{i\sigma}^*(\mathbf{r}') \varphi_{a\sigma}^*(\mathbf{r}) \varphi_{i\sigma}(\mathbf{r}) \varphi_{a\sigma}(\mathbf{r}')}{\varepsilon_{a\sigma} - \varepsilon_{i\sigma}} + \text{c.c.} \quad (\text{S.56})$$

As regards the functional derivative of E_x with respect to $v_s(\mathbf{r})$, it is obtained by using chain rule with

the orbitals and using Eqs. (S.49), (S.50), and (S.54)

$$\begin{aligned}
\frac{\delta E_x}{\delta v_s(\mathbf{r})} &= \sum_{\sigma \in \{\uparrow, \downarrow\}} \sum_{i=1}^{N_\sigma} \int \frac{\delta E_x}{\delta \varphi_{i\sigma}^*(\mathbf{r}_1)} \frac{\delta \varphi_{i\sigma}^*(\mathbf{r}_1)}{\delta v_s(\mathbf{r})} d\mathbf{r}_1 + \text{c.c.} \\
&= - \sum_{\sigma \in \{\uparrow, \downarrow\}} \sum_{i=1}^{N_\sigma} \sum_{\substack{p \geq 1 \\ p \neq i}} \iint v_{x,\sigma}^{\text{HF}}(\mathbf{r}_1, \mathbf{r}_2) \varphi_{i\sigma}(\mathbf{r}_2) \frac{\varphi_{p\sigma}(\mathbf{r}) \varphi_{i\sigma}^*(\mathbf{r})}{\varepsilon_{p\sigma} - \varepsilon_{i\sigma}} \varphi_{p\sigma}^*(\mathbf{r}_1) d\mathbf{r}_1 d\mathbf{r}_2 + \text{c.c.} \\
&= \sum_{\sigma \in \{\uparrow, \downarrow\}} \sum_{i=1}^{N_\sigma} \sum_{j=1}^{N_\sigma} \sum_{\substack{p \geq 1 \\ p \neq i}} \left[(\varphi_{p\sigma} \varphi_{j\sigma} | \varphi_{j\sigma} \varphi_{i\sigma}) \frac{\varphi_{p\sigma}(\mathbf{r}) \varphi_{i\sigma}^*(\mathbf{r})}{\varepsilon_{p\sigma} - \varepsilon_{i\sigma}} + \text{c.c.} \right] \\
&= \sum_{\sigma \in \{\uparrow, \downarrow\}} \sum_{i=1}^{N_\sigma} \sum_{j=1}^{N_\sigma} \sum_{a \geq N_\sigma + 1} (\varphi_{a\sigma} \varphi_{j\sigma} | \varphi_{j\sigma} \varphi_{i\sigma}) \frac{\varphi_{a\sigma}(\mathbf{r}) \varphi_{i\sigma}^*(\mathbf{r})}{\varepsilon_{a\sigma} - \varepsilon_{i\sigma}} + \text{c.c.} , \tag{S.57}
\end{aligned}$$

where again it was used that $\sum_{i=1}^{N_\sigma} \sum_{p \geq 1, p \neq i} = \sum_{i=1}^{N_\sigma} \sum_{p=1, p \neq i}^{N_\sigma} + \sum_{i=1}^{N_\sigma} \sum_{a \geq N_\sigma + 1} = \sum_{i=1}^{N_\sigma} \sum_{a \geq N_\sigma + 1}$ since the summand inside the square bracket is antisymmetric with respect to the exchange of i and p .

Exercise 22

In the GL2 correlation energy expression of Eq. (4.11), only double and single excitations contribute

$$E_c^{\text{GL2}} = - \sum_{i=1}^N \sum_{j=i+1}^N \sum_{a \geq N+1} \sum_{b \geq a+1} \frac{|\langle \Phi | \hat{W}_{ee} - \hat{V}_{\text{Hx}} | \Phi_{ij}^{ab} \rangle|^2}{\varepsilon_a + \varepsilon_b - \varepsilon_i - \varepsilon_j} - \sum_{i=1}^N \sum_{a \geq N+1} \frac{|\langle \Phi | \hat{W}_{ee} - \hat{V}_{\text{Hx}} | \Phi_i^a \rangle|^2}{\varepsilon_a - \varepsilon_i}. \tag{S.58}$$

Applying the standard Slater's rules, $\langle \Phi | \hat{W}_{ee} - \hat{V}_{\text{Hx}} | \Phi_{ij}^{ab} \rangle = \langle \Phi | \hat{W}_{ee} | \Phi_{ij}^{ab} \rangle = \langle ij || ab \rangle$, the first term gives the MP2-like contribution

$$E_c^{\text{MP2}} = - \sum_{i=1}^N \sum_{j=i+1}^N \sum_{a \geq N+1} \sum_{b \geq a+1} \frac{|\langle \psi_i \psi_j | \psi_a \psi_b \rangle|^2}{\varepsilon_a + \varepsilon_b - \varepsilon_i - \varepsilon_j} = - \frac{1}{4} \sum_{i=1}^N \sum_{j=1}^N \sum_{a \geq N+1} \sum_{b \geq N+1} \frac{|\langle \psi_i \psi_j | \psi_a \psi_b \rangle|^2}{\varepsilon_a + \varepsilon_b - \varepsilon_i - \varepsilon_j}. \tag{S.59}$$

As regards the second term, using $\langle \Phi | \hat{W}_{ee} - \hat{V}_{\text{Hx}} | \Phi_i^a \rangle = \sum_{j=1}^N \langle \psi_i \psi_j | \psi_a \psi_j \rangle - \langle \psi_i | \hat{V}_{\text{Hx}} | \psi_a \rangle = \langle \psi_i | \hat{V}_{\text{Hx}}^{\text{HF}} - \hat{V}_{\text{Hx}} | \psi_a \rangle = \langle \psi_i | \hat{V}_x^{\text{HF}} - \hat{V}_x | \psi_a \rangle$, we find the expected expression

$$E_c^{\text{S}} = - \sum_{i=1}^N \sum_{a \geq N+1} \frac{|\langle \psi_i | \hat{V}_x^{\text{HF}} - \hat{V}_x | \psi_a \rangle|^2}{\varepsilon_a - \varepsilon_i}. \tag{S.60}$$

Exercise 23

The linear-response function of Eq. (4.19) can be written as

$$\begin{aligned}
i\chi_\lambda(\mathbf{r}_1 t_1, \mathbf{r}_2 t_2) &= \theta(t_1 - t_2) \langle \Psi^\lambda | e^{i\hat{H}^\lambda t_1} \hat{n}(\mathbf{r}_1) e^{-i\hat{H}^\lambda t_1} e^{i\hat{H}^\lambda t_2} \hat{n}(\mathbf{r}_2) e^{-i\hat{H}^\lambda t_2} | \Psi^\lambda \rangle \\
&\quad + \theta(t_2 - t_1) \langle \Psi^\lambda | e^{i\hat{H}^\lambda t_2} \hat{n}(\mathbf{r}_2) e^{-i\hat{H}^\lambda t_2} e^{i\hat{H}^\lambda t_1} \hat{n}(\mathbf{r}_1) e^{-i\hat{H}^\lambda t_1} | \Psi^\lambda \rangle \\
&\quad - \langle \Psi^\lambda | \hat{n}(\mathbf{r}_1) | \Psi^\lambda \rangle \langle \Psi^\lambda | \hat{n}(\mathbf{r}_2) | \Psi^\lambda \rangle, \tag{S.61}
\end{aligned}$$

or, after introducing a complete set of orthonormal eigenstates of the Hamiltonian, $\hat{H}^\lambda | \Psi_n^\lambda \rangle = E_n^\lambda | \Psi_n^\lambda \rangle$ (with $\Psi_0^\lambda = \Psi^\lambda$),

$$\begin{aligned}
i\chi_\lambda(\mathbf{r}_1 t_1, \mathbf{r}_2 t_2) &= \theta(t_1 - t_2) \sum_n \langle \Psi^\lambda | e^{i\hat{H}^\lambda t_1} \hat{n}(\mathbf{r}_1) e^{-i\hat{H}^\lambda t_1} | \Psi_n^\lambda \rangle \langle \Psi_n^\lambda | e^{i\hat{H}^\lambda t_2} \hat{n}(\mathbf{r}_2) e^{-i\hat{H}^\lambda t_2} | \Psi^\lambda \rangle \\
&\quad + \theta(t_2 - t_1) \sum_n \langle \Psi^\lambda | e^{i\hat{H}^\lambda t_2} \hat{n}(\mathbf{r}_2) e^{-i\hat{H}^\lambda t_2} | \Psi_n^\lambda \rangle \langle \Psi_n^\lambda | e^{i\hat{H}^\lambda t_1} \hat{n}(\mathbf{r}_1) e^{-i\hat{H}^\lambda t_1} | \Psi^\lambda \rangle \\
&\quad - \langle \Psi^\lambda | \hat{n}(\mathbf{r}_1) | \Psi^\lambda \rangle \langle \Psi^\lambda | \hat{n}(\mathbf{r}_2) | \Psi^\lambda \rangle. \tag{S.62}
\end{aligned}$$

After applying the Hamiltonian \hat{H}^λ on its eigenstates, it is apparent that $\chi_\lambda(\mathbf{r}_1 t_1, \mathbf{r}_2 t_2)$ only depends on $\tau = t_1 - t_2$

$$\begin{aligned} i\chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \tau) &= \theta(\tau) \sum_n \langle \Psi^\lambda | \hat{n}(\mathbf{r}_1) | \Psi_n^\lambda \rangle \langle \Psi_n^\lambda | \hat{n}(\mathbf{r}_2) | \Psi^\lambda \rangle e^{-i\omega_n^\lambda \tau} \\ &\quad + \theta(-\tau) \sum_n \langle \Psi^\lambda | \hat{n}(\mathbf{r}_2) | \Psi_n^\lambda \rangle \langle \Psi_n^\lambda | \hat{n}(\mathbf{r}_1) | \Psi^\lambda \rangle e^{i\omega_n^\lambda \tau} \\ &\quad - \langle \Psi^\lambda | \hat{n}(\mathbf{r}_1) | \Psi^\lambda \rangle \langle \Psi^\lambda | \hat{n}(\mathbf{r}_2) | \Psi^\lambda \rangle, \end{aligned} \quad (\text{S.63})$$

where $\omega_n^\lambda = E_n^\lambda - E_0^\lambda$ are the excitation energies. Since the last term in Eq. (S.63) just corresponds to the $n = 0$ contribution from the sums, we finally find

$$\begin{aligned} i\chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \tau) &= \theta(\tau) \sum_{n \neq 0} \langle \Psi^\lambda | \hat{n}(\mathbf{r}_1) | \Psi_n^\lambda \rangle \langle \Psi_n^\lambda | \hat{n}(\mathbf{r}_2) | \Psi^\lambda \rangle e^{-i\omega_n^\lambda \tau} \\ &\quad + \theta(-\tau) \sum_{n \neq 0} \langle \Psi^\lambda | \hat{n}(\mathbf{r}_2) | \Psi_n^\lambda \rangle \langle \Psi_n^\lambda | \hat{n}(\mathbf{r}_1) | \Psi^\lambda \rangle e^{i\omega_n^\lambda \tau}. \end{aligned} \quad (\text{S.64})$$

Using the Fourier-transform formulas for the Heaviside step function, $\theta(\tau) = -1/(2\pi i) \int_{-\infty}^{\infty} d\omega e^{-i\omega\tau}/(\omega + i0^+)$ and $\theta(-\tau) = 1/(2\pi i) \int_{-\infty}^{\infty} d\omega e^{-i\omega\tau}/(\omega - i0^+)$, we can express $i\chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \tau)$ as

$$\begin{aligned} i\chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \tau) &= - \int_{-\infty}^{\infty} \frac{d\omega}{2\pi i} \sum_{n \neq 0} \frac{\langle \Psi^\lambda | \hat{n}(\mathbf{r}_1) | \Psi_n^\lambda \rangle \langle \Psi_n^\lambda | \hat{n}(\mathbf{r}_2) | \Psi^\lambda \rangle}{\omega + i0^+} e^{-i(\omega + \omega_n^\lambda)\tau} \\ &\quad + \int_{-\infty}^{\infty} \frac{d\omega}{2\pi i} \sum_{n \neq 0} \frac{\langle \Psi^\lambda | \hat{n}(\mathbf{r}_2) | \Psi_n^\lambda \rangle \langle \Psi_n^\lambda | \hat{n}(\mathbf{r}_1) | \Psi^\lambda \rangle}{\omega - i0^-} e^{-i(\omega - \omega_n^\lambda)\tau}, \end{aligned} \quad (\text{S.65})$$

which, after making the substitutions $\omega \rightarrow \omega - \omega_n^\lambda$ and $\omega \rightarrow \omega + \omega_n^\lambda$ in the first and second integrals, respectively, can be recast in the form

$$\chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \tau) = \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} e^{-i\omega\tau} \sum_{n \neq 0} \frac{\langle \Psi^\lambda | \hat{n}(\mathbf{r}_1) | \Psi_n^\lambda \rangle \langle \Psi_n^\lambda | \hat{n}(\mathbf{r}_2) | \Psi^\lambda \rangle}{\omega - \omega_n^\lambda + i0^+} - \frac{\langle \Psi^\lambda | \hat{n}(\mathbf{r}_2) | \Psi_n^\lambda \rangle \langle \Psi_n^\lambda | \hat{n}(\mathbf{r}_1) | \Psi^\lambda \rangle}{\omega + \omega_n^\lambda - i0^-}, \quad (\text{S.66})$$

meaning that the Fourier transform of $\chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \tau)$ is

$$\chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \omega) = \sum_{n \neq 0} \frac{\langle \Psi^\lambda | \hat{n}(\mathbf{r}_1) | \Psi_n^\lambda \rangle \langle \Psi_n^\lambda | \hat{n}(\mathbf{r}_2) | \Psi^\lambda \rangle}{\omega - \omega_n^\lambda + i0^+} - \frac{\langle \Psi^\lambda | \hat{n}(\mathbf{r}_2) | \Psi_n^\lambda \rangle \langle \Psi_n^\lambda | \hat{n}(\mathbf{r}_1) | \Psi^\lambda \rangle}{\omega + \omega_n^\lambda - i0^-}. \quad (\text{S.67})$$

After extending the function $\chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \omega)$ on the ω -complex plane by analytic continuation, and noting that the integral of $e^{i\omega 0^+} \chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \omega)$ is zero on the infinite upper semi-circle C according to Jordan's lemma (since $|\chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \omega)|$ goes to zero when $|\omega| \rightarrow +\infty$), the integral of $e^{i\omega 0^+} \chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \omega)$ over the real axis $(-\infty, +\infty)$ is identical to the integral over the closed path $\gamma = (-\infty, +\infty) \cup C$

$$\int_{-\infty}^{\infty} \frac{d\omega}{2\pi i} e^{i\omega 0^+} \chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \omega) = \oint_\gamma \frac{d\omega}{2\pi i} e^{i\omega 0^+} \chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \omega). \quad (\text{S.68})$$

Only the second term in Eq. (S.67) gives poles in the upper-half of the complex plane, $\omega = -\omega_n^\lambda + i0^-$, enclosed by γ , therefore according to the residue theorem only their associated residues contribute to the integral

$$\begin{aligned} \oint_\gamma \frac{d\omega}{2\pi i} e^{i\omega 0^+} \chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \omega) &= - \sum_{n \neq 0} \langle \Psi^\lambda | \hat{n}(\mathbf{r}_2) | \Psi_n^\lambda \rangle \langle \Psi_n^\lambda | \hat{n}(\mathbf{r}_1) | \Psi^\lambda \rangle \\ &= - \left[\sum_n \langle \Psi^\lambda | \hat{n}(\mathbf{r}_2) | \Psi_n^\lambda \rangle \langle \Psi_n^\lambda | \hat{n}(\mathbf{r}_1) | \Psi^\lambda \rangle - \langle \Psi^\lambda | \hat{n}(\mathbf{r}_2) | \Psi^\lambda \rangle \langle \Psi^\lambda | \hat{n}(\mathbf{r}_1) | \Psi^\lambda \rangle \right] \\ &= - \left[\langle \Psi^\lambda | \hat{n}(\mathbf{r}_2) \hat{n}(\mathbf{r}_1) | \Psi^\lambda \rangle - \langle \Psi^\lambda | \hat{n}(\mathbf{r}_2) | \Psi^\lambda \rangle \langle \Psi^\lambda | \hat{n}(\mathbf{r}_1) | \Psi^\lambda \rangle \right]. \end{aligned} \quad (\text{S.69})$$

We thus correctly recover the fluctuation-dissipation theorem of Eq. (4.23)

$$\begin{aligned}
-\int_{-\infty}^{\infty} \frac{d\omega}{2\pi i} e^{i\omega 0^+} [\chi_\lambda(\mathbf{r}_1, \mathbf{r}_2; \omega) - \chi_0(\mathbf{r}_1, \mathbf{r}_2; \omega)] &= \langle \Psi^\lambda | \hat{n}(\mathbf{r}_2) \hat{n}(\mathbf{r}_1) | \Psi^\lambda \rangle - \langle \Phi | \hat{n}(\mathbf{r}_2) \hat{n}(\mathbf{r}_1) | \Phi \rangle \\
&= n_{2,c}^\lambda(\mathbf{r}_1, \mathbf{r}_2).
\end{aligned} \tag{S.70}$$

Exercise 24

The variation of $F[f]$ induced by a variation of $f(x)$ is

$$\begin{aligned}
\delta F[f] &= \int \left[\frac{\partial h}{\partial f(x)} \delta f(x) + \frac{\partial h}{\partial f'(x)} \delta f'(x) + \frac{\partial h}{\partial f''(x)} \delta f''(x) + \cdots + \frac{\partial h}{\partial f^{(n)}(x)} \delta f^{(n)}(x) \right] dx \\
&= \int \left[\frac{\partial h}{\partial f(x)} - \frac{d}{dx} \left(\frac{\partial h}{\partial f'(x)} \right) + \frac{d^2}{dx^2} \left(\frac{\partial h}{\partial f''(x)} \right) + \cdots + (-1)^n \frac{d^n}{dx^n} \left(\frac{\partial h}{\partial f^{(n)}(x)} \right) \right] \delta f(x) dx,
\end{aligned} \tag{S.71}$$

where, in the n^{th} -order term, it was used that $\delta f^{(n)}(x) = [\delta f(x)]^{(n)}$ (since differentiation is a linear operation) and n consecutive integrations by parts were performed (and assuming that all boundary terms vanish). The functional derivative is thus

$$\frac{\delta F[f]}{\delta f(x)} = \frac{\partial h}{\partial f(x)} - \frac{d}{dx} \left(\frac{\partial h}{\partial f'(x)} \right) + \frac{d^2}{dx^2} \left(\frac{\partial h}{\partial f''(x)} \right) + \cdots + (-1)^n \frac{d^n}{dx^n} \left(\frac{\partial h}{\partial f^{(n)}(x)} \right). \tag{S.72}$$

References

- [1] R. G. Parr and W. Yang, *Density-Functional Theory of Atoms and Molecules* (Oxford University Press, New York, 1989).
- [2] R. M. Dreizler and E. K. U. Gross, *Density Functional Theory* (Springer-Verlag, Berlin, 1990).
- [3] W. Koch and M. C. Holthausen, *A Chemist's Guide To Density Functional Theory* (Wiley-VCH, New York, 2001).
- [4] C. Fiolhais, F. Nogueira and M. A. L. Marques, eds., *A Primer in Density Functional Theory*, Vol. 620 of Lecture Notes in Physics (Springer, Berlin, 2003).
- [5] E. Engel and R. M. Dreizler, *Density Functional Theory: An Advanced Course*, Theoretical and Mathematical Physics (Springer-Verlag, Berlin Heidelberg, 2011).
- [6] T. Tsuneda, *Density Functional Theory in Quantum Chemistry* (Springer, Tokyo, 2014).
- [7] J. P. Perdew and K. Schmidt, AIP Conf. Proc. **577**, 1 (2001).
- [8] G. E. Scuseria and V. N. Staroverov, in *Theory and Applications of Computational Chemistry: The First 40 years*, edited by C. E. Dykstra, G. Frenking, K. S. Kim and G. E. Scuseria (Elsevier, 2005), pp. 669–724.
- [9] K. Capelle, Braz. J. Phys. **36**, 1318 (2006).
- [10] A. J. Cohen, P. Mori-Sánchez and W. Yang, Chem. Rev. **112**, 289 (2012).
- [11] K. Burke, J. Chem. Phys. **136**, 150901 (2012).
- [12] V. N. Staroverov, in *A Matter of Density: Exploring the Electron Density Concept in the Chemical, Biological, and Materials Sciences*, edited by N. Sukumar (John Wiley & Sons, Hoboken, NJ, 2013), pp. 125–156.
- [13] A. D. Becke, J. Chem. Phys. **140**, 18A301 (2014).
- [14] R. O. Jones, Rev. Mod. Phys. **87**, 897 (2015).
- [15] H. S. Yu, S. L. Li and D. G. Truhlar, J. Chem. Phys. **145**, 130901 (2016).
- [16] N. Mardirossian and M. Head-Gordon, Mol. Phys. **115**, 2315 (2017).
- [17] J. Toulouse, in *Density Functional Theory*, edited by E. Cancès, G. Friesecke and L. Lin (Springer), to appear; arxiv.org/abs/2103.02645.
- [18] P. Hohenberg and W. Kohn, Phys. Rev. **136**, B 864 (1964).
- [19] M. Levy, Proc. Natl. Acad. Sci. U.S.A. **76**, 6062 (1979).
- [20] M. Levy, Phys. Rev. A **26**, 1200 (1982).
- [21] E. H. Lieb, Int. J. Quantum Chem. **24**, 243 (1983).
- [22] S. M. Valone, J. Chem. Phys. **73**, 4653 (1980).
- [23] W. Kohn and L. J. Sham, Phys. Rev. **140**, A1133 (1965).
- [24] A. D. Becke, J. Chem. Phys. **88**, 2547 (1988).
- [25] U. von Barth and L. Hedin, J. Phys. C **5**, 1629 (1972).
- [26] A. K. Rajagopal and J. Callaway, Phys. Rev. B **7**, 1912 (1973).
- [27] J. P. Perdew and A. Zunger, Phys. Rev. B **23**, 5048 (1981).
- [28] G. L. Oliver and J. P. Perdew, Phys. Rev. A **20**, 397 (1979).
- [29] A. Seidl, A. Görling, P. Vogl, J. A. Majewski and M. Levy, Phys. Rev. B **53**, 3764 (1996).
- [30] M. Levy and J. P. Perdew, Phys. Rev. A **32**, 2010 (1985).
- [31] M. Levy, Phys. Rev. A **43**, 4637 (1991).
- [32] M. Levy, in *Density Functional Theory*, edited by E. Gross and R. Dreizler (Plenum Press, New York, 1995).
- [33] A. Görling and M. Levy, Phys. Rev. B **47**, 13105 (1993).
- [34] A. Görling and M. Levy, Phys. Rev. A **50**, 196 (1994).
- [35] C.-J. Huang and C. J. Umrigar, Phys. Rev. A **56**, 290 (1997).
- [36] M. Seidl, J. P. Perdew and M. Levy, Phys. Rev. A **59**, 51 (1999).
- [37] M. Seidl, Phys. Rev. A **60**, 4387 (1999).

- [38] M. Seidl, P. Gori-Giorgi and A. Savin, Phys. Rev. A **75**, 042511 (2007).
- [39] P. Gori-Giorgi and M. Seidl, Phys. Chem. Chem. Phys. **12**, 14405 (2010).
- [40] E. H. Lieb and S. Oxford, Int. J. Quantum. Chem. **19**, 427 (1981).
- [41] J. P. Perdew, in *Electronic Structure of Solids '91*, edited by P. Ziesche and H. Eschrig (Akademie Verlag, Berlin, 1991).
- [42] J. P. Perdew, R. G. Parr, M. Levy and J. L. Balduz, Phys. Rev. Lett. **49**, 1691 (1982).
- [43] J. F. Janak, Phys. Rev. B **18**, 7165 (1978).
- [44] M. Levy, J. P. Perdew and V. Sahni, Phys. Rev. A **30**, 2745 (1984).
- [45] E. Sagvolden and J. P. Perdew, Phys. Rev. A **77**, 012517 (2008).
- [46] P. Gori-Giorgi and A. Savin, Int. J. Quantum Chem. **109**, 2410 (2009).
- [47] T. Gould and J. Toulouse, Phys. Rev. A **90**, 050502(R) (2014).
- [48] R. Peverati and D. G. Truhlar, Phil. Trans. R. Soc. A **372**, 20120476 (2014), preprint at <http://arxiv.org/abs/1212.0944>.
- [49] P. A. M. Dirac, Proc. Cambridge Phil. Soc. **26**, 376 (1930).
- [50] J. C. Slater, Phys. Rev. **81**, 385 (1951).
- [51] D. M. Ceperley and B. J. Alder, Phys. Rev. Lett. **45**, 566 (1980).
- [52] S. J. Vosko, L. Wilk and M. Nusair, Can. J. Phys. **58**, 1200 (1980).
- [53] J. P. Perdew and Y. Wang, Phys. Rev. B **45**, 13244 (1992).
- [54] P.-F. Loos and P. M. W. Gill, WIREs Comput. Mol. Sci. **6**, 410 (2016), doi: 10.1002/wcms.1257.
- [55] A. D. Becke, Phys. Rev. A **38**, 3098 (1988).
- [56] C. Lee, W. Yang and R. G. Parr, Phys. Rev. B **37**, 785 (1988).
- [57] R. Colle and O. Salvetti, Theor. Chim. Acta **37**, 329 (1975).
- [58] B. Miehlich, A. Savin, H. Stoll and H. Preuss, Chem. Phys. Lett. **157**, 200 (1989).
- [59] J. P. Perdew, J. A. Chevary, S. H. Vosko, K. A. Jackson, M. R. Pederson, D. J. Singh and C. Fiolhais, Phys. Rev. B **46**, 6671 (1992).
- [60] K. Burke, J. P. Perdew and Y. Wang, *Derivation of a generalized gradient approximation: The PW91 density functional* (Plenum, NY, 1997), p. 81.
- [61] J. P. Perdew, K. Burke and M. Ernzerhof, Phys. Rev. Lett. **77**, 3865 (1996).
- [62] A. D. Becke, Int. J. Quantum. Chem. **23**, 1915 (1983).
- [63] J. Tao, J. P. Perdew, V. N. Staroverov and G. E. Scuseria, Phys. Rev. Lett. **91**, 146401 (2003).
- [64] J. Sun, A. Ruzsinszky and J. P. Perdew, Phys. Rev. Lett. **115**, 036402 (2015).
- [65] A. D. Becke, J. Chem. Phys. **98**, 1372 (1993).
- [66] A. D. Becke, J. Chem. Phys. **98**, 5648 (1993).
- [67] P. J. Stephens, F. J. Devlin, C. F. Chabalowski and M. J. Frisch, J. Phys. Chem. **98**, 11623 (1994).
- [68] A. D. Becke, J. Chem. Phys. **104**, 1040 (1996).
- [69] J. P. Perdew, M. Ernzerhof and K. Burke, J. Chem. Phys. **105**, 9982 (1996).
- [70] C. Adamo and V. Barone, J. Chem. Phys. **110**, 6158 (1999).
- [71] M. Ernzerhof and G. E. Scuseria, J. Chem. Phys. **110**, 5029 (1999).
- [72] A. D. Becke, J. Chem. Phys. **107**, 8554 (1997).
- [73] Y. Zhao and D. G. Truhlar, Theor. Chem. Acc. **120**, 215 (2008).
- [74] A. Savin, in *Recent Developments of Modern Density Functional Theory*, edited by J. M. Seminario (Elsevier, Amsterdam, 1996), pp. 327–357.
- [75] H. Iikura, T. Tsuneda, T. Yanai and K. Hirao, J. Chem. Phys. **115**, 3540 (2001).
- [76] B. Mussard and J. Toulouse, Mol. Phys. **115**, 161 (2017).
- [77] O. A. Vydrov and G. E. Scuseria, J. Chem. Phys. **125**, 234109 (2006).
- [78] I. C. Gerber and J. G. Ángyán, Chem. Phys. Lett. **415**, 100 (2005).
- [79] T. Yanai, D. P. Tew and N. C. Handy, Chem. Phys. Lett. **393**, 51 (2004).
- [80] J.-D. Chai and M. Head-Gordon, J. Chem. Phys. **128**, 084106 (2008).

- [81] J. Heyd, G. E. Scuseria and M. Ernzerhof, *J. Chem. Phys.* **118**, 8207 (2003).
- [82] S. Grimme, *J. Chem. Phys.* **124**, 034108 (2006).
- [83] K. Sharkas, J. Toulouse and A. Savin, *J. Chem. Phys.* **134**, 064113 (2011).
- [84] E. Fromager, *J. Chem. Phys.* **135**, 244106 (2011).
- [85] K. Sharkas, A. Savin, H. J. A. Jensen and J. Toulouse, *J. Chem. Phys.* **137**, 044104 (2012).
- [86] J. G. Ángyán, I. C. Gerber, A. Savin and J. Toulouse, *Phys. Rev. A* **72**, 012510 (2005).
- [87] O. Franck, B. Mussard, E. Luppi and J. Toulouse, *J. Chem. Phys.* **142**, 074107 (2015).
- [88] E. Goll, H.-J. Werner and H. Stoll, *Phys. Chem. Chem. Phys.* **7**, 3917 (2005).
- [89] J. Toulouse, I. C. Gerber, G. Jansen, A. Savin and J. G. Ángyán, *Phys. Rev. Lett.* **102**, 096404 (2009).
- [90] B. G. Janesko, T. M. Henderson and G. E. Scuseria, *J. Chem. Phys.* **130**, 081105 (2009).
- [91] J. Toulouse, W. Zhu, A. Savin, G. Jansen and J. G. Ángyán, *J. Chem. Phys.* **135**, 084119 (2011).
- [92] Y. Cornaton and E. Fromager, *Int. J. Quantum Chem.* **114**, 1199 (2014).
- [93] A. J. Garza, I. W. Bulik, T. M. Henderson and G. E. Scuseria, *Phys. Chem. Chem. Phys.* **17**, 22412 (2015).
- [94] C. Kalai and J. Toulouse, *J. Chem. Phys.* **148**, 164105 (2018).
- [95] C. Kalai, B. Mussard and J. Toulouse, *J. Chem. Phys.* **151**, 074102 (2019).
- [96] M. Elstner, P. Hobza, T. Frauenheim, S. Suhai and E. Kaxiras, *J. Chem. Phys.* **114**, 5149 (2001).
- [97] Q. Wu and W. Yang, *J. Chem. Phys.* **116**, 515 (2002).
- [98] S. Grimme, *J. Comput. Chem.* **25**, 1463 (2004).
- [99] S. Grimme, J. Antony, S. Ehrlich and H. Krieg, *J. Chem. Phys.* **132**, 154104 (2010).
- [100] S. Grimme, *J. Comput. Chem.* **27**, 1787 (2006).
- [101] J.-D. Chai and M. Head-Gordon, *Phys. Chem. Chem. Phys.* **10**, 6615 (2008).
- [102] T. Schwabe and S. Grimme, *Phys. Chem. Chem. Phys.* **9**, 3397 (2007).
- [103] A. D. Becke and E. R. Johnson, *J. Chem. Phys.* **127**, 154108 (2007).
- [104] A. Tkatchenko and M. Scheffler, *Phys. Rev. Lett.* **102**, 073005 (2009).
- [105] T. Sato and H. Nakai, *J. Chem. Phys.* **133**, 194101 (2010).
- [106] J. F. Dobson and B. P. Dinte, *Phys. Rev. Lett.* **76**, 1780 (1996).
- [107] M. Dion, H. Rydberg, E. Schröder, D. C. Langreth and B. I. Lundqvist, *Phys. Rev. Lett.* **92**, 246401 (2004).
- [108] K. Lee, E. D. Murray, L. Kong, B. I. Lundqvist and D. C. Langreth, *Phys. Rev. B* **82**, 081101 (2010).
- [109] O. A. Vydrov and T. Van Voorhis, *J. Chem. Phys.* **130**, 104105 (2009).
- [110] O. A. Vydrov and T. Van Voorhis, *Phys. Rev. Lett.* **103**, 063004 (2009).
- [111] O. A. Vydrov and T. Van Voorhis, *J. Chem. Phys.* **133**, 244103 (2010).
- [112] N. Mardirossian and M. Head-Gordon, *Phys. Chem. Chem. Phys.* **16**, 9904 (2014).
- [113] N. Mardirossian and M. Head-Gordon, *J. Chem. Phys.* **144**, 214110 (2016).
- [114] R. T. Sharp and G. K. Horton, *Phys. Rev.* **90**, 317 (1953).
- [115] J. D. Talman and W. F. Shadwick, *Phys. Rev. A* **14**, 36 (1976).
- [116] S. Ivanov, S. Hirata and R. J. Bartlett, *Phys. Rev. Lett.* **83**, 5455 (1999).
- [117] A. Görling, *Phys. Rev. Lett.* **83**, 5459 (1999).
- [118] I. Grabowski, S. Hirata, S. Ivanov and R. J. Bartlett, *J. Chem. Phys.* **116**, 4415 (2002).
- [119] E. Engel, in *A Primer in Density Functional Theory*, edited by C. Fiolhais, F. Nogueira and M. A. L. Marques (Springer, Berlin, 2003), Vol. 620 of Lecture Notes in Physics, pp. 56–122.
- [120] P. Mori-Sánchez, Q. Wu and W. Yang, *J. Chem. Phys.* **123**, 062204 (2005).
- [121] F. Furche, *Phys. Rev. B* **64**, 195120 (2001).
- [122] G. E. Scuseria, T. M. Henderson and D. C. Sorensen, *J. Chem. Phys.* **129**, 231101 (2008).

- [123] J. Toulouse, W. Zhu, J. G. Ángyán and A. Savin, *Phys. Rev. A* **82**, 032502 (2010).
- [124] J. G. Ángyán, R.-F. Liu, J. Toulouse and G. Jansen, *J. Chem. Theory Comput.* **7**, 3116 (2011).
- [125] H. Eshuis, J. Bates and F. Furche, *Theor. Chem. Acc.* **131**, 1084 (2012).
- [126] B. Mussard, D. Rocca, G. Jansen and J. G. Ángyán, *J. Chem. Theory Comput.* **12**, 2191 (2016).
- [127] M. Hellgren, D. R. Rohr and E. K. U. Gross, *J. Chem. Phys.* **136**, 034106 (2012).
- [128] P. Bleiziffer, A. Heßelmann and A. Görling, *J. Chem. Phys.* **139**, 084113 (2013).
- [129] M. Hellgren and U. von Barth, *Phys. Rev. B* **78**, 115107 (2008).
- [130] M. Hellgren and U. von Barth, *J. Chem. Phys.* **132**, 044101 (2010).
- [131] A. Heßelmann and A. Görling, *Mol. Phys.* **108**, 359 (2010).
- [132] A. Heßelmann and A. Görling, *Phys. Rev. Lett.* **106**, 093001 (2011).
- [133] J. Erhard, P. Bleiziffer and A. Görling, *Phys. Rev. Lett.* **117**, 143002 (2016).
- [134] A. L. Fetter and J. D. Walecka, *Quantum Theory of Many-Particle Systems* (Dover, 2003).