

Introduction to density-functional theory

Julien Toulouse

Laboratoire de Chimie Théorique
Sorbonne Université and **CNRS**, Paris, France

Course 5CI228 of Master 2 Chemistry
Sorbonne Université, Paris

Why and how learning density-functional theory?

Density-functional theory (DFT) is:

- ▶ a **practical electronic-structure computational method**, widely used in quantum chemistry and condensed-matter physics;
- ▶ an **exact and elegant reformulation of the quantum many-body problem**, which has led to new ways of thinking in the field.

Classical books:

- ▶ R. G. Parr and W. Yang, *Density-Functional Theory of Atoms and Molecules*, Oxford University Press, 1989.
- ▶ R. M. Dreizler and E. K. U. Gross, *Density Functional Theory: An Approach to the Quantum Many-Body Problem*, Springer-Verlag, 1990.
- ▶ W. Koch and M. C. Holthausen, *A Chemist's Guide To Density Functional Theory*, Wiley-VCH, 2001.

My lecture notes:

http://www.lct.jussieu.fr/pagesperso/toulouse/enseignement/introduction_dft.pdf

A book chapter:

J. Toulouse, in *Density Functional Theory*, edited by E. Cancès and G. Friesecke, Springer, 2023.

- 1 Basic density-functional theory
 - Quantum many-electron problem
 - Universal density functional
 - Kohn-Sham method
 - Generalized Kohn-Sham method
- 2 Exact constraints for the exchange-correlation functional
 - Exact expressions for the exchange and correlation functionals
 - Uniform coordinate scaling
 - One-orbital spatial regions and self-interaction
 - Lieb-Oxford lower bound
 - Frontier orbital energies
- 3 Approximations for the exchange-correlation energy
 - Local-density approximation
 - Semilocal approximations
 - Single-determinant hybrid approximations
 - Multideterminant hybrid approximations
 - Dispersion corrections
 - Machine-learned density functionals
 - Benchmark
- 4 Time-dependent density-functional theory
 - Runge-Gross theorem
 - Linear-response TDDFT

1 Basic density-functional theory

- Quantum many-electron problem
- Universal density functional
 - The Hohenberg-Kohn theorem
 - Levy-Lieb constrained-search formulation
- Kohn-Sham method
 - Decomposition of the universal functional
 - The Kohn-Sham equations
 - Practical calculations in an atomic basis
 - Extension to spin density-functional theory
- Generalized Kohn-Sham method

- ▶ We consider an **N -electron system** in the Born-Oppenheimer and non-relativistic approximations.
- ▶ The **electronic Hamiltonian** in the position representation is, in atomic units,

$$\hat{H} = -\frac{1}{2} \sum_i^N \nabla_{\mathbf{r}_i}^2 + \frac{1}{2} \sum_i^N \sum_{j \neq i}^N \frac{1}{|\mathbf{r}_i - \mathbf{r}_j|} + \sum_i^N v_{\text{ne}}(\mathbf{r}_i)$$

where $v_{\text{ne}}(\mathbf{r}_i) = -\sum_{\alpha} Z_{\alpha}/|\mathbf{r}_i - \mathbf{R}_{\alpha}|$ is the nuclei-electron interaction potential.

- ▶ Stationary states satisfy the **time-independent Schrödinger equation**

$$H\Psi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N) = E\Psi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N)$$

where $\Psi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N)$ is a wave function written with space-spin coordinates $\mathbf{x}_i = (\mathbf{r}_i, \sigma_i)$ (with $\mathbf{r}_i \in \mathbb{R}^3$ and $\sigma_i \in \{\uparrow, \downarrow\}$) which is antisymmetric with respect to the exchange of two space-spin coordinates, and E is the associated energy.

- ▶ Using **Dirac notations** (representation-independent formalism):

$$\hat{H}|\Psi\rangle = E|\Psi\rangle$$

where

$$\hat{H} = \hat{T} + \hat{W}_{\text{ee}} + \hat{V}_{\text{ne}}$$

- ▶ The **ground-state electronic energy** E_0 can be expressed with the **wave-function variational principle**

$$E_0 = \min_{\Psi} \langle \Psi | \hat{H} | \Psi \rangle$$

where the minimization is over all N -electron (multi-determinant) wave functions Ψ normalized to unity $\langle \Psi | \Psi \rangle = 1$.

- ▶ DFT is based on a reformulation of this variational theorem in terms of the **one-electron density** defined as

$$n(\mathbf{r}) = N \int \cdots \int |\Psi(\mathbf{x}, \mathbf{x}_2, \dots, \mathbf{x}_N)|^2 d\sigma d\mathbf{x}_2 \dots d\mathbf{x}_N$$

which is normalized to the electron number, $\int n(\mathbf{r}) d\mathbf{r} = N$.

Remark: Integration over a spin coordinate σ means a sum over the two values of σ , i.e.
 $\int d\sigma = \sum_{\sigma \in \{\uparrow, \downarrow\}}$

1 Basic density-functional theory

- Quantum many-electron problem
- Universal density functional
 - The Hohenberg-Kohn theorem
 - Levy-Lieb constrained-search formulation
- Kohn-Sham method
 - Decomposition of the universal functional
 - The Kohn-Sham equations
 - Practical calculations in an atomic basis
 - Extension to spin density-functional theory
- Generalized Kohn-Sham method

The Hohenberg-Kohn theorem

- ▶ Consider an electronic system with an arbitrary external local potential $v(\mathbf{r})$ (that bounds N electrons) in place of $v_{\text{ne}}(\mathbf{r})$. For simplicity, we will assume that $v(\mathbf{r})$ gives an N -electron ground state which is not degenerate.
- ▶ For any such external potential $v(\mathbf{r})$, the ground-state wave function Ψ can be obtained by solving the Schrödinger equation, from which an associated **ground-state density** $n(\mathbf{r})$ can be deduced. Therefore, one has the mapping:

$$v(\mathbf{r}) \longrightarrow n(\mathbf{r})$$

- ▶ In 1964, Hohenberg and Kohn showed that this mapping can be inverted, i.e. a **ground-state density** $n(\mathbf{r})$ determines the potential $v(\mathbf{r})$ up to an arbitrary additive constant:

$$n(\mathbf{r}) \xrightarrow{\text{Hohenberg-Kohn}} v(\mathbf{r}) + \text{const}$$

Proof of the Hohenberg-Kohn theorem (1/2)

This is a two-step proof by contradiction.

Consider two local potentials differing by more than an additive constant:

$$v_1(\mathbf{r}) \neq v_2(\mathbf{r}) + \text{const}$$

We have two Hamiltonians:

$\hat{H}_1 = \hat{T} + \hat{W}_{\text{ee}} + \hat{V}_1$ with a ground state $\hat{H}_1|\Psi_1\rangle = E_1|\Psi_1\rangle$ and ground-state density $n_1(\mathbf{r})$

$\hat{H}_2 = \hat{T} + \hat{W}_{\text{ee}} + \hat{V}_2$ with a ground state $\hat{H}_2|\Psi_2\rangle = E_2|\Psi_2\rangle$ and ground-state density $n_2(\mathbf{r})$

1 We first show that $\Psi_1 \neq \Psi_2$:

Assume $\Psi_1 = \Psi_2 = \Psi$. Then we have:

$$(\hat{H}_1 - \hat{H}_2)|\Psi\rangle = (\hat{V}_1 - \hat{V}_2)|\Psi\rangle = (E_1 - E_2)|\Psi\rangle$$

or, in position representation,

$$\left(\sum_i^N [v_1(\mathbf{r}_i) - v_2(\mathbf{r}_i)] \right) \Psi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N) = (E_1 - E_2)\Psi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N)$$

If $\Psi(\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N) \neq 0$ for at least one fixed set of $(\sigma_1, \sigma_2, \dots, \sigma_N)$ and “almost” all $(\mathbf{r}_1, \mathbf{r}_2, \dots, \mathbf{r}_N)$, which is true for “reasonably well behaved potentials”, then it implies that $v_1(\mathbf{r}) - v_2(\mathbf{r}) = \text{const}$, in contradiction with the initial hypothesis.

$\implies$ Intermediate conclusion: two local potentials differing by more than an additive constant cannot have a common ground-state wave function.

Proof of the Hohenberg-Kohn theorem (2/2)

2 We now show that $n_1 \neq n_2$:

Assume $n_1 = n_2 = n$. Then, by the variational theorem, we have:

$$E_1 = \langle \Psi_1 | \hat{H}_1 | \Psi_1 \rangle < \langle \Psi_2 | \hat{H}_1 | \Psi_2 \rangle = \langle \Psi_2 | \hat{H}_2 + \hat{V}_1 - \hat{V}_2 | \Psi_2 \rangle = E_2 + \int [v_1(\mathbf{r}) - v_2(\mathbf{r})] n(\mathbf{r}) d\mathbf{r}$$

The strict inequality comes from the fact that Ψ_2 cannot be a ground-state wave function of $\hat{H}_1$, as shown in the first step of the proof.

So, we have shown

$$E_1 < E_2 + \int [v_1(\mathbf{r}) - v_2(\mathbf{r})] n(\mathbf{r}) d\mathbf{r}$$

Symmetrically, by exchanging the role of system 1 and 2, we have the strict inequality

$$E_2 < E_1 + \int [v_2(\mathbf{r}) - v_1(\mathbf{r})] n(\mathbf{r}) d\mathbf{r}$$

Adding the two inequalities gives the inconsistent result

$$E_1 + E_2 < E_2 + E_1$$

$\Rightarrow$ **Conclusion:** there cannot exist two local potentials differing by more than an additive constant which have a common ground-state density.

- ▶ The Hohenberg-Kohn theorem can be summarized as

$$n \longrightarrow v \longrightarrow \hat{H} \longrightarrow \text{everything}$$

v is a functional of the density n , i.e. $v[n]$, and all other quantities as well.

- ▶ The **ground-state wave function** Ψ is a functional of n , denoted by $\Psi[n]$. Hohenberg and Kohn defined the **universal density functional**

$$F[n] = \langle \Psi[n] | \hat{T} + \hat{W}_{ee} | \Psi[n] \rangle$$

and the **total electronic energy functional**

$$E[n] = F[n] + \int v_{ne}(\mathbf{r})n(\mathbf{r})d\mathbf{r}$$

- ▶ Hohenberg and Kohn showed that we have a **variational property** giving the exact ground-state energy

$$E_0 = \min_n \left\{ F[n] + \int v_{ne}(\mathbf{r})n(\mathbf{r})d\mathbf{r} \right\}$$

The minimum is reached for the exact ground-state density $n_0(\mathbf{r})$ of the potential $v_{ne}(\mathbf{r})$.


- In 1979 Levy, and later in 1983 Lieb, proposed to redefine the **universal density functional** as

$$F[n] = \min_{\Psi \rightarrow n} \langle \Psi | \hat{T} + \hat{W}_{ee} | \Psi \rangle = \langle \Psi[n] | \hat{T} + \hat{W}_{ee} | \Psi[n] \rangle$$

where “ $\Psi \rightarrow n$ ” means that the minimization is over all normalized N -electron (multi-determinant) wave functions **giving the density** n .

- The **variational property** is easily obtained using the constrained-search formulation:

$$\begin{aligned} E_0 &= \min_{\Psi} \langle \Psi | \hat{T} + \hat{W}_{ee} + \hat{V}_{ne} | \Psi \rangle \\ &= \min_n \min_{\Psi \rightarrow n} \langle \Psi | \hat{T} + \hat{W}_{ee} + \hat{V}_{ne} | \Psi \rangle \\ &= \min_n \left\{ \min_{\Psi \rightarrow n} \langle \Psi | \hat{T} + \hat{W}_{ee} | \Psi \rangle + \int v_{ne}(\mathbf{r}) n(\mathbf{r}) d\mathbf{r} \right\} \\ &= \min_n \left\{ F[n] + \int v_{ne}(\mathbf{r}) n(\mathbf{r}) d\mathbf{r} \right\} \end{aligned}$$



- Hence, in DFT, we replace “ $\min_{\Psi}$ ” by “ $\min_n$ ” which is a **tremendous simplification!** However, $F[n] = T[n] + W_{ee}[n]$ is **very difficult to approximate**, in particular the kinetic energy part $T[n]$.

1 Basic density-functional theory

- Quantum many-electron problem
- Universal density functional
 - The Hohenberg-Kohn theorem
 - Levy-Lieb constrained-search formulation
- Kohn-Sham method
 - Decomposition of the universal functional
 - The Kohn-Sham equations
 - Practical calculations in an atomic basis
 - Extension to spin density-functional theory
- Generalized Kohn-Sham method

- ▶ In 1965, Kohn and Sham proposed to decompose $F[n]$ as

$$F[n] = T_s[n] + E_{\text{Hxc}}[n]$$

- ▶ $T_s[n]$ is the **non-interacting kinetic-energy functional**:

$$T_s[n] = \min_{\Phi \rightarrow n} \langle \Phi | \hat{T} | \Phi \rangle = \langle \Phi[n] | \hat{T} | \Phi[n] \rangle$$

where the minimization is over normalized N -electron **single-determinant** wave functions Φ giving the fixed density n . The minimizing single-determinant wave function (assumed to be unique for simplicity) is called the **KS wave function** and is denoted by $\Phi[n]$.

- ▶ The remaining functional $E_{\text{Hxc}}[n]$ is called the **Hartree-exchange-correlation functional**.
- ▶ Any density n can be obtained from a single-determinant wave function, therefore the **Kohn-Sham decomposition does not introduce any approximation**.

- The **exact ground-state energy** can then be expressed as

$$\begin{aligned}
 E_0 &= \min_n \left\{ F[n] + \int v_{\text{ne}}(\mathbf{r})n(\mathbf{r})d\mathbf{r} \right\} \\
 &= \min_n \left\{ \min_{\Phi \rightarrow n} \langle \Phi | \hat{T} | \Phi \rangle + E_{\text{Hxc}}[n] + \int v_{\text{ne}}(\mathbf{r})n(\mathbf{r})d\mathbf{r} \right\} \\
 &= \min_n \min_{\Phi \rightarrow n} \left\{ \langle \Phi | \hat{T} + \hat{V}_{\text{ne}} | \Phi \rangle + E_{\text{Hxc}}[n_{\Phi}] \right\} \\
 &= \min_{\Phi} \left\{ \langle \Phi | \hat{T} + \hat{V}_{\text{ne}} | \Phi \rangle + E_{\text{Hxc}}[n_{\Phi}] \right\}
 \end{aligned}$$

and the minimizing single-determinant KS wave function gives the exact ground-state density $n_0(\mathbf{r})$.

- Hence, in KS DFT, we replace “ $\min_{\Psi}$ ” by “ $\min_{\Phi}$ ” which is still a **tremendous simplification!** The advantage of KS DFT over pure DFT is that a major part of the kinetic energy is treated explicitly with the single-determinant wave function Φ .
- KS DFT is similar to Hartree-Fock (HF)

$$E_{\text{HF}} = \min_{\Phi} \langle \Phi | \hat{T} + \hat{V}_{\text{ne}} + \hat{W}_{\text{ee}} | \Phi \rangle$$

but in KS DFT the exact ground-state energy and density are in principle obtained!

Kohn-Sham (KS) method: the Hartree-exchange-correlation functional

- ▶ $E_{\text{Hxc}}[n]$ is decomposed as

$$E_{\text{Hxc}}[n] = E_{\text{H}}[n] + E_{\text{xc}}[n]$$

- ▶ $E_{\text{H}}[n]$ is the **Hartree energy functional**

$$E_{\text{H}}[n] = \frac{1}{2} \iint \frac{n(\mathbf{r}_1)n(\mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2$$

representing the classical electrostatic repulsion energy for the charge distribution $n(\mathbf{r})$ and which is calculated exactly.

- ▶ $E_{\text{xc}}[n]$ is the **exchange-correlation energy functional** that remains to approximate. This functional is often decomposed as

$$E_{\text{xc}}[n] = E_{\text{x}}[n] + E_{\text{c}}[n]$$

where $E_{\text{x}}[n]$ is the **exchange energy functional**

$$E_{\text{x}}[n] = \langle \Phi[n] | \hat{W}_{\text{ee}} | \Phi[n] \rangle - E_{\text{H}}[n]$$

and $E_{\text{c}}[n]$ is the **correlation energy functional**

$$E_{\text{c}}[n] = \langle \Psi[n] | \hat{T} + \hat{W}_{\text{ee}} | \Psi[n] \rangle - \langle \Phi[n] | \hat{T} + \hat{W}_{\text{ee}} | \Phi[n] \rangle = T_{\text{c}}[n] + U_{\text{c}}[n]$$

containing a kinetic contribution $T_{\text{c}}[n] = \langle \Psi[n] | \hat{T} | \Psi[n] \rangle - \langle \Phi[n] | \hat{T} | \Phi[n] \rangle$

and a potential contribution $U_{\text{c}}[n] = \langle \Psi[n] | \hat{W}_{\text{ee}} | \Psi[n] \rangle - \langle \Phi[n] | \hat{W}_{\text{ee}} | \Phi[n] \rangle$.

The Kohn-Sham equations (1/2)

- ▶ The single determinant Φ is constructed from a set of N **orthonormal occupied spin-orbitals** $\chi_a(\mathbf{x}) = \psi_a(\mathbf{r})\alpha(\sigma)$ or $\chi_a(\mathbf{x}) = \psi_a(\mathbf{r})\beta(\sigma)$.
- ▶ The **total energy** to be minimized is

$$E[\{\psi_a\}] = \sum_a^N \int \psi_a^*(\mathbf{r}) \left(-\frac{1}{2} \nabla^2 + v_{\text{ne}}(\mathbf{r}) \right) \psi_a(\mathbf{r}) d\mathbf{r} + E_{\text{Hxc}}[n]$$

and the **density** is

$$n(\mathbf{r}) = \sum_a^N |\psi_a(\mathbf{r})|^2$$

- ▶ For minimizing over the orbitals $\{\psi_a\}$ with the constraint of keeping the orbitals orthonormalized, we introduce the **Lagrangian**

$$\mathcal{L}[\{\psi_a\}] = E[\{\psi_a\}] - \sum_a^N \varepsilon_a \left(\int \psi_a^*(\mathbf{r}) \psi_a(\mathbf{r}) d\mathbf{r} - 1 \right)$$

where ε_a is the Lagrange multiplier associated with the normalization condition of $\psi_a(\mathbf{r})$.

- ▶ The Lagrangian must be **stationary** with respect to variations of the orbitals $\psi_a(\mathbf{r})$

$$\frac{\delta \mathcal{L}}{\delta \psi_a^*(\mathbf{r})} = 0$$

Interlude: Review on functional derivatives

- ▶ For a **functional** $F : f \mapsto F[f]$ of the function $f : x \mapsto f(x)$, an infinitesimal variation δf of f leads to an infinitesimal variation of F which can be expressed as

$$\delta F[f] = \int \frac{\delta F[f]}{\delta f(x)} \delta f(x) dx$$

This defines the **functional derivative** of $F[f]$ with respect to $f(x)$: $\frac{\delta F[f]}{\delta f(x)}$

Remark: For a function $F(f_1, f_2, \dots)$ of several variables $f_1, f_2, \dots$, we have

$$dF = \sum_i \frac{\partial F}{\partial f_i} df_i$$

$\delta F[f]/\delta f(x)$ is the analog of $\partial F/\partial f_i$ for the case of an infinite continuous number of variables.

- ▶ For a functional $F[f]$ of a function $f[g](x)$ which is itself a functional of another function $g(x)$, we have the **chain rule**

$$\frac{\delta F}{\delta g(x)} = \int \frac{\delta F}{\delta f(x')} \frac{\delta f(x')}{\delta g(x)} dx'$$

Remark: It is the analog of the chain rule for a function $F(f_1, f_2, \dots)$ of several variables $f_j(g_1, g_2, \dots)$ which are themselves functions of other variables $g_1, g_2, \dots$

$$\frac{\partial F}{\partial g_i} = \sum_j \frac{\partial F}{\partial f_j} \frac{\partial f_j}{\partial g_i}$$

The Kohn-Sham equations (2/2)

- We find for the functional derivative of the Lagrangian

$$0 = \frac{\delta \mathcal{L}}{\delta \psi_a^*(\mathbf{r})} = \left(-\frac{1}{2} \nabla^2 + v_{\text{ne}}(\mathbf{r}) \right) \psi_a(\mathbf{r}) + \frac{\delta E_{\text{Hxc}}[n]}{\delta \psi_a^*(\mathbf{r})} - \varepsilon_a \psi_a(\mathbf{r})$$

- We calculate the term $\delta E_{\text{Hxc}}[n]/\delta \psi_a^*(\mathbf{r})$ using the chain rule

$$\frac{\delta E_{\text{Hxc}}[n]}{\delta \psi_a^*(\mathbf{r})} = \int \frac{\delta E_{\text{Hxc}}[n]}{\delta n(\mathbf{r}')} \frac{\delta n(\mathbf{r}')}{\delta \psi_a^*(\mathbf{r})} d\mathbf{r}' = v_{\text{Hxc}}(\mathbf{r}) \psi_a(\mathbf{r})$$

where we have used $\delta n(\mathbf{r}')/\delta \psi_a^*(\mathbf{r}) = \psi_a(\mathbf{r})\delta(\mathbf{r} - \mathbf{r}')$ and we have introduced the **Hartree-exchange-correlation potential** $v_{\text{Hxc}}(\mathbf{r})$

$$v_{\text{Hxc}}(\mathbf{r}) = \frac{\delta E_{\text{Hxc}}[n]}{\delta n(\mathbf{r})}$$

which is itself a functional of the density.

- We arrive at the **KS equations**

$$\left(-\frac{1}{2} \nabla^2 + v_{\text{ne}}(\mathbf{r}) + v_{\text{Hxc}}(\mathbf{r}) \right) \psi_a(\mathbf{r}) = \varepsilon_a \psi_a(\mathbf{r})$$

The orbitals $\psi_a(\mathbf{r})$ are called the KS orbitals and ε_a are the KS orbital energies.

- ▶ The KS orbitals are eigenfunctions of the KS one-electron Hamiltonian

$$\hat{h}_s = -\frac{1}{2}\nabla^2 + v_s(\mathbf{r})$$

where $v_s(\mathbf{r}) = v_{ne}(\mathbf{r}) + v_{Hxc}(\mathbf{r})$ is the KS potential.

- ▶ Mathematically, the KS equations are a set of coupled self-consistent equations since the potential $v_{Hxc}(\mathbf{r})$ depends on all the occupied orbitals $\{\psi_a\}_{a=1,\dots,N}$ through the density.
- ▶ Physically, $\hat{h}_s$ defines the KS system which is a system of N non-interacting electrons in an effective external potential $v_s(\mathbf{r})$ ensuring that its ground-state density $n(\mathbf{r})$ is the same as the exact ground-state density $n_0(\mathbf{r})$ of the physical system of N interacting electrons.
- ▶ The KS equations also defines virtual KS orbitals $\{\psi_r\}_{r\geq N+1}$.

The Hartree-exchange-correlation potential

- ▶ Following the decomposition of $E_{\text{Hxc}}[n]$, the potential $v_{\text{Hxc}}(\mathbf{r})$ is also decomposed as

$$v_{\text{Hxc}}(\mathbf{r}) = v_{\text{H}}(\mathbf{r}) + v_{\text{xc}}(\mathbf{r})$$

with the **Hartree potential**

$$v_{\text{H}}(\mathbf{r}) = \frac{\delta E_{\text{H}}[n]}{\delta n(\mathbf{r})} = \int \frac{n(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} d\mathbf{r}'$$

and the **exchange-correlation potential**

$$v_{\text{xc}}(\mathbf{r}) = \delta E_{\text{xc}}[n] / \delta n(\mathbf{r})$$

- ▶ The potential $v_{\text{xc}}(\mathbf{r})$ can be decomposed as

$$v_{\text{xc}}(\mathbf{r}) = v_{\text{x}}(\mathbf{r}) + v_{\text{c}}(\mathbf{r})$$

with the **exchange potential** $v_{\text{x}}(\mathbf{r}) = \delta E_{\text{x}}[n] / \delta n(\mathbf{r})$

and the **correlation potential** $v_{\text{c}}(\mathbf{r}) = \delta E_{\text{c}}[n] / \delta n(\mathbf{r})$

- ▶ We consider a **basis of K atom-centered functions** $\{\phi_\nu\}$, e.g. GTO basis functions. The orbitals are expanded as

$$\psi_i(\mathbf{r}) = \sum_{\nu}^K C_{\nu i} \phi_{\nu}(\mathbf{r})$$

- ▶ Inserting this expansion in the KS equations

$$\hat{h}_s \psi_i(\mathbf{r}) = \varepsilon_i \psi_i(\mathbf{r})$$

and multiplying on the left by $\phi_{\mu}^*(\mathbf{r})$ and integrating over $\mathbf{r}$, we arrive at the familiar **SCF generalized eigenvalue equation**

$$\sum_{\nu}^K F_{\mu\nu} C_{\nu i} = \varepsilon_i \sum_{\nu}^K S_{\mu\nu} C_{\nu i}$$

where $F_{\mu\nu} = \int \phi_{\mu}^*(\mathbf{r}) \hat{h}_s \phi_{\nu}(\mathbf{r}) d\mathbf{r}$ are the elements of the KS Fock matrix and $S_{\mu\nu} = \int \phi_{\mu}^*(\mathbf{r}) \phi_{\nu}(\mathbf{r}) d\mathbf{r}$ are the elements of the overlap matrix.

Practical calculations in an atomic basis (2/3)

► The Fock matrix is calculated as $F_{\mu\nu} = H_{\mu\nu}^{\text{core}} + J_{\mu\nu} + V_{\text{xc},\mu\nu}$

► $H_{\mu\nu}^{\text{core}}$ are the one-electron integrals: $H_{\mu\nu}^{\text{core}} = \int \phi_{\mu}^*(\mathbf{r}) \left(-\frac{1}{2} \nabla^2 + v_{\text{ne}}(\mathbf{r}) \right) \phi_{\nu}(\mathbf{r}) d\mathbf{r}$

► $J_{\mu\nu}$ is the Hartree potential matrix:

$$J_{\mu\nu} = \int \phi_{\mu}^*(\mathbf{r}) v_{\text{H}}(\mathbf{r}) \phi_{\nu}(\mathbf{r}) d\mathbf{r} = \sum_{\lambda}^K \sum_{\gamma}^K P_{\lambda\gamma} \langle \mu\gamma | \nu\lambda \rangle$$

where $\langle \mu\gamma | \nu\lambda \rangle = \iint_N \frac{\phi_{\mu}^*(\mathbf{r}_1) \phi_{\gamma}^*(\mathbf{r}_2) \phi_{\nu}(\mathbf{r}_1) \phi_{\lambda}(\mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2$ are the two-electron integrals and $P_{\lambda\gamma} = \sum_a C_{\lambda a} C_{\gamma a}^*$ is the density matrix.

► $V_{\text{xc},\mu\nu}$ is the exchange-correlation potential matrix: $V_{\text{xc},\mu\nu} = \int \phi_{\mu}^*(\mathbf{r}) v_{\text{xc}}(\mathbf{r}) \phi_{\nu}(\mathbf{r}) d\mathbf{r}$

► The total electronic energy is calculated as

$$E = \sum_{\mu}^K \sum_{\nu}^K P_{\nu\mu} H_{\mu\nu}^{\text{core}} + \frac{1}{2} \sum_{\mu}^K \sum_{\nu}^K P_{\nu\mu} J_{\mu\nu} + E_{\text{xc}}$$

► The density is calculated as $n(\mathbf{r}) = \sum_{\lambda}^K \sum_{\gamma}^K P_{\lambda\gamma} \phi_{\lambda}(\mathbf{r}) \phi_{\gamma}^*(\mathbf{r})$

- ▶ In the simplest approximation, the exchange-correlation energy functional has a local form

$$E_{xc}^{\text{local}} = \int f(n(\mathbf{r})) d\mathbf{r}$$

where $f(n(\mathbf{r}))$ has a complicated nonlinear dependence on the density $n(\mathbf{r})$.

- ▶ For example, in the local-density approximation (LDA), the exchange energy is

$$E_x^{\text{LDA}} = C_x \int n(\mathbf{r})^{4/3} d\mathbf{r}$$

where C_x is a constant, and the exchange potential is

$$v_x^{\text{LDA}}(\mathbf{r}) = \frac{4}{3} C_x n(\mathbf{r})^{1/3}$$

- ▶ Therefore, the integrals cannot be calculated analytically, but are instead evaluated by **numerical integration on a grid**

$$V_{xc,\mu\nu} \approx \sum_k w_k \phi_\mu^*(\mathbf{r}_k) v_{xc}(\mathbf{r}_k) \phi_\nu(\mathbf{r}_k)$$

and

$$E_{xc}^{\text{local}} \approx \sum_k w_k f(n(\mathbf{r}_k))$$

where $\mathbf{r}_k$ and w_k are quadrature points and weights.

Extension to spin density-functional theory (1/2)

- For dealing with an external magnetic field, DFT has been extended from the total density to **spin-resolved densities**

$$n_{\sigma}(\mathbf{r}) = N \int \cdots \int |\Psi(\mathbf{r}\sigma, \mathbf{x}_2, \dots, \mathbf{x}_N)|^2 d\mathbf{x}_2 \dots d\mathbf{x}_N \quad \text{with } \sigma \in \{\uparrow, \downarrow\}$$

which integrate to the numbers of σ -spin electrons, i.e. $\int n_{\sigma}(\mathbf{r}) d\mathbf{r} = N_{\sigma}$.

- Without magnetic fields, this is **in principle not necessary**. In practice, the dependence on the spin densities allows one to construct **more accurate approximate exchange-correlation functionals** for **open-shell systems**.
- The **universal density functional** is now defined as

$$F[n_{\uparrow}, n_{\downarrow}] = \min_{\Psi \rightarrow n_{\uparrow}, n_{\downarrow}} \langle \Psi | \hat{T} + \hat{W}_{\text{ee}} | \Psi \rangle$$

where the search is over wave functions Ψ giving the fixed spin densities $n_{\uparrow}$ and $n_{\downarrow}$.

- A **KS method** is obtained by decomposing $F[n_{\uparrow}, n_{\downarrow}]$ as

$$F[n_{\uparrow}, n_{\downarrow}] = T_{\text{s}}[n_{\uparrow}, n_{\downarrow}] + E_{\text{H}}[n] + E_{\text{xc}}[n_{\uparrow}, n_{\downarrow}]$$

where $T_{\text{s}}[n_{\uparrow}, n_{\downarrow}]$ is defined with a constrained search over (spin-unrestricted) Slater determinants Φ

$$T_{\text{s}}[n_{\uparrow}, n_{\downarrow}] = \min_{\Phi \rightarrow n_{\uparrow}, n_{\downarrow}} \langle \Phi | \hat{T} | \Phi \rangle$$

Extension to spin density-functional theory (2/2)

- ▶ The **exact ground-state energy** is expressed as

$$E_0 = \min_{\Phi} \left\{ \langle \Phi | \hat{T} + \hat{V}_{\text{ne}} | \Phi \rangle + E_{\text{H}}[n_{\Phi}] + E_{\text{xc}}[n_{\uparrow, \Phi}, n_{\downarrow, \Phi}] \right\}$$

- ▶ Writing the spatial orbitals of the determinant as $\psi_{a\sigma}(\mathbf{r})$ (with indices explicitly including spin now), we have now the **spin-dependent KS equations**

$$\left(-\frac{1}{2} \nabla^2 + v_{\text{ne}}(\mathbf{r}) + v_{\text{H}}(\mathbf{r}) + v_{\text{xc}, \sigma}(\mathbf{r}) \right) \psi_{a\sigma}(\mathbf{r}) = \varepsilon_{a\sigma} \psi_{a\sigma}(\mathbf{r})$$

with the spin-dependent exchange-correlation potential and density

$$v_{\text{xc}, \sigma}(\mathbf{r}) = \frac{\delta E_{\text{xc}}[n_{\uparrow}, n_{\downarrow}]}{\delta n_{\sigma}(\mathbf{r})} \quad \text{and} \quad n_{\sigma}(\mathbf{r}) = \sum_a^{N_{\sigma}} |\psi_{a\sigma}(\mathbf{r})|^2$$

- ▶ The spin-dependent exchange functional $E_{\text{x}}[n_{\uparrow}, n_{\downarrow}]$ can be obtained from the spin-independent exchange functional $E_{\text{x}}[n]$ with the **spin-scaling relation**

$$E_{\text{x}}[n_{\uparrow}, n_{\downarrow}] = \frac{1}{2} (E_{\text{x}}[2n_{\uparrow}] + E_{\text{x}}[2n_{\downarrow}])$$

Therefore, any approximation for the spin-independent exchange functional $E_{\text{x}}[n]$ can be easily extended to an approximation for the spin-dependent exchange functional $E_{\text{x}}[n_{\uparrow}, n_{\downarrow}]$. Unfortunately, there is no such relation for the correlation functional.

1 Basic density-functional theory

- Quantum many-electron problem
- Universal density functional
 - The Hohenberg-Kohn theorem
 - Levy-Lieb constrained-search formulation
- Kohn-Sham method
 - Decomposition of the universal functional
 - The Kohn-Sham equations
 - Practical calculations in an atomic basis
 - Extension to spin density-functional theory
- Generalized Kohn-Sham method

Generalized Kohn-Sham method

- An important extension of the KS method is the **generalized Kohn-Sham** (GKS) method (1996) in which the universal density functional $F[n]$ is decomposed as

$$F[n] = \min_{\Phi \rightarrow n} \left\{ \langle \Phi | \hat{T} | \Phi \rangle + E_H[n_\Phi] + S[\Phi] \right\} + \bar{S}[n]$$

where $S[\Phi]$ is any (reasonable) functional of a single-determinant wave function Φ and $\bar{S}[n]$ is the complementary density functional. E.g., in hybrids, $S[\Phi] = aE_x^{\text{HF}}[\Phi]$.

- Defining the **GKS exchange-correlation functional**, $E_{\text{xc}}^S[\Phi] = S[\Phi] + \bar{S}[n_\Phi]$, we can express the **exact ground-state energy** as

$$E_0 = \min_{\Phi} \left\{ \langle \Phi | \hat{T} + \hat{V}_{\text{ne}} | \Phi \rangle + E_H[n_\Phi] + E_{\text{xc}}^S[\Phi] \right\}$$

and any minimizing single-determinant wave function gives a ground-state density $n_0(\mathbf{r})$.

- The corresponding **GKS equations** are

$$\left(-\frac{1}{2}\nabla^2 + v_{\text{ne}}(\mathbf{r}) + v_{\text{H}}(\mathbf{r}) + v_{\bar{S}}(\mathbf{r}) \right) \psi_{a\sigma}(\mathbf{r}) + \frac{\delta S[\Phi]}{\delta \psi_{a\sigma}^*(\mathbf{r})} = \varepsilon_{a\sigma} \psi_{a\sigma}(\mathbf{r})$$

where $v_{\bar{S}}(\mathbf{r}) = \delta \bar{S}[n]/\delta n(\mathbf{r})$ is a local potential and $\delta S[\Phi]/\delta \psi_{a\sigma}^*(\mathbf{r})$ generates a one-electron (possibly nonlocal) operator.

- The GKS method gives **much more freedom** than the KS method (which corresponds to the special case $S[\Phi] = 0$).

- 1 Basic density-functional theory
 - Quantum many-electron problem
 - Universal density functional
 - Kohn-Sham method
 - Generalized Kohn-Sham method
- 2 Exact constraints for the exchange-correlation functional
 - Exact expressions for the exchange and correlation functionals
 - Uniform coordinate scaling
 - One-orbital spatial regions and self-interaction
 - Lieb-Oxford lower bound
 - Frontier orbital energies
- 3 Approximations for the exchange-correlation energy
 - Local-density approximation
 - Semilocal approximations
 - Single-determinant hybrid approximations
 - Multideterminant hybrid approximations
 - Dispersion corrections
 - Machine-learned density functionals
 - Benchmark
- 4 Time-dependent density-functional theory
 - Runge-Gross theorem
 - Linear-response TDDFT

2 Exact constraints for the exchange-correlation functional

- Exact expressions for the exchange and correlation functionals
 - The exchange and correlation holes
 - The adiabatic connection
- Uniform coordinate scaling
- One-orbital spatial regions and self-interaction
- Lieb-Oxford lower bound
- Frontier orbital energies
 - HOMO and LUMO energies
 - Fundamental gap

The exchange-correlation hole

- ▶ The **pair density** associated with the wave function $\Psi[n]$ is

$$n_2(\mathbf{r}_1, \mathbf{r}_2) = N(N-1) \int \cdots \int |\Psi[n](\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N)|^2 d\sigma_1 d\sigma_2 d\mathbf{x}_3 \dots d\mathbf{x}_N$$

which is a functional of the density. It is normalized to the number of electron pairs: $\iint n_2(\mathbf{r}_1, \mathbf{r}_2) d\mathbf{r}_1 d\mathbf{r}_2 = N(N-1)$. It is proportional to the probability density of finding two electrons at positions $(\mathbf{r}_1, \mathbf{r}_2)$ with all the other electrons anywhere.

- ▶ It can be used to express the electron-electron interaction energy

$$\langle \Psi[n] | \hat{W}_{ee} | \Psi[n] \rangle = \frac{1}{2} \iint \frac{n_2(\mathbf{r}_1, \mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2$$

- ▶ Mirroring the decomposition of $E_{Hxc}[n]$, the pair density can be decomposed as

$$n_2(\mathbf{r}_1, \mathbf{r}_2) = \underbrace{n(\mathbf{r}_1)n(\mathbf{r}_2)}_{\text{independent electrons}} + \underbrace{n_{2,xc}(\mathbf{r}_1, \mathbf{r}_2)}_{\text{exchange and correlation effects}}$$

- ▶ We also introduce the **exchange-correlation hole** $n_{xc}(\mathbf{r}_1, \mathbf{r}_2)$ by

$$n_{2,xc}(\mathbf{r}_1, \mathbf{r}_2) = n(\mathbf{r}_1)n_{xc}(\mathbf{r}_1, \mathbf{r}_2)$$

It can be interpreted as the modification due to exchange and correlation effects of the conditional probability of finding an electron at $\mathbf{r}_2$ knowing that one has been found at $\mathbf{r}_1$.

- ▶ We have the exact constraints: $n_{xc}(\mathbf{r}_1, \mathbf{r}_2) \geq -n(\mathbf{r}_2)$ and $\int n_{xc}(\mathbf{r}_1, \mathbf{r}_2) d\mathbf{r}_2 = -1$

The exchange hole

- Similarly, we define the **KS pair density** associated with the KS single determinant $\Phi[n]$

$$n_{2,\text{KS}}(\mathbf{r}_1, \mathbf{r}_2) = N(N-1) \int \cdots \int |\Phi[n](\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N)|^2 d\sigma_1 d\sigma_2 d\mathbf{x}_3 \dots d\mathbf{x}_N$$

- It can be decomposed as


$$n_{2,\text{KS}}(\mathbf{r}_1, \mathbf{r}_2) = n(\mathbf{r}_1)n(\mathbf{r}_2) + n_{2,x}(\mathbf{r}_1, \mathbf{r}_2)$$

and we introduce the **exchange hole** $n_x(\mathbf{r}_1, \mathbf{r}_2)$ by

$$n_{2,x}(\mathbf{r}_1, \mathbf{r}_2) = n(\mathbf{r}_1)n_x(\mathbf{r}_1, \mathbf{r}_2)$$

which satisfies the exact constraints:

$$n_x(\mathbf{r}_1, \mathbf{r}_2) \geq -n(\mathbf{r}_2) \quad \text{and} \quad \int n_x(\mathbf{r}_1, \mathbf{r}_2) d\mathbf{r}_2 = -1 \quad \text{and} \quad n_x(\mathbf{r}_1, \mathbf{r}_2) \leq 0$$



- The exchange energy functional is the electrostatic interaction energy between an electron and its exchange hole:

$$E_x[n] = \frac{1}{2} \iint \frac{n(\mathbf{r}_1)n_x(\mathbf{r}_1, \mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2 = \int n(\mathbf{r}_1)\varepsilon_x[n](\mathbf{r}_1) d\mathbf{r}_1$$

where $\varepsilon_x[n](\mathbf{r}_1)$ is the exchange energy per particle. In approximate exchange density functionals, the quantity $\varepsilon_x[n](\mathbf{r}_1)$ is usually what is approximated.

The correlation hole

- ▶ The **correlation hole** is defined as the difference

$$n_c(\mathbf{r}_1, \mathbf{r}_2) = n_{xc}(\mathbf{r}_1, \mathbf{r}_2) - n_x(\mathbf{r}_1, \mathbf{r}_2)$$

and satisfies the sum rule


$$\int n_c(\mathbf{r}_1, \mathbf{r}_2) d\mathbf{r}_2 = 0$$

which implies that the correlation hole has negative and positive contributions.

- ▶ The potential contribution to the correlation energy can be written in terms of the correlation hole

$$U_c[n] = \frac{1}{2} \iint \frac{n(\mathbf{r}_1) n_c(\mathbf{r}_1, \mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2$$

But in order to express the total correlation energy $E_c[n] = T_c[n] + U_c[n]$ in a similar form, we need to introduce the adiabatic-connection formalism.



The adiabatic connection (1/3)

- ▶ The idea of the **adiabatic connection** is to have a continuous path between the non-interacting KS system and the physical system while keeping the ground-state density constant.
- ▶ For this, we introduce a Hamiltonian depending on a **coupling constant** λ which switches on the electron-electron interaction

$$\hat{H}^\lambda = \hat{T} + \lambda \hat{W}_{ee} + \hat{V}^\lambda$$

where $\hat{V}^\lambda$ is the external local potential imposing that the ground-state density is the same as the ground-state density of the physical system for all λ , i.e. $n^\lambda(\mathbf{r}) = n_0(\mathbf{r}), \forall \lambda$.

- ▶ By varying λ , we connect the KS non-interacting system ($\lambda = 0$) to the physical interacting system ($\lambda = 1$):

$$\underbrace{\hat{H}^{\lambda=0}}_{\text{KS non-interacting system}} \longleftrightarrow^{0 \leq \lambda \leq 1} \underbrace{\hat{H}^{\lambda=1}}_{\text{Physical interacting system}}$$

- ▶ We define a universal functional for each value of the parameter λ

$$F^\lambda[n] = \min_{\Psi \rightarrow n} \langle \Psi | \hat{T} + \lambda \hat{W}_{ee} | \Psi \rangle = \langle \Psi^\lambda[n] | \hat{T} + \lambda \hat{W}_{ee} | \Psi^\lambda[n] \rangle$$

The adiabatic connection (2/3)

- ▶ The functional $F^\lambda[n]$ can be decomposed as

$$F^\lambda[n] = T_s[n] + E_H^\lambda[n] + E_x^\lambda[n] + E_c^\lambda[n]$$

- ▶ $E_H^\lambda[n]$ and $E_x^\lambda[n]$ are the Hartree and exchange functionals associated with the interaction $\lambda \hat{W}_{ee}$ and are simply linear in λ

$$E_H^\lambda[n] = \lambda E_H[n]$$

and

$$E_x^\lambda[n] = \lambda E_x[n]$$

- ▶ The correlation functional $E_c^\lambda[n]$ is nonlinear in λ

$$E_c^\lambda[n] = \langle \Psi^\lambda[n] | \hat{T} + \lambda \hat{W}_{ee} | \Psi^\lambda[n] \rangle - \langle \Phi[n] | \hat{T} + \lambda \hat{W}_{ee} | \Phi[n] \rangle$$

- ▶ We can get rid of $\hat{T}$ by taking the derivative with respect to λ and using the Hellmann-Feynman theorem for the wave function $\Psi^\lambda[n]$

$$\frac{\partial E_c^\lambda[n]}{\partial \lambda} = \langle \Psi^\lambda[n] | \hat{W}_{ee} | \Psi^\lambda[n] \rangle - \langle \Phi[n] | \hat{W}_{ee} | \Phi[n] \rangle$$

- ▶ Reintegrating over λ from 0 to 1, and using $E_c^{\lambda=1}[n] = E_c[n]$ and $E_c^{\lambda=0}[n] = 0$ (assuming no degeneracies at $\lambda = 0$), we arrive at the **adiabatic-connection formula**

$$E_c[n] = \int_0^1 d\lambda \langle \Psi^\lambda[n] | \hat{W}_{ee} | \Psi^\lambda[n] \rangle - \langle \Phi[n] | \hat{W}_{ee} | \Phi[n] \rangle$$

- ▶ Introducing the correlation hole $n_c^\lambda(\mathbf{r}_1, \mathbf{r}_2)$ associated with the wave function $\Psi^\lambda[n]$, the adiabatic-connection formula can also be written as

$$E_c[n] = \frac{1}{2} \int_0^1 d\lambda \iint \frac{n(\mathbf{r}_1) n_c^\lambda(\mathbf{r}_1, \mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2$$

- ▶ Introducing the λ -integrated correlation hole $\bar{n}_c(\mathbf{r}_1, \mathbf{r}_2) = \int_0^1 d\lambda n_c^\lambda(\mathbf{r}_1, \mathbf{r}_2)$, we finally write

$$E_c[n] = \frac{1}{2} \iint \frac{n(\mathbf{r}_1) \bar{n}_c(\mathbf{r}_1, \mathbf{r}_2)}{|\mathbf{r}_1 - \mathbf{r}_2|} d\mathbf{r}_1 d\mathbf{r}_2 = \int n(\mathbf{r}_1) \varepsilon_c[n](\mathbf{r}_1) d\mathbf{r}_1$$

where $\varepsilon_c[n](\mathbf{r}_1)$ is the correlation energy per particle, which is the quantity usually approximated in practice.

2 Exact constraints for the exchange-correlation functional

- Exact expressions for the exchange and correlation functionals
 - The exchange and correlation holes
 - The adiabatic connection
- Uniform coordinate scaling
- One-orbital spatial regions and self-interaction
- Lieb-Oxford lower bound
- Frontier orbital energies
 - HOMO and LUMO energies
 - Fundamental gap

- For a given density n , we consider the **scaled density** with a scaling factor $\gamma > 0$

$$n_\gamma(\mathbf{r}) = \gamma^3 n(\gamma\mathbf{r})$$

which preserves the number of electrons, i.e. $\int n_\gamma(\mathbf{r})d\mathbf{r} = \int n(\mathbf{r})d\mathbf{r} = N$.

- It can be shown that the Hartree and exchange density functionals scale linearly in γ

$$E_H[n_\gamma] = \gamma E_H[n] \quad \text{and} \quad E_x[n_\gamma] = \gamma E_x[n]$$

- The correlation density functional has the more complicated scaling

$$E_c[n_\gamma] = \gamma^2 E_c^{1/\gamma}[n]$$

where $E_c^{1/\gamma}[n]$ is the correlation density functional for coupling constant $\lambda = 1/\gamma$.

- ▶ In the **high-density limit** ($\gamma \rightarrow \infty$), the correlation functional goes to a constant, for nondegenerate KS systems,

$$\lim_{\gamma \rightarrow \infty} E_c[n_\gamma] = E_c^{\text{GL2}}[n]$$

where $E_c^{\text{GL2}}[n]$ is the second-order Görling-Levy (GL2) correlation energy.

- ▶ This is also called the **weak-correlation limit** since $E_c[n] \ll E_x[n]$.
 - ▶ Atomic and molecular systems are often close to the high-density limit. E.g., for the ground-state density of He, $E_c[n] = -0.0421$ a.u. and $\lim_{\gamma \rightarrow \infty} E_c[n_\gamma] = -0.0467$ a.u..
- ▶ In the **low-density limit** ($\gamma \rightarrow 0$), the Hartree-exchange-correlation functional goes to zero linearly in γ

$$E_{\text{Hxc}}[n_\gamma] \underset{\gamma \rightarrow 0}{\sim} \gamma W_{\text{ee}}^{\text{SCE}}[n]$$

where $W_{\text{ee}}^{\text{SCE}}[n] = \min_{\Psi \rightarrow n} \langle \Psi | \hat{W}_{\text{ee}} | \Psi \rangle$ is the strictly-correlated-electron (SCE) functional.

- ▶ This limit corresponds to a Wigner crystallization.
- ▶ This is also called the **strong-correlation limit** because $E_c[n] \sim E_x[n]$.

2 Exact constraints for the exchange-correlation functional

- Exact expressions for the exchange and correlation functionals
 - The exchange and correlation holes
 - The adiabatic connection
- Uniform coordinate scaling
- One-orbital spatial regions and self-interaction
- Lieb-Oxford lower bound
- Frontier orbital energies
 - HOMO and LUMO energies
 - Fundamental gap

- ▶ For **one-electron densities** $n_{1e}(\mathbf{r}) = |\psi_1(\mathbf{r})|^2$ where ψ_1 is the unique occupied KS orbital, we have

$$E_x[n_{1e}] = -E_H[n_{1e}] \quad \text{and} \quad E_c[n_{1e}] = 0$$

- ▶ For **opposite-spin two-electron densities** $n_{2e}^{\uparrow\downarrow}(\mathbf{r}) = 2|\psi_1(\mathbf{r})|^2$ where ψ_1 is the unique doubly occupied KS orbital, we have

$$E_x[n_{2e}^{\uparrow\downarrow}] = -\frac{1}{2}E_H[n_{2e}^{\uparrow\downarrow}]$$

- ▶ For systems with more electrons, **similar relations apply locally in one-orbital spatial regions**, i.e. in regions where only one occupied KS orbital is not zero. This situation can be approximately realized in chemical systems (unpaired electron in a radical, and electron pair in a single covalent bond, in a lone pair, or in a core orbital).
- ▶ If approximate exchange and correlation density functionals do not satisfy these constraints, we say that they introduce a **self-interaction error**.

2 Exact constraints for the exchange-correlation functional

- Exact expressions for the exchange and correlation functionals
 - The exchange and correlation holes
 - The adiabatic connection
- Uniform coordinate scaling
- One-orbital spatial regions and self-interaction
- **Lieb-Oxford lower bound**
- Frontier orbital energies
 - HOMO and LUMO energies
 - Fundamental gap

- ▶ Lieb and Oxford derived a useful **lower bound** which can be expressed as

$$E_x[n] \geq E_{xc}[n] \geq -C_{LO} \int n(\mathbf{r})^{4/3} d\mathbf{r}$$

where the optimal (i.e., smallest) constant C_{LO} (independent of the electron number N) has been narrowed to $1.4442 \leq C_{LO} \leq 1.5765$.

- ▶ This bound is approached in the low-density limit.
- ▶ For one-electron densities and opposite-spin two-electron densities, specific tighter bounds (i.e., with smaller C_{LO}) are known.

2 Exact constraints for the exchange-correlation functional

- Exact expressions for the exchange and correlation functionals
 - The exchange and correlation holes
 - The adiabatic connection
- Uniform coordinate scaling
- One-orbital spatial regions and self-interaction
- Lieb-Oxford lower bound
- **Frontier orbital energies**
 - HOMO and LUMO energies
 - Fundamental gap

The HOMO energy and the ionization energy

- ▶ For clarity, we will explicitly indicate the dependence on the electron number N in this section.
- ▶ For finite systems, the exact ground-state density of a N -electron system decays exponentially for $r = |\mathbf{r}| \rightarrow \infty$ with an exponent related to the **ionization energy**
 $I_N = E_0^{N-1} - E_0^N$

$$n^N(\mathbf{r}) \underset{r \rightarrow \infty}{\propto} e^{-2\sqrt{2I_N} r}$$

- ▶ Choosing the constant in the KS potential so that it goes to zero at infinity, i.e. $\lim_{|r| \rightarrow \infty} v_s^N(\mathbf{r}) = 0$, it can be shown the density calculated from the KS orbitals decays exponentially with an exponent related to the **HOMO energy** ε_H^N

$$n^N(\mathbf{r}) = \sum_a^N |\psi_a(\mathbf{r})|^2 \underset{r \rightarrow \infty}{\propto} e^{-2\sqrt{-2\varepsilon_H^N} r}$$

- ▶ This implies that the **KS HOMO energy is the opposite of the exact ionization energy**

$$\varepsilon_H^N = -I_N$$

- ▶ It is similar to Koopmans' theorem for HF, except that here it is exact (no neglect of correlation or orbital relaxation).

The LUMO energy, the electron affinity, the derivative discontinuity

- Contrary to what one could have expected, the **KS LUMO energy** ϵ_L^N is **not the opposite of the exact electron affinity** $A_N = E_0^N - E_0^{N+1}$ but instead

$$\epsilon_L^N = -A_N - \Delta_{xc}^N$$

where $\Delta_{xc}^N \geq 0$ is a constant.

- For the $(N+1)$ -electron system (with the same external potential v_{ne}), we have

$$\epsilon_H^{N+1} = -I_{N+1} = -A_N, \text{ so it means that}$$

$$\Delta_{xc}^N = \epsilon_H^{N+1} - \epsilon_L^N$$


i.e., the constant Δ_{xc}^N corresponds to the **“jump” of the LUMO energy** of the N -electron system upon adding an electron so that the HOMO energy of the $(N+1)$ -electron system correctly gives $-I_{N+1}$.

- In the extension of DFT to fractional electron numbers, it can be shown that the constant Δ_{xc}^N corresponds to the **uniform jump** that the **exchange-correlation potential** makes when going from $N - \delta$ electrons to $N + \delta$ electrons with $\delta \rightarrow 0^+$

$$\Delta_{xc}^N = v_{xc}^{N+\delta}(\mathbf{r}) - v_{xc}^{N-\delta}(\mathbf{r}) = \left(\frac{\delta E_{xc}[n]}{\delta n(\mathbf{r})} \right)_{N+\delta} - \left(\frac{\delta E_{xc}[n]}{\delta n(\mathbf{r})} \right)_{N-\delta}$$

i.e. Δ_{xc}^N is the **derivative discontinuity** in the exchange-correlation energy functional $E_{xc}[n]$.

Kohn-Sham frontier orbital energies: Graphical summary



- ▶ The **fundamental gap** of the N -electron system is defined as

$$E_{\text{gap}}^N = I_N - A_N$$

- ▶ In KS DFT, it can thus be expressed as

$$E_{\text{gap}}^N = \underbrace{\varepsilon_{\text{L}}^N - \varepsilon_{\text{H}}^N}_{\text{KS gap}} + \Delta_{\text{xc}}^N$$

So the **KS gap is not equal to the exact fundamental gap** of the system, the difference coming from the derivative discontinuity Δ_{xc}^N .

- ▶ The derivative discontinuity Δ_{xc}^N can represent an important contribution to the fundamental gap. In the special case of open-shell systems, we have $\varepsilon_{\text{L}}^N = \varepsilon_{\text{H}}^N$, and thus if the fundamental gap of an open-shell system is not zero (Mott insulator), it is entirely given by Δ_{xc}^N .

- 1 Basic density-functional theory
 - Quantum many-electron problem
 - Universal density functional
 - Kohn-Sham method
 - Generalized Kohn-Sham method
- 2 Exact constraints for the exchange-correlation functional
 - Exact expressions for the exchange and correlation functionals
 - Uniform coordinate scaling
 - One-orbital spatial regions and self-interaction
 - Lieb-Oxford lower bound
 - Frontier orbital energies
- 3 Approximations for the exchange-correlation energy
 - Local-density approximation
 - Semilocal approximations
 - Single-determinant hybrid approximations
 - Multideterminant hybrid approximations
 - Dispersion corrections
 - Machine-learned density functionals
 - Benchmark
- 4 Time-dependent density-functional theory
 - Runge-Gross theorem
 - Linear-response TDDFT

3 Approximations for the exchange-correlation energy

- Local-density approximation
- Semilocal approximations
 - Generalized-gradient approximations
 - Meta-generalized-gradient approximations
- Single-determinant hybrid approximations
 - Hybrid approximations
 - Range-separated hybrid approximations
- Multideterminant hybrid approximations
 - Double-hybrid approximations
 - Range-separated double-hybrid approximations
- Dispersion corrections
 - Semiempirical dispersion corrections
 - Nonlocal van der Waals density functionals
- Machine-learned density functionals
- Benchmark

Local-density approximation

- ▶ In the **local-density approximation** (LDA), introduced by Kohn and Sham (1965), the exchange-correlation functional is approximated as

$$E_{xc}^{LDA}[n] = \int n(\mathbf{r}) \varepsilon_{xc}^{UEG}(n(\mathbf{r})) d\mathbf{r}$$

where $\varepsilon_{xc}^{UEG}(n)$ is the exchange-correlation energy per particle of the infinite **uniform electron gas (UEG)** with the density n .

- ▶ The exchange energy per particle of the UEG can be calculated analytically

$$\varepsilon_x^{UEG}(n) = C_x n^{1/3} \quad \text{Dirac (1930) and Slater (1951)}$$

- ▶ For the correlation energy per particle $\varepsilon_c^{UEG}(n)$ of the UEG, there are some parametrized functions of n fitted to QMC data and imposing the high- and low-density expansions (using the Wigner-Seitz radius $r_s = (3/(4\pi n))^{1/3}$)

$$\varepsilon_c^{UEG} \underset{r_s \rightarrow 0}{=} A \ln r_s + B + C r_s \ln r_s + O(r_s)$$

high-density limit or
weak-correlation limit

$$\varepsilon_c^{UEG} \underset{r_s \rightarrow \infty}{=} \frac{a}{r_s} + \frac{b}{r_s^{3/2}} + \frac{c}{r_s^2} + O\left(\frac{1}{r_s^{5/2}}\right)$$

low-density limit or
strong-correlation limit

The two most used parametrizations are VWN and PW92. Generalization to spin densities $\varepsilon_c^{UEG}(n_{\uparrow}, n_{\downarrow})$ is sometimes referred to as local-spin-density (LSD) approximation.

3 Approximations for the exchange-correlation energy

- Local-density approximation
- Semilocal approximations
 - Generalized-gradient approximations
 - Meta-generalized-gradient approximations
- Single-determinant hybrid approximations
 - Hybrid approximations
 - Range-separated hybrid approximations
- Multideterminant hybrid approximations
 - Double-hybrid approximations
 - Range-separated double-hybrid approximations
- Dispersion corrections
 - Semiempirical dispersion corrections
 - Nonlocal van der Waals density functionals
- Machine-learned density functionals
- Benchmark

Generalized-gradient approximations

- ▶ The next logical step beyond the LDA is to add the dependence on the gradient of the density $\nabla n(\mathbf{r})$.
- ▶ From the 1980s, many **generalized-gradient approximations (GGAs)** have been developed with the generic form

$$E_{xc}^{GGA}[n] = \int f(n(\mathbf{r}), \nabla n(\mathbf{r})) d\mathbf{r}$$

- ▶ The function f is chosen so as to satisfy some exact constraints and often contains some parameters fitted to experimental or theoretical data.
- ▶ The GGAs provide a big improvement over LDA for molecular systems.
- ▶ The GGAs are often called **semilocal** approximations, which means that they involve a single integral on $\mathbf{r}$ using “semilocal information” through $\nabla n(\mathbf{r})$.
- ▶ For simplicity, we consider here only the spin-independent form, but in practice GGA functionals are more generally formulated in terms of spin densities and their gradients

$$E_{xc}^{GGA}[n_{\uparrow}, n_{\downarrow}] = \int f(n_{\uparrow}(\mathbf{r}), n_{\downarrow}(\mathbf{r}), \nabla n_{\uparrow}(\mathbf{r}), \nabla n_{\downarrow}(\mathbf{r})) d\mathbf{r}$$

- ▶ Examples of GGAs: B exchange functional (1988), LYP correlation functional (1988), PBE exchange-correlation functional (1996)

- ▶ The **meta-generalized-gradient approximations (mGGAs)** are of the generic form

$$E_{xc}^{mGGA}[n, \tau] = \int f(n(\mathbf{r}), \nabla n(\mathbf{r}), \nabla^2 n(\mathbf{r}), \tau(\mathbf{r})) d\mathbf{r}$$

where $\nabla^2 n(\mathbf{r})$ is the Laplacian of the density and $\tau(\mathbf{r})$ is the **non-interacting positive kinetic energy density**

$$\tau(\mathbf{r}) = \frac{1}{2} \sum_a^N |\nabla \psi_a(\mathbf{r})|^2$$

which, as we will see, contains useful information.

- ▶ A mGGA can be either considered as an implicit functional of the density in the KS method, i.e. $E_{xc}[n] = E_{xc}^{mGGA}[n, \tau_{\Phi}[n]]$, or more commonly as an explicit functional of a single-determinant Φ in the GKS method, i.e. $E_{xc}^S[\Phi] = E_{xc}^{mGGA}[n_{\Phi}, \tau_{\Phi}]$.
- ▶ In the GKS method, a mGGA functional generates a non-multiplicative potential. But don't worry, this is allowed in GKS!
- ▶ Nowadays, $\nabla^2 n(\mathbf{r})$ is rarely used to construct mGGAs because it contains similar information than $\tau(\mathbf{r})$.
- ▶ The mGGAs are considered as part of the family of semilocal approximations.
- ▶ The mGGAs provide a modest improvement over GGAs.

► Motivations for introducing the variable $\tau(\mathbf{r})$:

- Short-range expansion of the spherically average exchange hole (for closed-shell systems):

$$\tilde{n}_x(\mathbf{r}_1, r_{12}) = -\frac{n(\mathbf{r}_1)}{2} - \frac{1}{3} \left(\frac{1}{4} \nabla^2 n(\mathbf{r}_1) - 4\tau(\mathbf{r}_1) + \frac{|\nabla n(\mathbf{r}_1)|^2}{8n(\mathbf{r}_1)} \right) r_{12}^2 + O(r_{12}^4)$$

Thus $\tau(\mathbf{r})$ is needed to describe the curvature of the exchange hole.

- $\tau(\mathbf{r})$ can be used as an indicator of **one-orbital spatial regions** (regions containing one or two electrons in a single orbital).

This is done by comparing $\tau(\mathbf{r})$ with the **von Weizsäcker kinetic energy density**

$$\tau^W(\mathbf{r}) = \frac{|\nabla n(\mathbf{r})|^2}{8n(\mathbf{r})}$$

which is the exact $\tau(\mathbf{r})$ for one and two electrons in a single orbital.

- Examples of mGGAs: TPSS (2003), M06-L (2006), and SCAN (2015).

3 Approximations for the exchange-correlation energy

- Local-density approximation
- Semilocal approximations
 - Generalized-gradient approximations
 - Meta-generalized-gradient approximations
- **Single-determinant hybrid approximations**
 - Hybrid approximations
 - Range-separated hybrid approximations
- Multideterminant hybrid approximations
 - Double-hybrid approximations
 - Range-separated double-hybrid approximations
- Dispersion corrections
 - Semiempirical dispersion corrections
 - Nonlocal van der Waals density functionals
- Machine-learned density functionals
- Benchmark

- ▶ In 1993, Becke proposed to **mix Hartree-Fock (HF) exchange** with GGA functionals in a three-parameter hybrid (3H) approximation

$$E_{xc}^{3H}[\Phi] = a E_x^{HF}[\Phi] + b E_x^{GGA}[n_\Phi] + (1 - a - b) E_x^{LDA}[n_\Phi] + c E_c^{GGA}[n_\Phi] + (1 - c) E_c^{LDA}[n_\Phi]$$

where a , b , and c are empirical parameters. Example: B3LYP ($a = 0.20$)

- ▶ These hybrids are approximations within the **GKS method**. The term $S[\Phi] = a E_x^{HF}[\Phi]$ generates a **nonlocal HF exchange potential** $av_{x,\sigma}^{HF}(\mathbf{r}, \mathbf{r}')$. Again, this is perfectly allowed in the GKS method.
- ▶ Adding a fraction a of HF exchange decreases the **self-interaction error**, which tends to favor too much delocalized electron densities. However, a too large a tends to increase the **static-correlation error** (stretched chemical bonds, transition metal elements, ...).
- ▶ In 1996, Becke proposed a simpler **one-parameter hybrid (1H) approximation**

$$E_{xc}^{1H}[\Phi] = a E_x^{HF}[\Phi] + (1 - a) E_x^{DFA}[n_\Phi] + E_c^{DFA}[n_\Phi]$$

where E_x^{DFA} and E_c^{DFA} can be any semilocal density-functional approximations (DFAs).

- ▶ The optimal a is often around 0.25. Example: PBE0 = HF/PBE hybrid with $a = 0.25$.
- ▶ A strategy is to use flexible E_x^{DFA} and E_c^{DFA} in a hybrid approximation and optimize many parameters on molecular properties.
Example: B97 (13 parameters) and M06 and M06-2X (36 parameters).

Range-separated hybrid (RSH) approximations

- ▶ Based on ideas of Savin (1996), Hirao and coworkers (2001) proposed a **long-range correction (LC)** scheme

$$E_{xc}^{LC}[\Phi] = E_x^{lr,HF}[\Phi] + E_x^{sr,DFA}[n_\Phi] + E_c^{DFA}[n_\Phi]$$

where

- ▶ $E_x^{lr,HF}[\Phi]$ is the HF exchange energy for the **long-range electron-electron interaction** $\frac{\text{erf}(\mu r_{12})}{r_{12}}$ replacing the Coulomb interaction $\frac{1}{r_{12}}$,
- ▶ $E_x^{sr,DFA}[n]$ is a semilocal DFA exchange energy for the complement **short-range electron-electron interaction** (semilocal DFAs are more accurate if limited to short-range interactions),
- ▶ the range-separation parameter μ (also sometimes denoted as ω) is often taken as $\mu \approx 0.3 - 0.5 \text{ bohr}^{-1}$.

Example: LC- ω PBE

- ▶ In 2004, Yanai, Tew, and Handy introduced a more flexible decomposition called the **Coulomb-attenuating method (CAM)**

$$E_{xc}^{CAM}[\Phi] = a E_x^{sr,HF}[\Phi] + b E_x^{lr,HF}[\Phi] + (1 - a) E_x^{sr,DFA}[n_\Phi] + (1 - b) E_x^{lr,DFA}[n_\Phi] + E_c^{DFA}[n_\Phi]$$

Examples: CAM-B3LYP, ω B97X

3 Approximations for the exchange-correlation energy

- Local-density approximation
- Semilocal approximations
 - Generalized-gradient approximations
 - Meta-generalized-gradient approximations
- Single-determinant hybrid approximations
 - Hybrid approximations
 - Range-separated hybrid approximations
- **Multideterminant hybrid approximations**
 - Double-hybrid approximations
 - Range-separated double-hybrid approximations
- Dispersion corrections
 - Semiempirical dispersion corrections
 - Nonlocal van der Waals density functionals
- Machine-learned density functionals
- Benchmark

- ▶ In 2006, Grimme introduced a **two-parameter double-hybrid (2DH) approximation**

$$E_{xc}^{2DH} = a_x E_x^{HF}[\Phi] + (1 - a_x) E_x^{DFA}[n_\Phi] + (1 - a_c) E_c^{DFA}[n_\Phi] + a_c E_c^{MP2}$$

where the MP2-like correlation energy E_c^{MP2} is added a posteriori with the previously calculated orbitals. Example: B2-PLYP ($a_x = 0.53$ and $a_c = 0.27$).

- ▶ The presence of nonlocal MP2 correlation allows one to use a larger fraction of nonlocal HF exchange.
- ▶ In 2011, Sharkas, Toulouse, and Savin showed that double hybrids can be understood as approximations of a **multideterminant extension of the KS method** based on the adiabatic-connection formalism in which the exact ground-state energy is written as

$$E_0 = \min_{\Psi} \left\{ \langle \Psi | \hat{T} + \hat{V}_{ne} + \lambda \hat{W}_{ee} | \Psi \rangle + \bar{E}_{Hxc}^\lambda[n_\Psi] \right\}$$

where $\bar{E}_{Hxc}^\lambda[n] = (1 - \lambda)E_H[n] + (1 - \lambda)E_x[n] + \bar{E}_c^\lambda[n]$ and $\bar{E}_c^\lambda[n] = E_c[n] - \lambda^2 E_c[n_{1/\lambda}]$.

- ▶ At second order of a non-linear Møller-Plesset-like perturbation theory, and using $E_c[n_{1/\lambda}] \approx E_c[n]$, we obtain a **one-parameter double-hybrid (1DH) approximation**

$$E_{xc}^{1DH} = \lambda E_x^{HF}[\Phi] + (1 - \lambda) E_x^{DFA}[n_\Phi] + (1 - \lambda^2) E_c^{DFA}[n_\Phi] + \lambda^2 E_c^{MP2}$$

- ▶ The multideterminant extension of the KS method can also be used to rigorously combine wave-function methods such as MCSCF with DFT.

- ▶ In 1996, Savin introduced the **range-separated multideterminant extension of the KS scheme** in which the exact ground-state energy is written as

$$E_0 = \min_{\Psi} \left\{ \langle \Psi | \hat{T} + \hat{V}_{\text{ne}} + \hat{W}_{\text{ee}}^{\text{lr}} | \Psi \rangle + \bar{E}_{\text{Hxc}}^{\text{sr}}[n_{\Psi}] \right\}$$

where $\hat{W}_{\text{ee}}^{\text{lr}}$ is the long-range electron-electron operator for the pair potential $\text{erf}(\mu r_{12})/r_{12}$ and $\bar{E}_{\text{Hxc}}^{\text{sr}}[n]$ is the complementary short-range density functional.

- ▶ The approach can be used to rigorously combine any wave-function method with DFT.
- ▶ In 2005, Ángyán, Gerber, Savin, and Toulouse introduced a **range-separated double-hybrid (RSDH)** approximation (also called RSH+MP2)

$$E_{\text{xc}}^{\text{RSDH}} = E_{\text{x}}^{\text{lr,HF}}[\Phi] + E_{\text{x}}^{\text{sr,DFA}}[n_{\Phi}] + E_{\text{c}}^{\text{sr,DFA}}[n_{\Phi}] + E_{\text{c}}^{\text{lr,MP2}}$$

- ▶ Obtained as second order of a non-linear Møller-Plesset-like perturbation theory.
- ▶ Long-range MP2 is qualitatively correct for London dispersion interactions.
- ▶ Long-range MP2 has a fast convergence with the one-electron basis size.
- ▶ Extensions of this scheme to a more flexible CAM decomposition have also been proposed.

3 Approximations for the exchange-correlation energy

- Local-density approximation
- Semilocal approximations
 - Generalized-gradient approximations
 - Meta-generalized-gradient approximations
- Single-determinant hybrid approximations
 - Hybrid approximations
 - Range-separated hybrid approximations
- Multideterminant hybrid approximations
 - Double-hybrid approximations
 - Range-separated double-hybrid approximations
- Dispersion corrections
 - Semiempirical dispersion corrections
 - Nonlocal van der Waals density functionals
- Machine-learned density functionals
- Benchmark

- ▶ To explicitly account for London dispersion interactions, it has been proposed in the 2000s to add to the standard approximate functionals a **semiempirical dispersion correction** of the form

$$E_{\text{disp}} = -s \sum_{\alpha < \beta} f(R_{\alpha\beta}) \frac{C_6^{\alpha\beta}}{R_{\alpha\beta}^6}$$

where

- ▶ $R_{\alpha\beta}$ is the distance between a pair of atoms,
 - ▶ $C_6^{\alpha\beta}$ is the dispersion coefficient between these atoms,
 - ▶ $f(R_{\alpha\beta})$ is a damping function which tends to 1 at large $R_{\alpha\beta}$ and tends to 0 at small $R_{\alpha\beta}$,
 - ▶ s is a scaling parameter that can be adjusted for each approximate functional.
- ▶ The dispersion coefficients $C_6^{\alpha\beta}$ are empirically obtained from tabulated data.
 - ▶ The most recent versions also includes $C_8^{\alpha\beta}$ two-body terms and $C_9^{\alpha\beta\gamma}$ three-body terms.
 - ▶ This approach was named “DFT-D” by Grimme. Examples of DFT-D functionals: PBE-D, B97-D, B3LYP-D, ω B97X-D, B2PLYP-D.
 - ▶ There are also various proposals to make the determination of dispersion coefficients less empirical, e.g. Becke and Johnson (2007), Tkatchenko and Scheffler (2009), Sato and Nakai (2010).

- ▶ Another approach to describe dispersion interactions is to add to the standard approximate functionals a **nonlocal van der Waals density functional** of the form

$$E_c^{nl}[n] = \frac{1}{2} \iint n(\mathbf{r}_1)n(\mathbf{r}_2)\phi(\mathbf{r}_1,\mathbf{r}_2)d\mathbf{r}_1d\mathbf{r}_2$$

where $\phi(\mathbf{r}_1,\mathbf{r}_2)$ is a correlation kernel.

- ▶ Two main families of such nonlocal correlation functionals exist: the “van der Waals density functionals” (vdW-DF) of Langreth, Lundqvist and coworkers and the Vydrov-Van Voorhis (VV) functionals.
- ▶ For example, the VV10 nonlocal correlation functional (2010) uses a theory-based kernel $\phi(\mathbf{r}_1,\mathbf{r}_2)$ with two adjustable parameters.
- ▶ Nonlocal van der Waals density functionals are less empirical but more computationally expensive than semiempirical dispersion corrections.
- ▶ Examples of functionals using VV10: ω B97X-V and ω B97M-V.

3 Approximations for the exchange-correlation energy

- Local-density approximation
- Semilocal approximations
 - Generalized-gradient approximations
 - Meta-generalized-gradient approximations
- Single-determinant hybrid approximations
 - Hybrid approximations
 - Range-separated hybrid approximations
- Multideterminant hybrid approximations
 - Double-hybrid approximations
 - Range-separated double-hybrid approximations
- Dispersion corrections
 - Semiempirical dispersion corrections
 - Nonlocal van der Waals density functionals
- Machine-learned density functionals
- Benchmark

- ▶ Construction of exchange-correlation energy approximations using Machine Learning (ML) is a current topic of research.
- ▶ For example, such ML exchange-correlation energy approximations can be of the form

$$E_{xc}^{ML} = \int f_{ML}[n, \nabla n, \tau, \dots](\mathbf{r}) d\mathbf{r}$$

where f_{ML} is a very complicated function performing various sequential transformations (including neural networks) of the different input features $(n, \nabla n, \tau, \dots)$ and containing of the order of 10^5 parameters.


- ▶ A very large amount of reliable accurate data is necessary to optimize all the parameters.
- ▶ Example: DM21 (2021), Skala (2025)
- ▶ ML exchange-correlation approximations look very promising, but, as of 2025, they are not yet used in applications.

3 Approximations for the exchange-correlation energy

- Local-density approximation
- Semilocal approximations
 - Generalized-gradient approximations
 - Meta-generalized-gradient approximations
- Single-determinant hybrid approximations
 - Hybrid approximations
 - Range-separated hybrid approximations
- Multideterminant hybrid approximations
 - Double-hybrid approximations
 - Range-separated double-hybrid approximations
- Dispersion corrections
 - Semiempirical dispersion corrections
 - Nonlocal van der Waals density functionals
- Machine-learned density functionals
- Benchmark

Benchmark of exchange-correlation approximations

Mean errors on large sets of molecular energetic data (nearly 5000 data points) for one of the “best” approximation in each family:



GGA = B97-D3(BJ); **mGGA** = B97M-rV; **1H** = M06-2X-D3;
RSH = ω B97M-V; **2DH** = DSD-PBEPBE-D3(BJ); **RSDH** = ω B97M(2)

Data from N. Mardirossian and M. Head-Gordon, Journal of Chemical Physics **148**, 241736 (2018) and N. Mardirossian and M. Head-Gordon, Molecular Physics **115**, 2315 (2017).

- 4 Time-dependent density-functional theory
 - Runge-Gross theorem
 - Linear-response TDDFT

- 4 Time-dependent density-functional theory
 - Runge-Gross theorem
 - Linear-response TDDFT

- ▶ Consider the **time-dependent electronic Schrödinger equation** with an external **time-dependent potential** $\hat{V}(t)$

$$i \frac{\partial |\Psi(t)\rangle}{\partial t} = \left(\hat{T} + \hat{W}_{\text{ee}} + \hat{V}(t) \right) |\Psi(t)\rangle$$

- ▶ Similarly to the Hohenberg-Kohn theorem, **Runge and Gross** (1984) showed that, for a given initial wave function $\Psi(0)$, the time-dependent density $n(\mathbf{r}, t)$ determines the time-dependent potential $v(\mathbf{r}, t)$ up to an arbitrary additive time function:

$$n(\mathbf{r}, t) \xrightarrow{\text{Runge-Gross}} v(\mathbf{r}, t) + c(t)$$

- ▶ We can thus set up a **time-dependent non-interacting KS system**

$$i \frac{\partial \psi_a(\mathbf{r}, t)}{\partial t} = \left(-\frac{1}{2} \nabla^2 + v_s(\mathbf{r}, t) \right) \psi_a(\mathbf{r}, t)$$

where the time-dependent KS potential $v_s(\mathbf{r}, t) = v(\mathbf{r}, t) + v_{\text{Hxc}}(\mathbf{r}, t)$ **reproduces the evolution of the exact density** as $n(\mathbf{r}, t) = \sum_a^N |\psi_a(\mathbf{r}, t)|^2$.

- 4 Time-dependent density-functional theory
 - Runge-Gross theorem
 - Linear-response TDDFT

- ▶ Let us consider a time-periodic potential of frequency ω . In Fourier space, a variation of the KS potential $v_s(\mathbf{r}_1, \omega)$ caused by a variation of the density $n(\mathbf{r}_2, \omega)$ can be written as

$$\frac{\delta v_s(\mathbf{r}_1, \omega)}{\delta n(\mathbf{r}_2, \omega)} = \frac{\delta v(\mathbf{r}_1, \omega)}{\delta n(\mathbf{r}_2, \omega)} + \frac{\delta v_{\text{Hxc}}(\mathbf{r}_1, \omega)}{\delta n(\mathbf{r}_2, \omega)}$$

- ▶ This can be rewritten as

$$\chi_s^{-1}(\mathbf{r}_1, \mathbf{r}_2, \omega) = \chi^{-1}(\mathbf{r}_1, \mathbf{r}_2, \omega) + f_{\text{Hxc}}(\mathbf{r}_1, \mathbf{r}_2, \omega)$$

where

- ▶ $\chi_s(\mathbf{r}_1, \mathbf{r}_2, \omega) = \delta n(\mathbf{r}_1, \omega) / \delta v_s(\mathbf{r}_2, \omega)$ is the **KS non-interacting linear-response function**
 - ▶ $\chi(\mathbf{r}_1, \mathbf{r}_2, \omega) = \delta n(\mathbf{r}_1, \omega) / \delta v(\mathbf{r}_2, \omega)$ is the **interacting linear-response function**
 - ▶ $f_{\text{Hxc}}(\mathbf{r}_1, \mathbf{r}_2, \omega) = \delta v_{\text{Hxc}}(\mathbf{r}_1, \omega) / \delta n(\mathbf{r}_2, \omega)$ is the **Hartree-exchange-correlation kernel**
- ▶ The interacting linear-response function $\chi(\mathbf{r}_1, \mathbf{r}_2, \omega)$ is thus found from the **Dyson-like response equation**

$$\chi^{-1}(\mathbf{r}_1, \mathbf{r}_2, \omega) = \chi_s^{-1}(\mathbf{r}_1, \mathbf{r}_2, \omega) - f_{\text{Hxc}}(\mathbf{r}_1, \mathbf{r}_2, \omega)$$

or, equivalently,

$$\chi(\mathbf{r}_1, \mathbf{r}_2, \omega) = \chi_s(\mathbf{r}_1, \mathbf{r}_2, \omega) + \iint d\mathbf{r}_3 d\mathbf{r}_4 \chi_s(\mathbf{r}_1, \mathbf{r}_3, \omega) f_{\text{Hxc}}(\mathbf{r}_3, \mathbf{r}_4, \omega) \chi(\mathbf{r}_4, \mathbf{r}_2, \omega)$$

- ▶ The KS linear-response function has poles at the KS (de-)excitation energies

$$\chi_s(\mathbf{r}_1, \mathbf{r}_2, \omega) = \sum_{\sigma \in \{\uparrow, \downarrow\}} \sum_a^{\text{occ}} \sum_r^{\text{vir}} \left[\frac{\psi_{a\sigma}^*(\mathbf{r}_1) \psi_{r\sigma}(\mathbf{r}_1) \psi_{r\sigma}^*(\mathbf{r}_2) \psi_{a\sigma}(\mathbf{r}_2)}{\omega - (\varepsilon_r - \varepsilon_a) + i0^+} - \frac{\psi_{r\sigma}^*(\mathbf{r}_1) \psi_{a\sigma}(\mathbf{r}_1) \psi_{a\sigma}^*(\mathbf{r}_2) \psi_{r\sigma}(\mathbf{r}_2)}{\omega + (\varepsilon_r - \varepsilon_a) + i0^+} \right]$$

- ▶ Similarly, $\chi(\mathbf{r}_1, \mathbf{r}_2, \omega)$ has poles at the exact excitation energies $\omega_n = E_n - E_0$.

- ▶ Writing $\chi^{-1}(\omega) = \chi_s^{-1}(\omega) - f_{\text{Hxc}}(\omega)$ in the spin-orbital tensor-product basis $\{\psi_a^* \psi_r, \psi_r^* \psi_a\}$

$$\chi^{-1}(\omega) = - \left[\begin{pmatrix} \mathbf{A}(\omega) & \mathbf{B}(\omega) \\ \mathbf{B}(-\omega)^* & \mathbf{A}(-\omega)^* \end{pmatrix} - \omega \begin{pmatrix} \mathbf{1} & \mathbf{0} \\ \mathbf{0} & -\mathbf{1} \end{pmatrix} \right]$$

where the matrices $\mathbf{A}(\omega)$ and $\mathbf{B}(\omega)$ are

$$[\mathbf{A}(\omega)]_{ar,bs} = (\varepsilon_r - \varepsilon_a) \delta_{ab} \delta_{rs} + \langle rb | f_{\text{Hxc}}(\omega) | as \rangle$$

$$[\mathbf{B}(\omega)]_{ar,bs} = \langle rs | f_{\text{Hxc}}(\omega) | ab \rangle$$

- ▶ The excitation energies ω_n can be calculated from the generalized eigenvalue equation

$$\begin{pmatrix} \mathbf{A}(\omega_n) & \mathbf{B}(\omega_n) \\ \mathbf{B}(-\omega_n)^* & \mathbf{A}(-\omega_n)^* \end{pmatrix} \begin{pmatrix} \mathbf{X}_n \\ \mathbf{Y}_n \end{pmatrix} = \omega_n \begin{pmatrix} \mathbf{1} & \mathbf{0} \\ \mathbf{0} & -\mathbf{1} \end{pmatrix} \begin{pmatrix} \mathbf{X}_n \\ \mathbf{Y}_n \end{pmatrix}$$

The Hartree-exchange-correlation kernel

- ▶ In linear-response TDDFT, the key quantity to be approximated is the **Hartree-exchange-correlation kernel**

$$f_{\text{Hxc}}(\mathbf{r}_1, \mathbf{r}_2, \omega) = \frac{\delta v_{\text{Hxc}}(\mathbf{r}_1, \omega)}{\delta n(\mathbf{r}_2, \omega)}$$

- ▶ It can be decomposed as

$$f_{\text{Hxc}}(\mathbf{r}_1, \mathbf{r}_2, \omega) = f_{\text{H}}(\mathbf{r}_1, \mathbf{r}_2) + f_{\text{xc}}(\mathbf{r}_1, \mathbf{r}_2, \omega)$$

where the Hartree kernel is simply $f_{\text{H}}(\mathbf{r}_1, \mathbf{r}_2) = 1/|\mathbf{r}_1 - \mathbf{r}_2|$.

- ▶ In almost all TDDFT calculations, the frequency dependence of f_{xc} is neglected, which is called the **adiabatic approximation**

$$f_{\text{xc}}(\mathbf{r}_1, \mathbf{r}_2, \omega) \approx \frac{\delta v_{\text{xc}}(\mathbf{r}_1)}{\delta n(\mathbf{r}_2)} = \frac{\delta^2 E_{\text{xc}}[n]}{\delta n(\mathbf{r}_1) \delta n(\mathbf{r}_2)}$$

with the notorious consequence that only single-electron excitations are taken into account (double excitations and higher are missing).

- ▶ To describe nonlocal excitations, such as charge-transfer excitations, range-separated hybrid approximations are often used. The kernel has then the expression

$$f_{\text{xc}} = f_{\text{x}}^{\text{lr, HF}} + f_{\text{x}}^{\text{sr, DFA}} + f_{\text{c}}^{\text{DFA}}$$

where $f_{\text{x}}^{\text{lr, HF}}$ is the long-range HF exchange kernel.