

Relativistic electronic-structure methods based on effective quantum electrodynamics

Julien Toulouse

Laboratoire de Chimie Théorique
Sorbonne Université and **CNRS**, Paris, France
Institut Universitaire de France

27th ETSF Workshop
Marseille, June 2024

- 1 Relativistic electronic-structure theory
- 2 Effective quantum electrodynamics
- 3 1D model for effective quantum electrodynamics

- 1 Relativistic electronic-structure theory
- 2 Effective quantum electrodynamics
- 3 1D model for effective quantum electrodynamics

- ▶ It is known that **special relativity effects** can strongly influence **physical and chemical properties of heavy elements**.

Pyykkö, Annu. Rev. Phys. Chem., 2012

- ▶ **A simple explanation:** the average electron velocity in hydrogen-like atoms

$$\sqrt{\langle \hat{v}^2 \rangle} = \frac{Z}{n}$$

can be a significant fraction of the speed of light $c \approx 137$ a.u. for large Z .


- ▶ **Two striking examples:**

- **Gold would not be yellow without relativity!**

Relativity reduces the $5d \rightarrow 6s$ excitation energy from 3.6 eV (UV) to 2 eV (Vis)

Romaniello, de Boeij, JCP, 2005

- **Lead-acid batteries would not work without relativity!**



Relativity increases the electromotive force from 0.39 V to 2.13 V due to destabilization of PbO_2

Ahuja, Blomqvist, Larsson, Pyykkö, Zaleski-Ejgierd, PRL, 2011

One-electron Dirac equation

- We start with the **one-electron Dirac equation** with a potential v :

$$(\mathbf{D}_0 + v(\vec{r})) \psi_p(\vec{r}) = \varepsilon_p \psi_p(\vec{r})$$

with the 4×4 **free Dirac Hamiltonian**


$$\mathbf{D}_0 = c\vec{\alpha} \cdot \vec{p} + \beta mc^2$$

where $\vec{p} = -i\vec{\nabla}$ is the momentum operator, and $\vec{\alpha} = (\alpha_1, \alpha_2, \alpha_3)$ and β are the 4×4 Dirac matrices

$$\vec{\alpha} = \begin{pmatrix} \mathbf{0}_2 & \vec{\sigma} \\ \vec{\sigma} & \mathbf{0}_2 \end{pmatrix} \text{ and } \beta = \begin{pmatrix} \mathbf{I}_2 & \mathbf{0}_2 \\ \mathbf{0}_2 & -\mathbf{I}_2 \end{pmatrix}$$

and $\vec{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$ are the 2×2 Pauli matrices.

- The (4-component) eigenfunctions $\{\psi_p\}$ can be partitioned into
- **positive-energy states (PS)** $\{\psi_p\}_{p \in \text{PS}} \leftarrow$ **electrons**
 - **negative-energy states (NS)** $\{\psi_p\}_{p \in \text{NS}} \leftarrow$ **positrons**
- Note: What we call “electrons” and “positrons” depend on the potential v !



No-pair many-electron Dirac-Coulomb-Breit (DCB) Hamiltonian

- ▶ In relativistic electronic-structure theory, we normally neglect the negative-energy (positronic) states, which is called the **no-pair approximation**.
- ▶ Thus, we introduce the **Dirac electron field operator** (with only positive-energy states)

$$\hat{\psi}_+(\vec{r}) = \sum_{p \in \text{PS}} \hat{b}_p \psi_p(\vec{r})$$

where $\{\hat{b}_p\}$ are electron annihilation operators.

- ▶ The **no-pair many-electron Dirac-Coulomb-Breit (DCB) Hamiltonian** is

$$\hat{H}^{\text{np}} = \hat{T}^{\text{np}} + \hat{V}^{\text{np}} + \hat{W}^{\text{np}}$$

with the **no-pair** free-electron and external potential operators

$$\hat{T}^{\text{np}} = \int \hat{\psi}_+^\dagger(\vec{r}) \mathbf{D}_0 \hat{\psi}_+(\vec{r}) d\vec{r} \quad \text{and} \quad \hat{V}^{\text{np}} = \int \hat{\psi}_+^\dagger(\vec{r}) v(\vec{r}) \hat{\psi}_+(\vec{r}) d\vec{r}$$

and the **no-pair** Coulomb-Breit two-electron interaction operator

$$\hat{W}^{\text{np}} = \frac{1}{2} \iint \hat{\psi}_+^\dagger(\vec{r}_1) \hat{\psi}_+^\dagger(\vec{r}_2) \mathbf{w}(\vec{r}_{12}) \hat{\psi}_+(\vec{r}_2) \hat{\psi}_+(\vec{r}_1) d\vec{r}_1 d\vec{r}_2$$

where

$$\mathbf{w}(\vec{r}_{12}) = \underbrace{\frac{1}{r_{12}}}_{\text{Coulomb}} - \underbrace{\frac{1}{2r_{12}} \left(\vec{\alpha} \otimes \vec{\alpha} + \frac{(\vec{\alpha} \cdot \vec{r}_{12}) \otimes (\vec{\alpha} \cdot \vec{r}_{12})}{r_{12}^2} \right)}_{\text{Breit}}$$

Beyond the no-pair approximation

- ▶ The next challenge in relativistic electronic-structure theory is to go **beyond the no-pair approximation**, i.e. to include the **quantum-electrodynamics (QED)** effect of **vacuum polarization due to virtual electron-positron pairs**.



- ▶ This is necessary for high accuracy, but also to put relativistic electronic-structure theory on firmer theoretical grounds.
- ▶ How to include QED effects in relativistic electronic-structure theory?
 - Highly accurate QED calculations for very few electron systems
E.g., Indelicato, Mohr, Book chapter, 2016
 - Model one-electron QED correction potentials for many-electron systems
E.g., Sunaga, Salman, Saue, JCP, 2022
- ▶ Here, we explore an **effective QED theory**, i.e. without photons but with the static Coulomb-Breit two-particle interaction.

Chaix, Iracane, JPB, 1989; Saue, Visscher, Book chapter, 2003;
Hainzl, Lewin, Séré, Solovej, PRA, 2007; Kutzelnigg, CP, 2012; Liu, Lindgren, JCP, 2013;
Toulouse, SciPost Chem., 2021.

- 1 Relativistic electronic-structure theory
- 2 Effective quantum electrodynamics
- 3 1D model for effective quantum electrodynamics

Effective quantum electrodynamics (1/2)

- ▶ We start from the **free vacuum state** $|0\rangle$ where all the free negative-energy states are filled (called the “Dirac sea”).
- ▶ The **Dirac field operator** in terms of free electron and positron states $\{\psi_p^0\}$ is then:

$$\hat{\psi}(\vec{r}) = \sum_{p \in \text{PS}} \hat{b}_p \psi_p^0(\vec{r}) + \sum_{p \in \text{NS}} \hat{d}_p^\dagger \psi_p^0(\vec{r})$$


where $\{\hat{b}_p\}$ are electron annihilation operators and $\{\hat{d}_p^\dagger\}$ are positron creation operators such that $\hat{b}_p|0\rangle = \hat{d}_p|0\rangle = 0$.

- ▶ We work in the **electron-positron Fock space** which can be decomposed into charge sectors:

$$\mathcal{F} = \bigoplus_N \mathcal{F}_N$$

where \mathcal{F}_N is the Fock space for N negative charges.

- ▶ In \mathcal{F}_N , the number of electrons N_e and the number of positrons N_p **are not fixed**, but only the net (opposite) charge $N = N_e - N_p$ is fixed.



- The **effective QED Hamiltonian** is defined as

$$\hat{H} = \hat{T} + \hat{V} + \hat{W}$$

with the **normal-ordered** free-particle and external potential operators

$$\hat{T} = \int \mathcal{N}_0 \left[\hat{\psi}^\dagger(\vec{r}) \mathbf{D}_0 \hat{\psi}(\vec{r}) \right] d\vec{r} \quad \text{and} \quad \hat{V} = \int \mathcal{N}_0 \left[\hat{\psi}^\dagger(\vec{r}) v(\vec{r}) \hat{\psi}(\vec{r}) \right] d\vec{r}$$

and the **normal-ordered** Coulomb-Breit two-particle interaction operator

$$\hat{W} = \frac{1}{2} \iint \mathcal{N}_0 \left[\hat{\psi}^\dagger(\vec{r}_1) \hat{\psi}^\dagger(\vec{r}_2) \mathbf{w}(\vec{r}_{12}) \hat{\psi}(\vec{r}_2) \hat{\psi}(\vec{r}_1) \right] d\vec{r}_1 d\vec{r}_2$$

- The **normal ordering** $\mathcal{N}_0[\dots]$ is taken **with respect to the free vacuum state** $|0\rangle$:
- the annihilation electron and positron operators \hat{b}_p and \hat{d}_p are put to the right of the creation electron and positron operators \hat{b}_p^\dagger and \hat{d}_p^\dagger
 - the energy is calculated with respect to the free vacuum state, i.e. $\langle 0 | \hat{H} | 0 \rangle = 0$
 - the Hamiltonian correctly has charge-conjugation symmetry: $\hat{C} \hat{H}[v] \hat{C}^\dagger = \hat{H}[-v]$

Relativistic DFT based on effective QED (1/2)

- ▶ We can formulate a **relativistic density-functional theory (DFT)** based on **effective QED** in a very similar way to non-relativistic DFT.

Toulouse, SciPost Chem., 2021.

- ▶ In effective QED, the **ground-state energy for N negative charges** is

$$E_N = \min_{|\Psi\rangle \in \mathcal{W}_N} \langle \Psi | \hat{T} + \hat{W} + \hat{V} | \Psi \rangle$$

where $\mathcal{W}_N = \{|\Psi\rangle \in \mathcal{F}_N \mid \langle \Psi | \Psi \rangle = 1\}$.

- ▶ Thus, we define the corresponding **Levy-Lieb universal density functional** as

$$F[n] = \min_{\substack{|\Psi\rangle \in \mathcal{W}_N \\ |\Psi\rangle \rightsquigarrow n}} \langle \Psi | \hat{T} + \hat{W} | \Psi \rangle$$

where n is the **opposite charge density** associated with the **normal-ordered density operator** $\hat{n}(\vec{r}) = \mathcal{N}_0 \left[\hat{\psi}^\dagger(\vec{r}) \hat{\psi}(\vec{r}) \right]$.

- ▶ The **exact ground-state energy for N negative charges** can then be expressed as

$$E_N = \min_{n \in \mathcal{D}_N} \left(F[n] + \int v(\vec{r}) n(\vec{r}) d\vec{r} \right)$$

where $\mathcal{D}_N = \{n \mid \exists |\Psi\rangle \in \mathcal{W}_N \text{ s.t. } |\Psi\rangle \rightsquigarrow n\}$ is the set of N -representable densities.

- ▶ We can set up a **Kohn-Sham DFT scheme** by decomposing the universal functional as

$$F[n] = \min_{\substack{|\Phi\rangle \in \mathcal{S}_N \\ |\Phi\rangle \rightsquigarrow n}} \langle \Phi | \hat{T} | \Phi \rangle + E_{\text{Hxc}}[n]$$

where \mathcal{S}_N is the set of all N -electron Slater determinant states and $E_{\text{Hxc}}[n]$ is a Hartree-exchange-correlation density functional.

- ▶ The **exact ground-state energy for N negative charges** can then be expressed as

$$E_N = \min_{|\Phi\rangle \in \mathcal{S}_N} \left(\langle \Phi | \hat{T} + \hat{V} | \Phi \rangle + E_{\text{Hxc}}[n_{|\Phi\rangle}] \right)$$

- ▶ The corresponding **Kohn-Sham equations** are

$$(\mathbf{D}_0 + v(\vec{r}) + v_{\text{Hxc}}(\vec{r})) \psi_p(\vec{r}) = \varepsilon_p \psi_p(\vec{r})$$

where $v_{\text{Hxc}}(\vec{r}) = \delta E_{\text{Hxc}}[n] / \delta n(\vec{r})$ and the density is written as

$$n(\vec{r}) = \sum_{i=1}^N \psi_i^\dagger(\vec{r}) \psi_i(\vec{r}) + n^{\text{vp}}(\vec{r})$$

where $n^{\text{vp}}(\vec{r})$ is the **vacuum-polarization density**

$$n^{\text{vp}}(\vec{r}) = \sum_{p \in \text{NS}} \psi_p^\dagger(\vec{r}) \psi_p(\vec{r}) - \sum_{p \in \text{NS}} \psi_p^{0\dagger}(\vec{r}) \psi_p^0(\vec{r})$$

How to implement effective QED in practice?

- ▶ Effective QED, like standard QED, suffers from **singularities**.
- ▶ In particular, in the complete-basis limit, the **vacuum-polarization density diverges!**
- ▶ In standard QED, this divergence can be regularized with a UV cutoff and the dependence on the cutoff is absorbed into a redefinition of the elementary charge. This is called **charge renormalization**.
- ▶ In effective QED, **charge renormalization** can also be performed in the complete-basis limit.
Hainzl, Lewin, Séré, Solovej, PRA, 2007
- ▶ **But we do not know yet how to handle the situation in a finite basis!**
- ▶ To progress toward a practical implementation in a finite basis, we will now study a **1D effective QED model**.
Audinet, Toulouse, JCP, 2023

- 1 Relativistic electronic-structure theory
- 2 Effective quantum electrodynamics
- 3 1D model for effective quantum electrodynamics

1D hydrogen-like Dirac equation

- ▶ We consider a **1D hydrogen-like Dirac equation** with a **delta potential**:

$$(\mathbf{D}_0 - Z\delta(x))\psi_p(x) = \varepsilon_p\psi_p(x)$$

with the **1D 2×2 free Dirac Hamiltonian**


$$\mathbf{D}_0 = c\sigma_1 p_x + \sigma_3 mc^2$$

where $p_x = -i\hbar d/dx$ is the momentum operator, and σ_1 and σ_3 are the Pauli matrices

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \text{ and } \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

Lapidus, AJP, 1983

- ▶ **The delta potential simulates the Coulomb interaction in 1D.**
Herrick, Stillinger, PRA, 1975
- ▶ **The bound and continuum eigenfunctions** can be calculated **analytically**.
Nogami, Beachey, EL, 1986
- ▶ There are subtleties on the mathematical interpretation of the delta potential.
Audinet, Morellini, Levitt, Toulouse, in preparation




Vacuum-polarization density in a *complete* basis

- ▶ At first order in Z , the **(Uehling) vacuum-polarization density** is

$$n^{\text{vp}}(x) = \frac{Z}{\pi c} \delta(x) + n_{\text{reg}}^{\text{vp}}(x)$$

where the regular part is $n_{\text{reg}}^{\text{vp}}(x) = -\frac{Zm}{\pi} \int_1^\infty \frac{e^{-2mc|x|t}}{t\sqrt{t^2-1}} dt$.

- ▶ The vacuum-polarization density has a **singular delta contribution at the nucleus**, similarly to the 3D case, but with a **finite coefficient**.
- ▶ Vacuum-polarization density $n^{\text{vp}}(x)$ for $Z = c = m = 1$:




Vacuum-polarization density in a *finite* basis

- ▶ We use a **finite plane-wave basis** $\left\{ e^{ikx}, k = \frac{2\pi\mathbb{Z}}{L}, |k| \leq \Lambda \right\}$ on the interval $[-L/2, L/2]$.

- ▶ The vacuum-polarization density is calculated from the eigenfunctions at Z and $Z = 0$:

$$n^{\text{vp}}(x) = \sum_{p \in \text{NS}} \psi_p^{Z\dagger}(x) \psi_p^Z(x) - \sum_{p \in \text{NS}} \psi_p^{0\dagger}(x) \psi_p^0(x)$$

- ▶ Vacuum-polarization density $n^{\text{vp}}(x)$ for $Z = c = m = 1$ in the **finite basis**:




Regularized vacuum-polarization density in a finite basis

- ▶ The vacuum-polarization density calculated in the finite basis is **regularized in Fourier space by removing the large-momentum contributions**:

$$n_{\text{regularized}}^{\text{vp}}(x) = F^{-1} \left[\left(F[n^{\text{vp}}](k) - F[n^{\text{vp}}](k_{\text{max}}) \right) \theta(k_{\text{max}} - k) \right]$$

for some k_{max} depending on the basis.

- ▶ **Regularized** vacuum-polarization density $n^{\text{vp}}(x)$ for $Z = c = m = 1$ in the finite basis:



► **Summary:**

- Effective QED includes electron-positron pairs without photons
- More tractable alternative to standard QED for electronic-structure calculations
- Need to handle singularities in a finite basis for a practical implementation
- Our 1D model suggests that it can be done

► **Outlook:**

- Develop relativistic DFT for the 1D effective QED model
- Extension to real 3D systems

www.lct.jussieu.fr/pagesperso/toulouse