

# Two strategies for continuum excited states: Robin boundary conditions and semiclassical approximations

**Julien Toulouse**

Laboratoire de Chimie Théorique

**Sorbonne Université** and **CNRS**, Paris, France

**Institut Universitaire de France**

Workshop on excited-state methods

April 2023, Toulouse

- 1 Difficulties with continuum states
- 2 Robin boundary conditions
- 3 Semiclassical approximations

- 1 Difficulties with continuum states
- 2 Robin boundary conditions
- 3 Semiclassical approximations

# There is more to electronic-structure theory than bound states!

- ▶ The Coulomb nuclei-electron potential is non-confining, which leads to a **continuous energy spectrum** with “**continuum states**”.
- ▶ Mathematically, **the continuum states are not truly energy eigenstates** since they do not belong to the  $L^2$  Hilbert space.
- ▶ In principle, **pointwise/distribution convergence to continuum states** can be obtained with  $L^2$  basis functions.
- ▶ In practice, **convergence to continuum states** (or to properties crucially depending on continuum states) **can be very slow or impossible with straightforward basis expansions**.



Reinhardt, *Comp. Phys. Comm.*, 1979

# Photoionization cross section

- ▶ The **photoabsorption cross section** is

$$\sigma(\omega) = \lim_{\eta \rightarrow 0^+} \frac{4\pi\omega}{c} \text{Im}[\alpha(\omega + i\eta)]$$

where  $\alpha(\omega)$  is the dynamic dipole polarizability.

- ▶ For  $\omega \geq \text{IP}$ , this is the **photoionization cross section**, which crucially depends on continuum states.
- ▶ If  $\hat{H}$  were diagonalizable in a complete eigenfunction basis  $\{\Psi_n\}_{n \in \mathbb{N}}$ , the cross section could be written as

$$\sigma(\omega) = \frac{4\pi^2\omega}{c} \sum_{n=0}^{\infty} |\langle \Psi_0 | \hat{d} | \Psi_n \rangle|^2 \delta(\omega - (E_n - E_0))$$

where  $\hat{d}$  is the dipole-moment operator (along the direction of the electric field).

- ▶ Formally, it can be expressed as an expectation value

$$\sigma(\omega) = \frac{4\pi^2\omega}{c} \langle \Psi_0 | \hat{d} \delta(\omega + E_0 - \hat{H}) \hat{d} | \Psi_0 \rangle$$

where  $\delta(\omega + E_0 - \hat{H})$  is the spectral-density operator.



- In a **finite basis set**, the linear-response (adiabatic) TDDFT/TDHF equations are

$$\begin{pmatrix} \mathbf{A} & \mathbf{B} \\ -\mathbf{B}^* & -\mathbf{A}^* \end{pmatrix} \begin{pmatrix} \mathbf{X}_n \\ \mathbf{Y}_n \end{pmatrix} = \omega_n \begin{pmatrix} \mathbf{X}_n \\ \mathbf{Y}_n \end{pmatrix}$$

with  $A_{ia,jb} = (\varepsilon_a - \varepsilon_i)\delta_{ij}\delta_{ab} + \langle aj|\hat{f}_{\text{Hxc}}|ib\rangle$  and  $B_{ia,jb} = \langle ab|\hat{f}_{\text{Hxc}}|ij\rangle$ .

- We obtain  $M$  **excitation energies**  $\omega_n$  and associated **oscillator strengths**  $f_n$ . The photoabsorption cross section is then

$$\sigma(\omega) = \frac{2\pi^2}{c} \sum_{n=1}^M f_n \delta(\omega - \omega_n)$$

- For  $\omega \geq \text{IP}$ , the **photoionization cross section** at  $\omega = \omega_n$  may be approximated as

$$\sigma(\omega_n) \approx \frac{2\pi^2}{c} f_n \rho_{\text{DOS}}(\omega_n)$$

where  $\rho_{\text{DOS}}(\omega_n)$  is a finite-difference estimate of the **density of states** at  $\omega = \omega_n$

$$\rho_{\text{DOS}}(\omega_n) = \frac{dn}{d\omega_n} \approx \frac{2}{\omega_{n+1} - \omega_{n-1}}$$



Macías, Martín, Riera, Yáñez, *Int. J. Quantum Chem.*, 1988

Yang, van Faassen, Burke, *J. Chem. Phys.*, 2009

Zapata, Luppi, Toulouse, *J. Chem. Phys.*, 2019

# Photoionization cross sections of He and Be

- ▶ We use a **B-spline basis set**, i.e. localized piecewise polynomial functions in a spherical box  $\Omega$ , with Dirichlet (zero) boundary conditions.
- ▶ TDHF photoionization cross sections of the He and Be atoms:



⇒ Reasonable spectrum for He but it does not work for core ionization in Be

Zapata, Luppi, Toulouse, J. Chem. Phys., 2019

# So, how to deal with continuum states?

## Some approaches for calculating cross sections involving continuum states:

- ▶ Various techniques involving the **complex-frequency plane**

E.g., analytical continuation of  $\alpha(\omega)$ :

Tenorio, Coriani, Rocha, Nascimento, *Prog. Theor. Chem. Phys.*, 2021

- ▶ Imposing **boundary conditions adapted to continuum states**

E.g., in TDDFT:

Stener, Decleva, Lisini, *J. Phys. B*, 1995

⇒ **Here, I will talk about using Robin boundary conditions**

- ▶ **Semiclassical approximations**

E.g., in nuclear physics:

Schuck, Hasse, Jaenicke, Grégoire, Rémaud, Sébille, Suraud, *Prog. Part. Nucl. Phys.*, 1989

⇒ **Here, I will talk about semiclassical approximations for photoionization**



1 Difficulties with continuum states

2 Robin boundary conditions

with E. Cancès, A. Levitt, E. Luppi, K. Schwinn, F. Zapata

3 Semiclassical approximations

- ▶ Let us look at a **basis-independent** formulation of linear-response TDDFT/TDHF.
- ▶ In (adiabatic) TDDFT/TDHF, the **occupied orbitals evolve in time** according to

$$i \frac{\partial}{\partial t} \psi_i(t) = \hat{h}[\gamma(t)] \psi_i(t) + \hat{v}(t) \psi_i(t)$$

where  $\hat{h}[\gamma(t)]$  is the KS/HF Hamiltonian depending on the density matrix  $\gamma(t)$  and  $\hat{v}(t) = -\hat{d}\mathcal{E}(e^{-i\omega t} + e^{+i\omega t})$  is the electric-dipole interaction.

- ▶ At **first order in the electric field**, the perturbed occupied orbitals (in the interaction picture) are

$$\psi_i^{(1)}(t) = \psi_i^{(+)}(\omega) e^{-i\omega t} + \psi_i^{(-)}(\omega) e^{+i\omega t}$$

where the Fourier modes  $\psi_i^{(\pm)}(\omega)$  satisfy the **TDDFT/TDHF Sternheimer equations**

$$\left( \pm\omega + \varepsilon_i - \hat{h}[\gamma^{(0)}] \right) \psi_i^{(\pm)}(\omega) = \left( \hat{v}_{\text{Hxc}}^{(\pm)}(\omega) - \hat{d} \right) \psi_i^{(0)}$$

- ▶ The **dynamic dipole polarizability** is then obtained as

$$\alpha(\omega) = \sum_{i=1}^N \langle \psi_i^{(0)} | \hat{d} | \psi_i^{(+)}(\omega) \rangle + \langle \psi_i^{(-)}(\omega) | \hat{d} | \psi_i^{(0)} \rangle$$

# Tackling the continuum with Robin boundary conditions

- ▶ For  $\omega \geq -\varepsilon_i$ , the first-order perturbed occupied orbital  $\psi_i^{(+)}(\omega)$  is a “**continuum wave function**” (i.e., not belonging to the Hilbert space).
- ▶ **Key idea: Expand  $\psi_i^{(+)}(\omega)$  in a basis set only in a box  $\Omega$  and use an analytical asymptotic approximation  $\bar{\psi}_i^{(+)}(\omega)$  to it outside the box. Match the two functions on the box surface  $\partial\Omega$  by equating the normal logarithmic derivatives.**
- ▶ For spherical symmetry, this leads to **local Robin boundary conditions** for  $\psi_i^{(+)}(\omega)$

$$\forall \mathbf{r} \in \partial\Omega, \frac{\mathbf{n}(\mathbf{r}) \cdot \nabla \psi_i^{(+)}(\mathbf{r}, \omega)}{\psi_i^{(+)}(\mathbf{r}, \omega)} = \frac{\mathbf{n}(\mathbf{r}) \cdot \nabla \bar{\psi}_i^{(+)}(\mathbf{r}, \omega)}{\bar{\psi}_i^{(+)}(\mathbf{r}, \omega)}$$

where  $\bar{\psi}_i^{(+)}(\mathbf{r}, \omega)$  are chosen as hydrogen-like continuum wave functions.

- ▶ For non-spherical symmetry, it is extended to **nonlocal Robin boundary conditions**

$$\forall \mathbf{r} \in \partial\Omega, \mathbf{n}(\mathbf{r}) \cdot \nabla \psi_i^{(+)}(\mathbf{r}, \omega) = \int_{\partial\Omega} \bar{K}_i(\mathbf{r}, \mathbf{r}'; \omega) \psi_i^{(+)}(\mathbf{r}', \omega) d\mathbf{r}'$$


where the kernel  $\bar{K}_i(\mathbf{r}, \mathbf{r}'; \omega)$  contains information about  $\bar{\psi}_i^{(+)}(\omega)$ .

- ▶ Using now a basis set in the box, it amounts to using the **kinetic integrals**

$$t_{i,\mu,\nu}(\omega) = \frac{1}{2} \int_{\Omega} \nabla \chi_{\mu}^*(\mathbf{r}) \cdot \nabla \chi_{\nu}(\mathbf{r}) d\mathbf{r} - \frac{1}{2} \int_{\partial\Omega^2} \chi_{\mu}^*(\mathbf{r}) \bar{K}_i(\mathbf{r}, \mathbf{r}'; \omega) \chi_{\nu}(\mathbf{r}') d\mathbf{r} d\mathbf{r}'$$

# Photoionization cross section of Be



- ▶ We use a **B-spline basis set**, i.e. localized piecewise polynomial functions in a spherical box  $\Omega$ , **with Robin boundary conditions**.
- ▶ TDLDA and TDHF photoionization cross sections of the Be atom:



⇒ We can now easily converge the spectra for all frequencies

# Core resonances in the Be atom


- ▶ The  $1s \rightarrow np$  core excitations are embedded in the continuum of the valence excitations from the  $2s \Rightarrow$  they are **resonances**, i.e. **quasi-bound states with finite lifetimes**.
- ▶ Example of the  $1s \rightarrow 2p$  core resonance (resonance energy  $E_R$  and inverse lifetime  $\Gamma$ ):



	$E_R$ (eV)	$\Gamma$ (meV)
TDLDA	103.0	2.3
TDHF	118.3	0.2
TDLRSH	114.8	0.1
Reference	115.5	37

$\Rightarrow$  **Good resonance energy with TDLRSH but much too small inverse lifetime**

- ▶ The resonance decays via the **Auger process**



The configuration  $1s^22p$  is a double excitation with respect to the ground state  $1s^22s^2$  and thus cannot be described by adiabatic TDDFT/TDHF.

## ▶ **Summary:**

- ▶ TDDFT/TDHF with special boundary conditions for describing continuum states
- ▶ It allows for calculations of photoionization spectra of atoms, including core resonances
- ▶ TDLRSH gives good resonance energies but unreliable lifetimes

## ▶ **Outlook:**

- ▶ Extension to Gaussian basis sets
- ▶ Extension to molecules
- ▶ Extension to time propagation for nonlinear optical properties

- 1 Difficulties with continuum states
- 2 Robin boundary conditions
- 3 Semiclassical approximations**  
with P. Schuck

- ▶ We start from the **photoionization cross section in the velocity gauge**

$$\sigma(\omega) = \frac{4\pi^2}{c\omega} \langle \Psi_0 | \hat{P} \delta(\omega + E_0 - \hat{H}) \hat{P} | \Psi_0 \rangle$$

where  $\hat{P}$  is the total momentum operator (along the direction of the field).

- ▶ We introduce the operators

$$\hat{A} = \delta(\omega + E_0 - \hat{H}) \quad \text{and} \quad \hat{B} = \hat{P} \hat{A} \hat{P}$$

and the ground-state density matrix

$$\hat{\rho}_0 = |\Psi_0\rangle\langle\Psi_0|$$

- ▶ We arrive at the following expression for the **photoionization cross section**

$$\sigma(\omega) = \frac{4\pi^2}{c\omega} \text{Tr}[\hat{B} \hat{\rho}_0] = \frac{4\pi^2}{c\omega} \int_{\mathbb{R}^{6N}} d\mathbf{r} d\mathbf{r}' B(\mathbf{r}, \mathbf{r}') \rho_0(\mathbf{r}', \mathbf{r})$$

where  $B(\mathbf{r}, \mathbf{r}') = \langle \mathbf{r} | \hat{B} | \mathbf{r}' \rangle$  and  $\rho_0(\mathbf{r}', \mathbf{r}) = \langle \mathbf{r}' | \hat{\rho}_0 | \mathbf{r} \rangle$ .



# Wigner representation of the photoionization cross section

- ▶ Let us introduce the **Wigner transform** of an operator  $\hat{C}$

$$[\hat{C}]_W(\mathbf{q}, \mathbf{p}) \equiv C_W(\mathbf{q}, \mathbf{p}) = \int_{\mathbb{R}^{3N}} d\mathbf{s} e^{-i\mathbf{p}\cdot\mathbf{s}} \underbrace{\langle \mathbf{q} + \mathbf{s}/2 |}_{=\mathbf{r}} \hat{C} \underbrace{| \mathbf{q} - \mathbf{s}/2 \rangle}_{=\mathbf{r}'}$$

where  $\mathbf{q} = (\mathbf{r} + \mathbf{r}')/2 \in \mathbb{R}^{3N}$  is the average position vector,  $\mathbf{s} = \mathbf{r} - \mathbf{r}' \in \mathbb{R}^{3N}$  is the relative position vector,  $\mathbf{p} \in \mathbb{R}^{3N}$  is the conjugate momentum vector of  $\mathbf{s}$ .

Ring, Schuck, *The Nuclear Many-Body Problem*, Springer, 2004

Case, *Am. J. Phys.*, 2008

- ▶ The **Wigner transformation preserves the trace** of a product of operators, so we have

$$\sigma(\omega) = \frac{4\pi^2}{c\omega} \int_{\mathbb{R}^{6N}} \frac{d\mathbf{q}d\mathbf{p}}{(2\pi)^{3N}} B_W(\mathbf{q}, \mathbf{p}) \rho_{0,W}(\mathbf{q}, \mathbf{p})$$

- ▶ We have put the photoionization cross section in the form of a **phase-space integral**.
- ▶ So far, everything is exact. We will assume that we know the Wigner function of the ground state  $\rho_{0,W}(\mathbf{q}, \mathbf{p})$ , and we will now use a **semiclassical expansion approximation** for  $B_W(\mathbf{q}, \mathbf{p})$ .

# Semiclassical expansion

- ▶ The **Wigner transform of a product of operators** is given by the Groenewold/Moyal/star-product formula:

$$[\hat{C}\hat{D}]_W(\mathbf{q}, \mathbf{p}) = C_W(\mathbf{q}, \mathbf{p}) e^{(i\hbar/2)\overleftrightarrow{\Lambda}} D_W(\mathbf{q}, \mathbf{p})$$

where  $\overleftrightarrow{\Lambda} = \overleftarrow{\nabla}_{\mathbf{q}} \cdot \overrightarrow{\nabla}_{\mathbf{p}} - \overleftarrow{\nabla}_{\mathbf{p}} \cdot \overrightarrow{\nabla}_{\mathbf{q}}$  is the Poisson bracket differential operator.

- ▶ Using this formula, we find the Wigner transform of  $\hat{B} = \hat{P}\hat{A}\hat{P}$

$$B_W(\mathbf{q}, \mathbf{p}) = \frac{1}{3} \mathbf{P}^2 A_W(\mathbf{q}, \mathbf{p}) + \frac{\hbar^2}{12} \mathbf{D}^2 A_W(\mathbf{q}, \mathbf{p})$$

where  $\mathbf{P} = \sum_{i=1}^N \mathbf{p}_i$  and  $\mathbf{D} = \sum_{i=1}^N \nabla_{\mathbf{q}_i}$ .

- ▶ We also find the semiclassical expansion of the Wigner transform of  $\hat{A} = \delta(\omega + E_0 - \hat{H})$

$$A_W(\mathbf{q}, \mathbf{p}) = A_W^{(0)}(\mathbf{q}, \mathbf{p}) + \hbar^2 A_W^{(2)}(\mathbf{q}, \mathbf{p}) + O(\hbar^4)$$

where  $A_W^{(0)}(\mathbf{q}, \mathbf{p}) = \delta(\omega + E_0 - H(\mathbf{q}, \mathbf{p}))$  and  $H(\mathbf{q}, \mathbf{p}) = \frac{\mathbf{p}^2}{2} + V(\mathbf{q})$

$$A_W^{(2)}(\mathbf{q}, \mathbf{p}) = \frac{1}{8} \left[ -\nabla_{\mathbf{q}}^2 V(\mathbf{q}) \delta''(\omega + E_0 - H(\mathbf{q}, \mathbf{p})) + \frac{1}{3} \left( (\nabla_{\mathbf{q}} V(\mathbf{q}))^2 + (\mathbf{p} \cdot \nabla_{\mathbf{q}})^2 V(\mathbf{q}) \right) \delta'''(\omega + E_0 - H(\mathbf{q}, \mathbf{p})) \right]$$

# Semiclassical expansion of the photoionization cross section

- ▶ We obtain the **semiclassical expansion of the photoionization cross section** (for  $\hbar = 1$ ):

$$\sigma(\omega) = \sigma^{(0)}(\omega) + \sigma^{(2)}(\omega) + \dots$$

- ▶ The zeroth-order term is

$$\sigma^{(0)}(\omega) = \frac{4\pi^2}{3c\omega} \int_{\mathbb{R}^{6N}} \frac{d\mathbf{q}d\mathbf{p}}{(2\pi)^{3N}} \mathbf{P}^2 A_W^{(0)}(\mathbf{q}, \mathbf{p}) \rho_{0,W}(\mathbf{q}, \mathbf{p})$$


- ▶ The second-order term is

$$\sigma^{(2)}(\omega) = \frac{4\pi^2}{3c\omega} \int_{\mathbb{R}^{6N}} \frac{d\mathbf{q}d\mathbf{p}}{(2\pi)^{3N}} \left[ \mathbf{P}^2 A_W^{(2)}(\mathbf{q}, \mathbf{p}) + \frac{1}{4} \mathbf{D}^2 A_W^{(0)}(\mathbf{q}, \mathbf{p}) \right] \rho_{0,W}(\mathbf{q}, \mathbf{p})$$

- ▶ We have arrived at an approximation to the photoionization cross section that only requires to know the ground-state Wigner function  $\rho_{0,W}(\mathbf{q}, \mathbf{p})$  but does not require the calculation of the continuum states.
- ▶ Note that it is not a full expansion in powers of  $\hbar$  since we do not expand  $\rho_{0,W}(\mathbf{q}, \mathbf{p})$ .

# Photoionization cross section of the hydrogen atom


Calculation of  $\sigma^{(0)}(\omega)$  and  $\sigma^{(2)}(\omega)$  by numerical integration:



⇒ As expected, the semiclassical expansion correctly captures the high-energy part of the spectrum

# Photoionization cross section of the helium atom

Calculation of  $\sigma^{(0)}(\omega)$  and an approximation to  $\sigma^{(2)}(\omega)$  by numerical integration:



$\Rightarrow$  Again, the semiclassical expansion correctly captures the high-energy part of the spectrum

## ▶ Summary:

- ▶ We derived semiclassical approximations for photoionization cross sections
- ▶ Tests on atoms confirm that they correctly captures the high-energy part of the spectrum

## ▶ Outlook:

- ▶ Extension to linear-response TDHF/TDDFT
- ▶ Extension to many-body calculations with Monte Carlo integration
- ▶ Extension to other properties such as second-order correlation energy
- ▶ Development of hybrid methods: basis set for low-energy part + semiclassical approximations for high-energy part

[www.lct.jussieu.fr/pagesperso/toulouse/presentations/presentation\\_toulouse\\_23.pdf](http://www.lct.jussieu.fr/pagesperso/toulouse/presentations/presentation_toulouse_23.pdf)